

BRUFEN® PAIN

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I taking BRUFEN PAIN?

BRUFEN PAIN contains the active ingredient ibuprofen. BRUFEN PAIN is used for the temporary relief of pain and/or inflammation associated with headache, migraine headache, tension headache, sinus pain, toothache, dental procedures, backache, muscular aches and pains, period pain, sore throat, tennis elbow, arthritis, rheumatic pain, and the aches and pains associated with colds and flu. It also reduces fever.

For more information, see Section [1. Why am I taking BRUFEN PAIN?](#) in the full CMI.

2. What should I know before I take BRUFEN PAIN?

Do not use if you have allergy to ibuprofen, aspirin, NSAIDs or any of the ingredients listed at the end of the CMI.

Talk to your doctor or pharmacist if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I take BRUFEN PAIN?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with BRUFEN PAIN and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I take BRUFEN PAIN?

- Adults and Children from 12 years: One tablet every 4 to 6 hours as necessary. Do not take more than 3 tablets in 24 hours.
- Do not give to children under 12.
- Do not exceed recommended dose. Excessive use can be harmful and increase the risk of heart attack, stroke or liver damage.

More instructions can be found in Section [4. How do I take BRUFEN PAIN?](#) in the full CMI.

5. What should I know while taking BRUFEN PAIN?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using BRUFEN PAIN.• If you become pregnant while taking this medicine tell your doctor immediately.• If you are going to have surgery, make sure the surgeon and anaesthetist knows you are taking BRUFEN PAIN. NSAID medicines can slow down blood clotting.
Things you should not do	<ul style="list-style-type: none">• Do not take BRUFEN PAIN with any other medicines containing ibuprofen, aspirin or other anti-inflammatory medicines unless your doctor tells you to.• Do not take more than the recommended dose unless your pharmacist or doctor tells you to.

Driving or using machines	<ul style="list-style-type: none"> • Be careful before you drive or use any machines or tools until you know how BRUFEN PAIN affects you. BRUFEN PAIN may cause dizziness in some people. If this happens, do not drive or operate machinery.
Drinking alcohol	<ul style="list-style-type: none"> • Tell your doctor or pharmacist if you drink alcohol.
Looking after your medicine	<ul style="list-style-type: none"> • Keep your medicine in a cool dry place where the temperature stays below 25°C.

For more information, see Section [5. What should I know while taking BRUFEN PAIN?](#) in the full CMI.

6. Are there any side effects?

- Side effects are uncommon. The most common of these include headache, nausea, abdominal pain and heartburn.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

BRUFEN® PAIN

Active ingredient(s): *Ibuprofen*

Consumer Medicine Information (CMI)

This leaflet provides important information about using BRUFEN PAIN. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using BRUFEN PAIN.**

Where to find information in this leaflet:

- [1. Why am I taking BRUFEN PAIN?](#)
- [2. What should I know before I take BRUFEN PAIN?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I take BRUFEN PAIN?](#)
- [5. What should I know while taking BRUFEN PAIN?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I taking BRUFEN PAIN?

BRUFEN PAIN contains the active ingredient ibuprofen. BRUFEN PAIN belongs to a group of medicines called non-steroidal anti-inflammatory drugs (or NSAIDs).

BRUFEN PAIN is used for the temporary relief of pain and/or inflammation associated with headache, migraine headache, tension headache, sinus pain, toothache, dental procedures, backache, muscular aches and pains, period pain, sore throat, tennis elbow, arthritis, rheumatic pain, and the aches and pains associated with colds and flu. It also reduces fever.

2. What should I know before I take BRUFEN PAIN?

Warnings

Do not use BRUFEN PAIN if:

- you are allergic to ibuprofen, aspirin, NSAIDs or any of the ingredients listed at the end of this leaflet.
Many medicines used to treat headache, period pain and other aches and pains contain aspirin or NSAID medicines. If you are not sure if you are taking any of these medicines, ask your pharmacist.
Always check the ingredients to make sure you can use this medicine.
Some of the symptoms of an allergic reaction may include:
 - shortness of breath;
 - wheezing or difficulty breathing;
 - swelling of the face, lips, tongue or other parts of the body; and rash, itching or hives on skin.
- you have asthma that is sensitive to aspirin or NSAIDs

- you have a peptic ulcer (ie stomach or duodenal ulcer), or a recent history of one, or have had peptic ulcers before;
- you have recently (or have previously) vomited blood or material that looks like coffee grounds
- you have recently (or previously) bled from the back passage (rectum), had black sticky bowel motions (stools) or bloody diarrhoea
- you have a condition resulting in an increased tendency to bleed
- you have or have had a history of, Ulcerative Colitis or Crohn's Disease
- you have severe kidney disease
- you have severe heart failure
- you have severe liver disease
- you are being treated for pain following heart bypass surgery

Check with your doctor or pharmacist if you:

- have any other medical conditions
- take any medicines for any other condition
- have asthma
- have heart disease or high blood pressure
- have heartburn, indigestion, stomach ulcer or any other stomach problems
- are vomiting blood or bleeding from the back passage
- have severe skin reactions such as Stevens-Johnson syndrome
- have vision problems

- have liver or kidney disease
- have a tendency to bleed or other blood problems
- have bowel or intestinal problems such as ulcerative colitis
- have heart failure
- have swelling of ankles or feet
- have diarrhoea
- If you are over 65 years of age, talk to your pharmacist or doctor about taking BRUFEN PAIN. Taking this medicine may increase the risk of you getting unwanted effects, such as stomach or heart problems.
- Tell your doctor or pharmacist if you currently have an infection.
- If you take BRUFEN PAIN while you have an infection, it may hide some of the signs and symptoms of an infection. This may make you think, **mistakenly, that you are better or that it is not serious.**
- **Tell your doctor or pharmacist if you plan to have surgery.**
- During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

BRUFEN PAIN is only available from your pharmacist.

BRUFEN PAIN is not addictive.

Pregnancy and breastfeeding

Do not take BRUFEN PAIN if you are pregnant, or intend to become pregnant or during the first 6 months of pregnancy, except on doctor's advice. Do not use at all in the last three months of pregnancy.

It may affect your developing baby if you take it during pregnancy. It may also impair female fertility.

Talk to your doctor or pharmacist if you are breastfeeding or intend to breastfeed. Like most NSAID medicines, BRUFEN PAIN is not recommended while you are breast-feeding.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and ibuprofen may interfere with each other. These include:

- warfarin or clopidogrel, medicines used to stop blood clots
- lithium, a medicine used to treat mood swings and some types of depression
- SSRIs such as sertraline, medicines used to treat depression
- medicines used to treat high blood pressure or other heart conditions

- methotrexate, a medicine used to treat arthritis and some types of cancer
- medicines used to treat heart failure such as digoxin
- medicines such as prednisone, prednisolone and cortisone, which reduce the activity of your immune system
- cyclosporine or tacrolimus, medicines used to treat certain problems with the immune system or to help prevent organ transplant rejection
- aminoglycosides, medicines used to treat certain infections
- quinolone antibiotics, medicines used to treat certain infections
- mifepristone, a medicine used to bring about an abortion
- zidovudine, a medicine used to treat HIV infection
- aspirin, salicylates and other non-steroidal anti-inflammatory drugs (NSAIDs)
- medicines used to treat diabetes
- probenecid, a medicine used to treat gout
- phenytoin, a medicine used to treat epilepsy.

These medicines may be affected by BRUFEN PAIN, or may affect how well it works. You may need to take different amounts of your medicine, or you may need to take different medicines. Your doctor or pharmacist will advise you.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect BRUFEN PAIN.

4. How do I take BRUFEN PAIN?

How much to take

- Adults and Children from 12 years: One tablet every 4 to 6 hours as necessary.
- Do not take more than 3 tablets in 24 hours.
- Do not give to children under 12.
- Follow the instructions provided with the medicine.
- Do not exceed the recommended dosage.
- Excessive use can be harmful and increase the risk of heart attack, stroke and liver damage.

When to take BRUFEN PAIN

- Take BRUFEN PAIN by mouth with water.
- BRUFEN PAIN should not be used for more than 3 days at a time, except on medical advice.

If you take too much BRUFEN PAIN

If you think that you have taken too much BRUFEN PAIN, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**Australia telephone 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while taking BRUFEN PAIN?

Things you should do

- If you become pregnant while taking this medicine tell your doctor immediately.
- If you are going to have surgery, make sure the surgeon and anaesthetist know you are taking BRUFEN PAIN.

NSAID medicines can slow down blood clotting.

- Tell your pharmacist or doctor if your symptoms do not improve. Your pharmacist or doctor will assess your conditions and decide if you should continue to take the medicine.

Remind any doctor, dentist or pharmacist you visit that you are using BRUFEN PAIN.

Things you should not do

- Do not take BRUFEN PAIN with any other medicines containing ibuprofen, aspirin or other anti-inflammatory medicines unless your doctor tells you to.
- Do not take BRUFEN PAIN to treat any other complaints unless your pharmacist or doctor tells you to.

- Do not give your medicine to anyone else, even if they have the same condition as you.
- Do not take more than the recommended dose unless your pharmacist or doctor tells you to.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how BRUFEN PAIN affects you.

BRUFEN PAIN may cause dizziness in some people. If this happens, do not drive or operate machinery.

Drinking alcohol

Tell your doctor or pharmacist if you drink alcohol.

Looking after your medicine

- Keep your medicine in a cool dry place where the temperature stays below 25°C.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions.

Less serious side effects

Less serious side effects	What to do
<p>Stomach related:</p> <ul style="list-style-type: none">• nausea or vomiting• cramps, wind, constipation or diarrhoea• heartburn, or pain in the upper part of your stomach• loss of appetite <p>Head related:</p>	<p>Speak to your doctor or pharmacist if you have any of these less serious side effects and they worry you.</p>

Less serious side effects	What to do
<ul style="list-style-type: none"> • buzzing or ringing in the ears or other trouble hearing • headache • sleeplessness, • changes in mood, for example depression, confusion, nervousness • dizziness • sleepiness <p>Skin related:</p> <ul style="list-style-type: none"> • a rash that always appears in the exact same spot on your skin (fixed eruption) 	

Serious side effects

Serious side effects	What to do
<p>Blood related:</p> <ul style="list-style-type: none"> • vomiting blood or material that looks like coffee grounds • bleeding from the back passage, black sticky bowel motions (stools) or bloody diarrhoea 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Serious side effects	What to do
<ul style="list-style-type: none"> • blood disorders may occur with this medicine. • bleeding or bruising more easily than normal, reddish or purplish blotches under the skin • signs of anaemia, such as tiredness, headaches, being short of breath and looking pale <p>Skin related:</p> <ul style="list-style-type: none"> • severe blisters and bleeding in the lips, eyes, mouth, nose and genitals (Stevens Johnson Syndrome) • symptoms of sunburn (such as redness, itching swelling, blistering) which may occur more quickly than usual <p>Heart related:</p> <ul style="list-style-type: none"> • This medicine may be associated with a small increased risk of heart attack (myocardial infarction) 	

Serious side effects	What to do
<ul style="list-style-type: none"> • fast or irregular heartbeats, also called palpitations <p>Stomach related:</p> <ul style="list-style-type: none"> • severe pain or tenderness in the stomach • nausea, stomach ache <p>Head related:</p> <ul style="list-style-type: none"> • severe or persistent headache <p>Immune system related:</p> <ul style="list-style-type: none"> • headache and stiff neck • fever, generally feeling unwell, • signs of frequent or worrying infections such as fever, severe chills, sore throat or mouth ulcers <p>Eye related:</p> <ul style="list-style-type: none"> • eye problems such as blurred vision, sore red eyes, itching <p>Liver related:</p>	

Serious side effects	What to do
<ul style="list-style-type: none"> • yellowing of the skin and/or eyes, also called jaundice <p>Kidney related:</p> <ul style="list-style-type: none"> • unusual weight gain, swelling of ankles or legs • fluid retention • tingling of the hands and feet <p>Allergy related:</p> <ul style="list-style-type: none"> • asthma, wheezing, shortness of breath or difficulty breathing • swelling of the face, lips or tongue which may cause difficulty in swallowing or breathing • hives, itching or skin rash • stomach ache, fever, chills, nausea and vomiting • fainting <p>If you are allergic to aspirin or NSAID medicines and take BRUFEN PAIN,</p>	

Serious side effects	What to do
these allergy related symptoms may be severe.	

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

7. Product details

This medicine is available over-the-counter without a doctor's prescription.

What BRUFEN PAIN contains

Active ingredient (main ingredient)	Ibuprofen
Other ingredients	Microcrystalline cellulose

(inactive ingredients)	Povidone Croscarmellose sodium Sodium lauryl sulfate Colloidal anhydrous silica Stearic acid Macrogol 6000 Hypromellose Titanium dioxide Purified talc Lactose monohydrate
Potential allergens	Lactose

Do not take this medicine if you are allergic to any of these ingredients.

What BRUFEN PAIN looks like

BRUFEN PAIN tablets (AUST R 401754) are white to off- white, pillow-shaped, film coated tablets, plain on both sides.

Who distributes BRUFEN PAIN

Viatrix Pty Ltd

Level 1, 30 The Bond

30-34 Hickson Road

Millers Point NSW 2000

www.viatris.com.au

Phone: 1800 274 276

This leaflet was prepared in April 2025.

BRUFEN® is a Viatris company trade mark

BRUFEN PAIN_cmi\Apr25/00