

▼ This vaccine is subject to additional monitoring **in Australia**. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse events at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

# AUSTRALIAN PRODUCT INFORMATION – COMIRNATY® Original/Omicron BA.4-5 COVID-19 VACCINE

## 1. NAME OF THE MEDICINE

Tozinameran and Famtozinameran

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Tozinameran and famtozinameran are single-stranded, 5'-capped messenger RNA (mRNA) produced using a cell-free *in vitro* transcription from the corresponding DNA templates, encoding the viral spike (S) protein of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (Original and Omicron BA.4-5).

COMIRNATY Original/Omicron BA.4-5			
Age group	12 years and older	5 to <12 years	6 months to <5 years
Strength per dose	15/15 micrograms	5/5 micrograms	1.5/1.5 micrograms

Each dose contains COVID-19 mRNA Vaccine embedded in lipid nanoparticles.  
For the full list of excipients, see Section 6.1 List of excipients.

## 3. PHARMACEUTICAL FORM

Age group	12 years and older		5 to <12 years			6 months to <5 years
AUST R	413718	400874	412350	413720	413719	417266
Cap & Label colour code	Light Grey	Dark Grey	Orange	Light Blue	Dark Blue	Maroon
Pharmaceutical form	Suspension for injection		Concentrate for suspension for injection	Suspension for injection		Concentrate for suspension for injection
Strength per dose	15/15 micrograms (0.3 mL dose)		5/5 micrograms (0.2 mL dose)	5/5 micrograms (0.3 mL dose)		1.5/1.5 micrograms (0.2 mL dose)
Fill volume	0.48 mL	2.25 mL	1.3 mL	0.48 mL	2.25 mL	0.4 mL
No. of doses per vial	1	6	10	1	6	10
Dilution	Do not dilute		Requires dilution	Do not dilute		Requires dilution

COMIRNATY Original/Omicron BA.4-5 (Grey, Orange and Maroon cap) is a white to off-white frozen suspension.

COMIRNATY Original/Omicron BA.4-5 (Blue cap) is a clear to slightly opalescent solution.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

COMIRNATY Original/Omicron BA.4-5 Vaccine has **provisional approval** for the indication below:

Active immunisation to prevent coronavirus disease 2019 (COVID-19) caused by SARS-CoV-2, in individuals 6 months of age and older.

The use of this vaccine should be in accordance with official recommendations.

The decision has been made on the basis of short term immunogenicity and safety data. Continued approval depends on the evidence of longer term efficacy and safety from ongoing clinical trials and post-market assessment.

### 4.2 Dose and method of administration

#### Dosage

Strength and Age of Individual	Cap and Label Color	Volume of Each Dose	Dose Schedule for Primary Series and Additional dose(s)
<b>1.5/1.5 micrograms per dose</b> 6 months to <5 years	Maroon	0.2 mL	<b>Primary series:</b> 3 doses Dose 1 and 2: at least 3 weeks apart Dose 3: at least 8 weeks after second dose. <b>Additional dose(s):</b> at least 3 months after a previous dose
<b>5/5 micrograms per dose</b> 5 to <12 years	Orange	0.2 mL	<b>Primary series:</b> 2 doses at least to 21 days (preferably 3 weeks) apart <b>Additional dose(s):</b> at least 3 months after a previous dose
	Blue	0.3 mL	
<b>15/15 micrograms per dose</b> 12 years and older	Grey	0.3 mL	<b>Additional dose(s):</b> at least 3 months after a previous dose

Primary series, when clinically indicated, can be given to the individuals such as those who are vaccine-naïve and immunocompromised.

The use of this vaccine should be in accordance with clinical recommendations in Australia, made by ATAGI in the Australian Immunisation Handbook.

#### *Severely immunocompromised aged 12 years and older*

In accordance with official recommendations, a third dose may be given, as part of the primary series, at least 28 days after the second dose to individuals who are severely immunocompromised (see Section 4.4 Special warnings and precautions for use).

#### *Elderly population*

No dosage adjustment is required in elderly individuals  $\geq 65$  years of age.

#### *Paediatric population*

Children who will turn from 4 years to 5 years of age or from 11 years to 12 years of age between their doses in the vaccination series should receive their age-appropriate dose at the time of the vaccination and the interval between doses is determined by the individual's age at the start of the vaccination series.

### ***Interchangeability***

There are limited data on the interchangeability of COMIRNATY with other COVID-19 vaccines to complete the primary vaccination course or any subsequent doses. Individuals who have received 1 dose of COMIRNATY Original/Omicron BA.4-5 should continue to receive COMIRNATY Original/Omicron BA.4-5 to complete the primary vaccination course and for any additional doses.

### **Method of administration**

In individuals 5 years of age and older, administer the vaccine intramuscularly in the deltoid muscle.

In individuals 1 to <5 years of age and older, administer the vaccine intramuscularly in the anterolateral aspect of the thigh or the deltoid muscle.

In individuals from 6 to <12 months of age, administer the vaccine intramuscularly in the anterolateral aspect of the thigh.

Do not inject the vaccine intravascularly, subcutaneously or intradermally.

COMIRNATY Original/Omicron BA.4-5 should not be mixed in the same syringe with any other vaccines or medicinal products.

For precautions to be taken before administering COMIRNATY Original/Omicron BA.4-5, see Section 4.4 Special warnings and precautions for use.

### ***Handling Instructions***

#### ***Handling prior to use***

Frozen vials must be completely thawed prior to use. Frozen vials should be transferred to 2 °C to 8 °C to thaw. Thaw times for 10-vial packs are noted in table below:

<b>Vial Cap Color</b>	<b>Time That May Be Required For a 10-vial Pack to Thaw (at 2 °C to 8 °C)</b>
Light Grey Light Blue Maroon	2 hours
Orange	4 hours
Dark Grey Dark Blue	6 hours

- Upon moving frozen vaccine to 2 °C to 8 °C storage, update the expiry date on the carton. The updated expiry date should reflect 10 weeks from the date of transfer to refrigerated conditions (2 °C to 8 °C) and not exceeding the original printed expiry date (EXP).
- Alternatively, individual frozen vials may be thawed for 30 minutes at temperatures up to 30 °C for immediate use.
- If the vaccine is received at 2 °C to 8 °C it should continue to be stored at 2 °C to 8 °C. Check that the carton has been previously updated to reflect the 10-week refrigerated expiry date.
- Unopened vials can be stored for up to 12 hours at temperatures up to 30 °C. Total storage time between 8 °C to 30 °C, inclusive of storage before and after puncture, should not exceed 24 hours.

## ***COMIRNATY Original/Omicron BA.4-5 Suspension for Injection***

### ***Preparation for administration***

COMIRNATY Original/Omicron BA.4-5 Suspension for Injection should be prepared by a healthcare professional using aseptic technique to ensure the sterility of the prepared suspension.

Vials of COMIRNATY Original/Omicron BA.4-5 Suspension for Injection have either a grey or a blue cap, contain either 1 or 6 doses of 0.3 mL of vaccine and do not require dilution.

- Light Grey or Light Blue cap: single dose vial
- Dark Grey or Dark Blue cap: 6 dose multidose vial

### ***Vial verification***

Prior to administration, check the name and strength of the vaccine on the vial label and the colour of the vial cap and vial label border to ensure it is the intended presentation. Check whether the vial is a single dose vial or a multidose vial and check if the vial requires dilution.

- Check appearance of vaccine prior to mixing and administration.
  - Grey cap vials: Prior to mixing, the vaccine is a white to off-white dispersion and may contain white to off-white opaque amorphous particles.
  - Blue cap vials: Prior to mixing, the vaccine is a clear to slightly opalescent dispersion and may contain white to off-white opaque amorphous particles.
- Gently invert the vial 10 times. **Do not shake.**
- Do not use the vaccine if particulates or discoloration are present after mixing.

### ***Preparation of individual doses***

- Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab.
- Withdraw a 0.3 mL single dose.
- *For Dark Grey or Dark Blue cap multidose vials (6 doses per vial):*
  - After first puncture, record appropriate date and time on the vial and store at 2 °C to 30 °C for up to 12 hours. Do not re-freeze.
  - Each dose must contain 0.3 mL of vaccine. Low dead-volume syringes and/or needles should be used in order to extract all doses from a single vial. The low dead-volume syringe and needle combination should have a dead volume of no more than 35 microliters.
  - If the amount of vaccine remaining in the vial cannot provide a full dose, discard the vial and any excess volume.
  - Do not pool excess vaccine from multiple vials.

## ***COMIRNATY Original/Omicron BA.4-5 Concentrated Suspension for Injection***

### ***Preparation for administration***

COMIRNATY Original/Omicron BA.4-5 Concentrated Suspension for Injection should be prepared by a healthcare professional using aseptic technique to ensure the sterility of the prepared diluted suspension.

Vials of COMIRNATY Original/Omicron BA.4-5 Concentrated Suspension for Injection have either an Orange or a Maroon cap, contains 10 doses of 0.2 mL of vaccine after dilution.

### ***Vial verification***

Prior to administration, check the name and strength of the vaccine on the vial label and the colour of the vial cap and vial label border to ensure it is the intended presentation. Check whether the vial is a single dose vial or a multidose vial and check if the vial requires dilution.

#### Prior to dilution

- After the thawed vial has reached room temperature, gently invert it 10 times prior to dilution. **Do not shake.**
- Check appearance of vaccine.
  - *Orange or Maroon cap vials:* Prior to dilution, the vaccine is a white to off-white dispersion and may contain white to off-white opaque amorphous particles.

#### Dilution instructions

- Thawed vaccine must be diluted in its original vial with sodium chloride 9 mg/mL (0.9%) solution for injection, using a 21 gauge or narrower needle and aseptic techniques. Volume of sodium chloride 9 mg/mL (0.9%) required are noted below:
  - *Orange cap vials:* 1.3 mL of sodium chloride 9 mg/mL
  - *Maroon cap vials:* 2.2 mL of sodium chloride 9 mg/mL
- Equalize vial pressure before removing the needle from the vial stopper by withdrawing air into the empty diluent syringe. Volume of air required are noted below:
  - *Orange cap vials:* 1.3 mL of air
  - *Maroon cap vials:* 2.2 mL of air
- Gently invert the diluted dispersion 10 times. **Do not shake.**
- Check appearance of vaccine after dilution.
  - *Orange or Maroon cap vials:* The diluted vaccine should present as a white to off-white dispersion with no particulates visible. Do not use the diluted vaccine if particulates or discoloration are present.
- After dilution, mark vial with appropriate date/time, store at 2 °C to 30 °C and use within 12 hours. Do not re-freeze.

#### Preparation of individual doses

- Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab.
- Withdraw a single dose.
  - *Orange or Maroon cap multidose vials (10 doses per vial):* each dose must contain 0.2 mL of vaccine. Low dead-volume syringes and/or needles should be used in order to extract all doses from a single vial. The low dead-volume syringe and needle combination should have a dead volume of no more than 35 microliters.
  - If the amount of vaccine remaining in the vial cannot provide a full dose, discard the vial and any excess volume.
  - Do not pool excess vaccine from multiple vials.

### 4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in Section 6.1 List of excipients.

### 4.4 Special warnings and precautions for use

#### Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be recorded in the Australian Immunisation Register.

## **General recommendations**

### ***Hypersensitivity and anaphylaxis***

Events of anaphylaxis have been reported. Appropriate medical treatment and supervision should always be readily available in case of an anaphylactic reaction following the administration of the vaccine.

The individual should be kept under close observation for at least 15 minutes following vaccination. A second dose of COMIRNATY Original/Omicron BA.4-5 should not be given to those who have experienced anaphylaxis to the first dose of COMIRNATY COVID-19 vaccines.

### ***Myocarditis and pericarditis***

Very rare cases of myocarditis and pericarditis have been observed following vaccination with COMIRNATY. Cases have occurred following first and second vaccinations and following booster doses. These cases have primarily occurred within 14 days following vaccination, more often after the second vaccination, and more often, but not exclusively in younger males. There have been reports in females. Based on accumulating data, the reporting rates of myocarditis and pericarditis after primary series in children ages 5 to <12 years are lower than in ages 12 to 17 years. Rates of myocarditis and pericarditis in booster doses do not appear to be higher than after the second dose in the primary series. Available data suggest that the course of myocarditis and pericarditis following vaccination is not different from myocarditis or pericarditis in general. Cases of myocarditis and pericarditis following vaccination have rarely been associated with severe outcomes including death.

Healthcare professionals should be alert to the signs and symptoms of myocarditis and pericarditis, including atypical presentations. Vaccinees should be instructed to seek immediate medical attention if they develop symptoms indicative of myocarditis or pericarditis such as (acute and persisting) chest pain, shortness of breath, or palpitations following vaccination. Non-specific symptoms of myocarditis and pericarditis also include fatigue, nausea and vomiting, abdominal pain, dizziness or syncope, oedema and cough. Healthcare professionals should consult guidance and/or specialists to diagnose and treat this condition.

For further details, please refer to the relevant clinical guidelines developed by the Australian Technical Advisory Group on Immunisation.

### ***Anxiety-related reactions***

Anxiety-related reactions, including vasovagal reactions (syncope), hyperventilation or stress-related reactions may occur in association with vaccination as a psychogenic response to the needle injection. It is important that precautions are in place to avoid injury from fainting.

Some individuals may have stress-related responses associated with the process of vaccination itself. Stress-related responses are temporary and resolve on their own. They may include dizziness, fainting, palpitations, increases in heart rate, alterations in blood pressure, feeling short of breath, tingling sensations, sweating and/or anxiety. Individuals should be advised to bring symptoms to the attention of the vaccination provider for evaluation and precautions should be in place to avoid injury from fainting.

### ***Syncope***

Syncope (fainting) may occur in association with administration of injectable vaccines. Procedures should be in place to avoid injury from fainting.

### ***Concurrent illness***

Vaccination should be postponed in individuals suffering from acute severe febrile illness or acute infection. The presence of a minor infection and/or low grade fever should not delay vaccination.

### ***Thrombocytopenia and coagulation disorders***

As with other intramuscular injections, the vaccine should be given with caution in individuals receiving anticoagulant therapy or those with thrombocytopenia or any coagulation disorder (such as haemophilia) because bleeding or bruising may occur following an intramuscular administration in these individuals.

### ***Immunocompromised individuals***

The efficacy, safety and immunogenicity of the vaccine has not been assessed in immunocompromised individuals, including those receiving immunosuppressant therapy.

Safety and immunogenicity after administration of COMIRNATY (tozinameran) have been assessed in a limited number of immunocompromised individuals (see Sections 4.8 Adverse effects and 5.1 Pharmacodynamic properties). The efficacy of COMIRNATY Original/Omicron BA.4-5 may be lower in immunosuppressed individuals.

### ***Duration of protection***

The duration of protection afforded by the vaccine is unknown as it is still being determined by ongoing clinical trials and observational studies.

### ***Limitations of vaccine effectiveness***

As with any vaccine, vaccination with COMIRNATY Original/Omicron BA.4-5 may not protect all vaccine recipients. Individuals may not be fully protected until 7 days after completion of their primary course of the vaccine.

### ***Use in the elderly***

Clinical studies of COMIRNATY Original/Omicron BA.4-5 include participants 55 years of age and older and their data contributes to the overall assessment of safety and immunogenicity. The safety of Comirnaty Original/Omicron BA.4-5 as a second booster in individuals 65 years of age and older is based on safety data in 159 booster dose recipients 65 years of age and older and 35 booster dose recipients 75 years of age and older in C4591044. See Section 5.1 Pharmacodynamic properties, Clinical trials, Efficacy against COVID-19. No dosage adjustment is required in elderly individuals  $\geq 65$  years of age.

The data for use in the frail elderly is limited. The potential benefits of vaccination versus the potential risk and clinical impact of even relatively mild systemic adverse events in the frail elderly should be carefully assessed on a case-by-case basis.

The safety of a booster dose of COMIRNATY (tozinameran) in individuals 65 years of age and older is based on safety data in 12 booster dose recipients 65 to 85 years of age in Study C4591001, 306 booster dose recipients 18 to 55 years of age in Study C4591001, and 1,175 booster dose recipients 65 years of age and older in Study C4591031.

The effectiveness of a booster dose of COMIRNATY (tozinameran) in individuals 65 years of age and older is based on effectiveness data in 306 booster dose recipients 18 to 55 years of age in Study C4591001, and an efficacy analysis from participants 16 years of age and older in 9,945 participants in Study C4591031.

## **Paediatric use**

The safety and efficacy of COMIRNATY Original/Omicron BA.4-5 in infants aged less than 6 months of age have not yet been established.

Limited safety and effectiveness data is available for COMIRNATY (tozinameran) booster dose in adolescents 12 to 15 years of age and no immunogenicity data is available for booster dose in this age group. The safety and effectiveness of a booster dose of COMIRNATY (tozinameran) in individuals 12 to 17 years of age is based on safety and effectiveness data in adults at least 18 to 55 years of age.

Real world evidence from the Ministry of Health of Israel and surveillance by CDC in USA on the administration of third doses of COMIRNATY (tozinameran) given after the primary course revealed no new safety concerns in adolescents 12 to 17 years of age.

Very rare cases of myocarditis and pericarditis have been observed following vaccination with COMIRNATY in adolescents (see Section 4.4 Special warnings and precautions for use, Myocarditis and pericarditis).

## **Effects on laboratory tests**

No data available.

## **4.5 Interactions with other medicines and other forms of interactions**

COMIRNATY (15/15 micrograms/dose only) may be administered concomitantly with seasonal influenza vaccine (see Section 5.1 Pharmacodynamic properties, Concomitant vaccine administration with influenza vaccine).

Different injectable vaccines should be given at different injection sites.

Do not mix COMIRNATY with other vaccines or products in the same syringe.

## **4.6 Fertility, pregnancy and lactation**

### **Effects on fertility**

There are no animal reproductive toxicity studies with COMIRNATY Original/Omicron BA.4-5.

In a combined fertility and developmental toxicity study, female rats were intramuscularly administered COMIRNATY prior to mating and during gestation (4 full human doses of 30 µg each, spanning between pre-mating day 21 and gestation day 20). SARS CoV-2 neutralising antibodies were present in maternal animals from prior to mating to the end of the study on postnatal day 21 as well as in fetuses and offspring. There were no vaccine related effects on female fertility and pregnancy rate.

### **Use in pregnancy - Pregnancy Category B1**

No data are available yet regarding the use of COMIRNATY Original/Omicron BA.4-5 during pregnancy. However, a large amount of information from pregnant women vaccinated with the initially approved COMIRNATY (tozinameran) vaccine during the second and third trimester have not shown negative effects on the pregnancy or the newborn baby. While information on effects on pregnancy or the newborn baby after vaccination during the first trimester is limited, no change to the risk for miscarriage has been seen.



There are limited clinical study data with use of COMIRNATY (tozinameran) in pregnant women (see Section 4.8 Adverse effects (undesirable effects)).

Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryo/fetal development, parturition or post-natal development (see Section 4.6 Fertility, pregnancy and lactation, Effects on fertility). Administration of COMIRNATY Original/Omicron BA.4-5 in pregnancy should only be considered when the potential benefits outweigh any potential risks for the mother and fetus.

### **Use in lactation**

No data are available yet regarding the use of COMIRNATY Original/Omicron BA.4-5 during breast-feeding. COMIRNATY Original/Omicron BA.4-5 can be used while breast-feeding, when the potential benefits outweigh any potential risks for the mother and baby.

It is unknown whether tozinameran is excreted in human milk. A combined fertility and developmental toxicity study in rats did not show harmful effects on offspring development before weaning (see Section 4.6 Fertility, pregnancy and lactation, Effects on fertility).

## **4.7 Effects on ability to drive and use machines**

COMIRNATY Original/Omicron BA.4-5 has no, or negligible, influence on the ability to drive and use machines. However, some of the effects mentioned under Section 4.8 Adverse effects (undesirable effects) may temporarily affect the ability to drive or use machines.

## **4.8 Adverse effects (undesirable effects)**

### **Summary of safety profile**

#### **COMIRNATY Original/Omicron BA.4-5 (tozinameran/famtozinameran)**

##### ***Participants 12 years of age and older – after bivalent Omicron BA.4-5 booster dose***

In a subset from Study C4591044 (Phase 2/3), 107 participants 12 to 17 years of age, 313 participants 18 to 55 years of age and 306 participants 56 years of age and older previously vaccinated with a 2-dose primary series and 1 booster dose of COMIRNATY (tozinameran) went on to receive a second booster dose with COMIRNATY Original/Omicron BA.4-5 (15/15 micrograms) 5.4 to 16.9 months after receiving the first booster dose and had a median follow up time of at least 1.5 months up to a data cut-off date 12 October 2022 (Cohort 2) and 31 October 2022 (Cohort 3).

The overall safety profile for the COMIRNATY Original/Omicron BA.4-5 booster (fourth dose) was similar to that seen after the COMIRNATY (tozinameran) booster (third dose). The most frequent adverse reactions in participants 12 years of age and older were injection site pain (> 60%), fatigue (> 50%), headache (> 40%), muscle pain (> 20%), chills (> 10%) and joint pain (> 10%). No new adverse reactions were identified for COMIRNATY Original/Omicron BA.4-5.

At present, data relating only to short term (1 month post booster) local and systemic effects are available. Long term safety data for COMIRNATY Original/Omicron BA.4-5 (tozinameran/famtozinameran) are not available.

### ***Participants 5 to <12 years of age – after bivalent Omicron BA.4-5 booster dose***

In a subset from Study C4591048 (Phase 3), 113 participants 5 to 11 years of age who had completed a 2-dose primary series and 1 booster dose of COMIRNATY (tozinameran), received a second booster dose of COMIRNATY Original/Omicron BA.4-5 (5/5 micrograms) 2.6 to 8.5 months after receiving the first booster dose. Participants who received a booster dose of COMIRNATY Original/Omicron BA.4-5 had a median follow-up time of 6.3 months up to a data cut-off date of 20 April 2023.

The overall safety profile for the COMIRNATY Original/Omicron BA.4-5 booster was similar to that seen after 3 doses of COMIRNATY (tozinameran). The most frequent adverse reactions in participants 5 to <12 years of age were injection site pain (>60%), fatigue (>40%), headache (>20%), and muscle pain (>10%).

### ***Participants 2 to <5 years of age – after bivalent Omicron BA.4-5 booster dose***

In 2 groups in Study C4591048 (Phase 3, Groups 2 and 3), 1, 207 participants (n = 218 Group 2, n = 989 Group 3) 2 to 4 years of age who had completed a 3-dose primary series, received a booster dose of COMIRNATY Original/Omicron BA.4-5 (1.5/1.5 micrograms) 2.1 to 8.6 months after receiving Dose 3 for Group 2 and 2.8 to 17.5 months after receiving Dose 3 for Group 3.

Participants who received a booster dose of COMIRNATY Original/Omicron BA.4-5 had a median follow-up time of 4.6 months up to a data cut-off date of 03 March 2023 for Group 2 and had a median follow-up time of 6.3 months for up to a data cut off date of 08 February 2024 for Group 3.

The overall safety profile for the COMIRNATY Original/Omicron BA.4-5 booster was similar to that seen after 3 doses of COMIRNATY (tozinameran). The most frequent adverse reactions in participants 2 to <5 years of age were injection site pain (>30%) and fatigue (>20%). Pyrexia was commonly observed (>5%).

### ***Participants 6 months to <2 years of age – after bivalent Omicron BA.4-5 booster dose***

In 2 groups in Study C4591048 (Phase 3, Groups 2 and 3), 160 participants (n = 92 Group 2, n = 68 Group 3) 6 to 23 months of age who had completed a 3-dose primary series, received a booster dose of COMIRNATY Original/Omicron BA.4-5 (1.5/1.5 micrograms) 2.1 to 8.6 months after receiving Dose 3 for Group 2 and 3.8 to 12.5 months after receiving Dose 3 for Group 3.

Participants who received a booster dose of COMIRNATY Original/Omicron BA.4-5 had a median follow-up time of 4.4 months up to a data cut-off date of 03 March 2023 for Group 2 and had a median follow up time of 6.4 months for up to a data cut-off date of 08 February 2024 for Group 3.

The overall safety profile for the COMIRNATY Original/Omicron BA.4-5 booster was similar to that seen after 3 doses of COMIRNATY (tozinameran). The most frequent adverse reaction in participants 6 to 23 months of age were irritability (>30%), decreased appetite (>20%), drowsiness (>19%), tenderness at the injection site (>10%), and fever (>10%).

### **COMIRNATY (tozinameran)**

The safety of COMIRNATY (tozinameran) was evaluated in participants aged 6 months and older in clinical studies (comprised of 22,026 participants 16 years of age and older and 1,131 adolescents 12 to 15 years of age from Study C4591001, and 3,109 children 5 to <12 years of

age, 2,368 participants 2 to <5 years of age and 1,458 participants 6 months to <2 years of age from Study C4591007) that have received at least one dose of COMIRNATY (tozinameran).

Additionally, 306 existing Phase 3 participants 18 to 55 years of age received a booster dose of COMIRNATY (tozinameran) approximately 6 months after the second dose in the non-placebo-controlled booster dose portion of Study C4591001. The overall safety profile for the booster dose was similar to that seen after 2 doses.

In Study C4591031, a placebo-controlled booster study, 5,081 participants 16 years of age and older were recruited from Study C4591001 to receive a booster dose of COMIRNATY (tozinameran) at least 6 months after the second dose. The overall safety profile for the booster dose was similar to that seen after 2 doses.

In a subset of C4591007 Phase 2/3 participants, 2,408 participants 5 to <12 years of age received a booster dose of COMIRNATY (tozinameran) at least 5 months (range 5.3 to 19.4 months) after completing the primary series. The overall safety profile for the booster dose was similar to that seen after the primary series.

### ***Participants 16 years of age and older – after 2 doses***

In Study C4591001, a total of 22,026 participants 16 years of age or older received at least 1 dose of COMIRNATY (tozinameran) 30 micrograms and a total of 22,021 participants 16 years of age or older received placebo (including 138 and 145 adolescents 16 and 17 years of age in the COMIRNATY (tozinameran) and placebo groups, respectively). A total of 20,519 participants 16 years of age or older received 2 doses of COMIRNATY (tozinameran).

At the time of the analysis of Study C4591001 with a data cut-off of 13 March 2021 for the placebo-controlled blinded follow-up period up to the participants' unblinding dates, a total of 25,651 (58.2%) participants [13,031 COMIRNATY (tozinameran) and 12,620 placebo] 16 years of age and older were followed up for  $\geq 4$  months after the second dose. This included a total of 15,111 [7,704 COMIRNATY (tozinameran) and 7,407 placebo] participants 16 to 55 years of age and a total of 10,540 [5,327 COMIRNATY (tozinameran) and 5,213 placebo] participants 56 years and older.

The most frequent adverse reactions in participants 16 years of age and older that received 2 doses were injection site pain (>80%), fatigue (>60%), headache (>50%), myalgia (>40%), chills (>30%), arthralgia (>20%), pyrexia and injection site swelling (>10%) and were usually mild or moderate in intensity and resolved within a few days after vaccination. A slightly lower frequency of reactogenicity events was associated with greater age.

The safety profile in 545 subjects receiving COMIRNATY (tozinameran), that were seropositive for SARS-CoV-2 at baseline, was similar to that seen in the general population.

Study C4591001 also included 200 participants with confirmed stable human immunodeficiency virus (HIV) infection. The safety profile of the participants receiving COMIRNATY (tozinameran) (n=100) in the individuals with stable HIV infection was similar to that seen in the general population.

### ***Adolescents 12 to 15 years of age – after 2 doses***

In an analysis of long term safety follow-up in Study C4591001, 2,260 adolescents [1,131 COMIRNATY (tozinameran) 30 micrograms; 1,129 placebo] were 12 to 15 years of age. Of these, 1,559 adolescents [786 COMIRNATY (tozinameran); 773 placebo] were followed for  $\geq 4$  months after the second dose of COMIRNATY (tozinameran).

The most frequent adverse reactions in adolescents 12 to 15 years of age that received 2 doses were injection site pain (>90%), fatigue and headache (>70%), myalgia and chills (>40%), arthralgia and pyrexia (>20%).

### ***Children 5 to <12 years of age – after 2 doses***

In an analysis of Study C4591007 Phase 2/3, 4,647 children [3,109 COMIRNATY (tozinameran) 10 micrograms; 1,538 placebo] were 5 to <12 years of age. Of these, 2,206 [1,481 COMIRNATY (tozinameran) 10 micrograms; 725 placebo] children have been followed for at least  $\geq 4$  months after the second dose in the placebo-controlled blinded follow-up period. The safety evaluation in Study C4591007 is ongoing.

The most frequent adverse reactions in children 5 to <12 years of age that received 2 doses included injection site pain (>80%), fatigue (>50%), headache (>30%), injection site redness and swelling ( $\geq 20\%$ ), myalgia, chills and diarrhoea (>10%).

### ***Children 2 to <5 years of age – after 3 doses***

In an analysis of Study C4591007 (Phase 2/3), 3,541 children [2,368 COMIRNATY (tozinameran) 3 micrograms; 1,173 placebo] were 2 to <5 years of age. Based on data in the blinded placebo-controlled follow-up period up to the cutoff date of 28 February 2023, 1,268 children 2 to <5 years of age who received a 3-dose primary course [863 COMIRNATY (tozinameran) 3 micrograms; 405 placebo] have been followed a median of 2.2 months after the third dose.

The most frequent adverse reactions in children 2 to <5 years of age that received any primary course dose included pain at injection site and fatigue (>40%), injection site redness and fever (>10%).

### ***Infants 6 months to <2 years of age – after 3 doses***

In an analysis of Study C4591007 (Phase 2/3), 1,776 infants [1,178 COMIRNATY (tozinameran) 3 micrograms and 598 placebo] were 6 months to <2 years of age. Based on data in the blinded placebo-controlled follow-up period up to the cutoff date of 29 April 2022, 570 infants 6 months to <2 years of age who received a 3-dose primary course [386 COMIRNATY 3 micrograms and 184 placebo] have been followed for a median of 1.3 months after the third dose.

The most frequent adverse reactions in infants 6 months to <2 years of age that received any primary course dose included irritability (>60%), decrease appetite (>30%), tenderness at the injection site (>20%), injection site redness and fever (>10%).

### ***Participants 16 years of age and older – after booster dose***

A subset from Study C4591001 Phase 2/3 participants of 306 adults 18 to 55 years of age who completed the original COMIRNATY (tozinameran) 2-dose course, received a booster dose of COMIRNATY (tozinameran) approximately 6 months (range of 4.8 to 8.0 months) after receiving Dose 2. Of these, 301 participants have been followed for  $\geq 4$  months after the booster dose of COMIRNATY (tozinameran).

The most frequent adverse reactions in participants 18 to 55 years of age were injection site pain (>80%), fatigue (>60%), headache (>40%), myalgia (>30%), chills and arthralgia (>20%).

In Study C4591031, a placebo-controlled booster study, participants 16 years of age and older recruited from Study C4591001 received a booster dose of COMIRNATY (tozinameran) (5,081 participants), or placebo (5,044 participants) at least 6 months after the second dose of

COMIRNATY (tozinameran). Overall, participants who received a booster dose, had a median follow-up time of 2.8 months (range 0.3 to 7.5 months) after the booster dose in the blinded placebo-controlled follow-up period to the cut-off date (8 February 2022). Of these, 1281 participants [895 COMIRNATY (tozinameran) and 386 placebo] were followed for  $\geq 4$  months after the booster dose of COMIRNATY (tozinameran). The overall safety profile for the booster dose was similar to that seen after 2 doses.

In another subset from Study C4591001, 825 adolescents 12 to 15 years of age who completed the COMIRNATY (tozinameran) 2-dose course, received a booster dose of COMIRNATY (tozinameran) approximately 11.2 months (range of 6.3 to 20.1 months) after receiving Dose 2. Overall, participants who received a booster dose, had a median follow-up time of 9.5 months (range 1.5 to 10.7 months) based on data up to the cut-off date (3 November 2022). No new adverse reactions of COMIRNATY (tozinameran) were identified.

### ***Participants 18 years of age and older – after subsequent booster doses***

In a subset from study C4591031 (Phase 3), 325 adults 18 to  $\leq 55$  years of age who had completed 3 doses of COMIRNATY (tozinameran) received a booster (fourth dose) of COMIRNATY (tozinameran 30 micrograms) 90 to 180 days after receiving Dose 3. Participants who received a booster (fourth dose) of COMIRNATY (tozinameran 30 micrograms) had a median follow-up time of 1.4 months. The most frequent adverse reactions in these participants were injection site pain ( $>70\%$ ), fatigue ( $>60\%$ ), headache ( $>40\%$ ), myalgia and chills ( $>20\%$ ) and arthralgia ( $>10\%$ ).

In a subset from Study C4591031 (Phase 3), 305 adults greater than 55 years of age who had completed 3 doses of COMIRNATY (tozinameran), received a booster (fourth dose) of COMIRNATY (tozinameran 30 micrograms) 5.3 to 13.1 months after receiving Dose 3. Participants who received a booster (fourth dose) of COMIRNATY (tozinameran 30 micrograms) had a median follow-up time of at least 1.7 months up to a data cutoff date of 16 May 2022. The most frequent adverse reactions in participants greater than 55 years of age were injection site pain (60%), fatigue ( $>40\%$ ), headache ( $>20\%$ ), myalgia and chills ( $>10\%$ ).

### ***Children 5 to $<12$ years of age – after booster dose***

In a subset from C4591007, a total of 2,408 children 5 to  $<12$  years of age received a booster dose of COMIRNATY (tozinameran) 10 micrograms at least 5 months (range 5.3 to 19.4 months) after completing the primary series. The analysis of the C4591007 Phase 2/3 subset is based on data up to the cut-off date of 28 February 2023 (median follow-up time of 6.4 months).

The most frequent adverse reactions in participants 5 to  $<12$  years of age were injection site pain ( $>60\%$ ), fatigue ( $>30\%$ ), headache ( $>20\%$ ), myalgia, chills, injection site redness, and swelling ( $>10\%$ ).

A higher frequency of lymphadenopathy was observed in C4591007 in participants receiving a booster dose compared to participants receiving 2 doses (2.5% vs. 0.9%).

### **Tabulated list of adverse reactions from clinical studies**

Adverse reactions observed during clinical studies are listed below according to the following frequency categories:

Very common ( $\geq 1/10$ ),  
Common ( $\geq 1/100$  to  $<1/10$ ),  
Uncommon ( $\geq 1/1,000$  to  $<1/100$ ),  
Rare ( $\geq 1/10,000$  to  $<1/1,000$ ),

Very rare (<1/10,000),  
Not known (cannot be estimated from the available data).

**Table 1: Adverse reactions from COMIRNATY Original/Omicron BA.4-5 clinical trial (C4591044 Cohort 2 and Cohort 3 combined): Individuals 18 to 55 years and >55 years of age (Cohort 2 12 October 2022 Data Cut-off Date and Cohort 3 31 October 2022 Data Cut-off Date)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders			Lymphadenopathy (18-55)	Lymphadenopathy (>55)	
Immune system disorders				Urticaria <sup>b</sup> (18-55) Pruritus <sup>b</sup> (>55)	
Nervous system disorders	Headache				
Gastrointestinal disorders		Vomiting <sup>a</sup> Diarrhoea <sup>a</sup> (>55)	Diarrhoea <sup>a</sup> (18-55)		
Musculoskeletal and connective tissue disorders	Arthralgia; Myalgia			Pain in extremity (arm) <sup>b</sup> (>55)	
General disorders and administration site conditions	Injection site pain; Fatigue; Chills;	Pyrexia; Injection site swelling; Injection site redness			

a. These adverse reactions were identified in the post-authorisation period.

b. The following events are categorised as hypersensitivity reactions: urticaria, pruritus, rash, and angioedema.

**Table 2: Adverse reactions from COMIRNATY Original/Omicron BA.4-5 clinical trial (C4591044 Cohort 2): Individuals 12 to 17 years of age (12 October 2022 Data Cut-off Date)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders					
Nervous system disorders	Headache				
Gastrointestinal disorders		Diarrhoea <sup>a</sup> ; Vomiting <sup>a</sup>			
Musculoskeletal and connective tissue disorders	Arthralgia; Myalgia				
General disorders and administration site conditions	Injection site pain; Fatigue; Chills;	Pyrexia; Injection site swelling; Injection site redness			

<sup>a</sup> These adverse reactions were identified in the post-authorisation period.

**Table 3: Adverse reactions from COMIRNATY Original/Omicron BA.4-5 clinical trial (C4591048 SSD): Individuals 5 to <12 years of age (25 November 2022 Data Cut-off Date)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders			Lymphadenopathy		
Nervous system disorders	Headache				
Gastrointestinal disorders		Diarrhoea <sup>a</sup> ; Vomiting <sup>a</sup>			
Musculoskeletal and connective tissue disorders	Myalgia	Arthralgia			
General disorders and administration site conditions	Injection site pain; Fatigue	Pyrexia; Chills; Injection site swelling; Injection site redness			

<sup>a</sup>. These adverse reactions were identified in the post-authorisation period

**Table 4: Adverse reactions from COMIRNATY Original/Omicron BA.4-5 clinical trial (C4591048 SSB): Individuals 2 Years to <5 Years of Age (Data Cut-off Date: Group 2- 3 Mar 2023 and Group 3-08 Feb 2024)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders			Lymphadenopathy		
Immune system disorders			Rash <sup>a,b</sup> Urticaria <sup>a,b</sup>		
Nervous system disorders		Headache			
Gastrointestinal disorders		Vomiting <sup>a</sup> ; Diarrhoea <sup>a</sup>			
Musculoskeletal and connective tissue disorders		Myalgia;	Arthralgia		
General disorders and administration site conditions	Injection site pain; Fatigue	Pyrexia; Injection site redness; Chills; Injection site swelling			

a. These adverse reactions were identified in the post-authorisation period.

b. The following events are categorised as hypersensitivity reactions: urticaria and rash.

**Table 5: Adverse reactions from COMIRNATY Original/Omicron BA.4-5 clinical trial (C4591048 SSB): Individuals 6 Months to <2 Years of Age (Data Cut-off Date: Group 2- 3 Mar 2023 and Group 3-08 Feb 2024)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
Immune system disorders			Rash <sup>b</sup>		
Metabolism and nutrition disorders	Decreased appetite				
Psychiatric disorders	Irritability				
Gastrointestinal disorders		Diarrhoea <sup>a</sup> ; Vomiting <sup>a</sup>			
Musculoskeletal and connective tissue disorders			Pain in extremity (arm)		
General disorders and administration site conditions	Pyrexia; Injection site tenderness; Drowsiness	Fatigue; Injection site swelling; Injection site redness	Chills		

a. These adverse reactions were identified in the post-authorisation period. At the time of the data cut-off date, the following reactions were not reported in participants 6 months to <2 years of age in Study C4591048: pruritus, angioedema, dizziness, lethargy, myocarditis, pericarditis, hyperhidrosis, night sweats, and malaise but are still considered ADRs for this age group.

b. The following event is categorised as a hypersensitivity reaction: rash

**Table 6: Adverse reactions from COMIRNATY (tozinameran) clinical trial (C4591001): Individuals 12 years of age and older**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders			Lymphadenopathy <sup>a</sup>		
Psychiatric disorders			Insomnia		
Metabolism and nutrition disorders			Decreased appetite		
Nervous system disorders	Headache		Lethargy	Acute peripheral facial paralysis <sup>b</sup>	
Gastrointestinal disorders		Nausea			
Skin and subcutaneous disorders			Hyperhidrosis Night sweats		
Musculoskeletal and connective tissue disorders	Arthralgia; Myalgia				



System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
General disorders and administration site conditions	Injection site pain; Fatigue; Chills; Pyrexia <sup>c</sup> ; Injection site swelling	Injection site redness	Asthenia Malaise		Facial swelling <sup>d</sup>

a A higher frequency of lymphadenopathy (2.8% vs 0.4%) was observed in participants receiving a booster dose in study C4591031 compared to participants receiving 2 doses.

b Through the clinical trial safety follow-up period to 14 November 2020, acute peripheral facial paralysis (or palsy) was reported by four participants in the COMIRNATY (tozinameran) group. Onset was Day 37 after Dose 1 (participant did not receive Dose 2) and Days 3, 9, and 48 after Dose 2. No cases of acute peripheral facial paralysis (or palsy) were reported in the placebo group.

c A higher frequency of pyrexia was observed after the second dose.

d Facial swelling in vaccine recipients with a history of injection of dermatological fillers

The safety profile in 545 subjects receiving COMIRNATY (tozinameran), that were seropositive for SARS-CoV-2 at baseline, was similar to that seen in the general population.

**Table 7. Adverse Reactions from COMIRNATY (tozinameran) clinical trial (C4591007): Individuals 5 to <12 Years of Age (20 May 2022 Data Cut-off Date)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders			Lymphadenopathy <sup>a</sup>		
Immune system disorders			Urticaria <sup>b,c</sup> ; Pruritus <sup>b,c</sup> ; Rash <sup>b,c</sup>	Angioedema <sup>b,c</sup>	Anaphylaxis <sup>b</sup>
Metabolism and nutrition disorders			Decreased appetite		
Nervous system disorders	Headache				
Gastrointestinal disorders	Diarrhoea <sup>b</sup>	Vomiting <sup>b</sup>	Nausea		
Skin and subcutaneous tissue disorders				Night sweats	
Musculoskeletal and connective tissue disorders	Myalgia	Arthralgia	Pain in extremity (arm) <sup>b</sup>		
General disorders and administration site conditions	Injection site pain; Fatigue; Chills; Injection site swelling; Injection site redness	Pyrexia	Malaise		

<sup>a</sup> A higher frequency of lymphadenopathy was observed in C4591007 (2.5% vs. 0.7%) in participants receiving a booster dose compared to participants receiving 2 doses.

<sup>b</sup> These adverse reactions were identified in the post-authorisation period. The following events were not reported in participants 5 to <12 Years of Age in Study C4591007 but were reported in individuals ≥16 years of age in Study C4591001: angioedema, lethargy, asthenia, hyperhidrosis, and night sweats.

<sup>c</sup> The following events are categorised as hypersensitivity reactions: urticaria, pruritus, rash and angioedema.

**Table 8. Adverse Reactions from COMIRNATY (tozinameran) clinical trial: Individuals 2 to <5 Years of Age (28 February 2023 Data Cut-off Date)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Very Rare <1/10,000	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders				Lymphadenopathy		
Immune system disorders			Rash <sup>a,b</sup> ; Urticaria <sup>a,b</sup>			Anaphylaxis <sup>a</sup>
Metabolism and nutrition disorders				Decreased appetite		
Nervous system disorders		Headache				
Cardiac disorders					Myocarditis <sup>a</sup> Pericarditis <sup>a</sup>	
Gastrointestinal disorders	Diarrhoea <sup>a</sup>	Vomiting <sup>a</sup>	Nausea			
Musculoskeletal and connective tissue disorders		Myalgia Arthralgia	Pain in extremity (arm) <sup>a</sup>			
General disorders and administration site conditions	Injection site pain; Fatigue; Injection site redness; Pyrexia	Injection site swelling; Chills		Asthenia		

a. These adverse reactions were identified in the post-authorisation period. At the time of the data-lock, the following reactions were not reported in participants 2 to <5 Years of Age in Study C4591007: pruritus, angioedema, lethargy, myocarditis, pericarditis, hyperhidrosis, night sweats, and malaise but are still considered adverse reactions for this age group.

b. The following events are categorised as hypersensitivity reactions: rash and urticaria

**Table 9. Adverse Reactions from COMIRNATY (tozinameran) clinical trial: Individuals 6 Months to <2 Years of Age (28 February 2023 Data Cut-off Date)**

System Organ Class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Very Rare <1/10,000	Not known (cannot be estimated from the available data)
Blood and lymphatic system disorders			Lymphadenopathy			
Immune system disorders		Rash <sup>a,b</sup>	Urticaria <sup>a,b</sup> ;			Anaphylaxis <sup>a</sup>
Metabolism and nutrition disorders	Decreased appetite					
Cardiac disorders					Myocarditis <sup>a</sup> Pericarditis <sup>a</sup>	
Psychiatric disorders	Irritability					
Nervous system disorders			Headache Lethargy			
Gastrointestinal disorders		Vomiting <sup>a</sup> ; Diarrhoea <sup>a</sup>				

General disorders and administration site conditions	Injection site tenderness; Injection site redness; Pyrexia	Injection site swelling	Fatigue; Chills			
--	--	-------------------------	-----------------	--	--	--

a. These adverse reactions were identified in the post-authorisation period. At the time of data-lock, the following events were not reported in participants 6 months to <2 Years of Age in Study C4591007: pruritus, angioedema, nausea, hyperhidrosis, night sweats, myalgia, arthralgia, pain in extremity (arm), malaise, and asthenia but are still considered adverse reactions for this age group.

b. The following events are categorised as hypersensitivity reactions: rash and urticaria

## Special populations

### ***Pregnant women and infants born to maternal participants – after 2 doses of COMIRNATY (tozinameran)***

Study C4591015, a Phase 2/3, placebo-controlled study, evaluated COMIRNATY (tozinameran) or placebo administered in 2 doses, approximately 21 days apart, in pregnant women 18 years of age and older, with the first dose given at 24 to 34 weeks gestation. A total of 346 pregnant women received COMIRNATY (tozinameran) (n=173) or placebo (n=173).

The most frequent adverse reactions in pregnant women who received any primary series dose with COMIRNATY (tozinameran) included injection site pain (>80%), fatigue (>60%), headache (>50%), myalgia (>30%), chills, arthralgia, and injection site swelling (>10%). No new adverse reactions were identified.

In Study C4591015, safety in infants born to maternal participants who received COMIRNATY (tozinameran) (n=167) or placebo (n=168) was evaluated up to 6 months after birth. No safety concerns were identified that were attributable to maternal vaccination with COMIRNATY (tozinameran).

### ***Immunocompromised participants (adults and children)***

In Study C4591024, 124 immunocompromised participants received at least 1 and up to 4 doses of COMIRNATY(tozinameran) ; 37 participants 2 to <5 years of age, 65 participants 5 to <12 years of age, 15 participants 12 to < 18 years of age, and 7 participants 18 years of age and older. Doses 1 and 2 were separated by 21 days, Doses 2 and 3 were separated by 28 days and Dose 4 was administered 3 to 6 months after Dose 3 (see Section 5.1 Pharmacodynamic properties).

The safety profile of COMIRNATY (tozinameran) in immunocompromised participants 2 years of age and older was consistent with that expected for an immunocompromised population and with the safety profile of COMIRNATY (tozinameran). No new adverse reactions were identified.

## Post-marketing experience

Although the events listed in Table 10 were not observed in the clinical trials, they are considered adverse drug reactions for COMIRNATY as they were reported in the post-marketing experience. As these reactions were derived from spontaneous reports, the frequencies could not be determined and are thus considered as not known.

**Table 10: Adverse reactions from COMIRNATY post marketing experience**

System Organ Class	Adverse Drug Reaction
Immune system disorders	Anaphylaxis Hypersensitivity reactions (e.g. rash, pruritis, urticaria, angioedema, erythema multiforme)
Cardiac disorders	Myocarditis Pericarditis

System Organ Class	Adverse Drug Reaction
Gastrointestinal disorders	Diarrhoea Vomiting
Musculoskeletal and connective tissue disorders	Pain in extremity (arm) <sup>a</sup>
General disorders and administration site conditions	Extensive swelling of vaccinated limb
Nervous system disorders	Paraesthesia Hypoaesthesia Dizziness Headache (including migraine)
Reproductive system and breast disorders	Non-sexually acquired genital ulceration Heavy menstrual bleeding* Breast swelling and Mastitis

<sup>a</sup> A higher frequency of pain in extremity (1.1% vs. 0.8%) was observed in participants receiving a booster dose in Study C4591031 compared to participants receiving 2 doses.

\* Most cases appear to be non-serious and temporary in nature

### Safety with concomitant vaccine administration

In Study C4591030, a Phase 3 study, participants 18 to 64 years of age who received COMIRNATY (tozinameran) coadministered with seasonal inactivated influenza vaccine (SIIV), quadrivalent followed 1 month later by placebo (n=564), were compared to participants who received an inactivated influenza vaccine with placebo followed 1 month later by COMIRNATY (tozinameran) alone (n=564). Reactogenicity events were reported more frequently by participants who received COMIRNATY (tozinameran) coadministered with SIIV, quadrivalent, compared to participants who received COMIRNATY (tozinameran) alone, but overall the reactogenicity events were mostly mild to moderate in severity. The most common adverse reactions reported in the coadministration group and after COMIRNATY (tozinameran) alone were injection site pain (86.2% and 84.4%, respectively), fatigue (64.0% and 50.8%, respectively) and headache (47.2% and 37.8%, respectively).

### Reporting suspected adverse effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

## 4.9 Overdose

In clinical trials, participants who received up to 2 times the recommended dose of COMIRNATY did not have an increase in reactogenicity or adverse reactions.

In post-authorisation experience, there have been reports of higher than recommended doses of COMIRNATY. In general, adverse events reported with overdoses have been similar to the known adverse reaction profile of COMIRNATY.

In the event of overdose, monitoring of vital functions and individualised symptomatic treatment is recommended.

For information on the management of overdose, contact the Poisons Information Centre on 131126 (Australia).

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: vaccines, Covid-19 RNA-based vaccines, ATC code: J07BN01

#### Mechanism of action

The nucleoside-modified messenger RNA in the vaccine is formulated in lipid nanoparticles, which enable delivery of the non-replicating RNA into host cells to direct transient expression of the SARS-CoV-2 spike (S) antigen. The mRNA codes for membrane-anchored, full-length S with two point mutations within the central helix. Mutation of these two amino acids to proline locks S in an antigenically preferred prefusion conformation. The vaccine elicits both neutralising antibody and cellular immune responses to the antigen, which may contribute to protection against COVID-19.

#### Clinical trials

#### Immunogenicity

##### COMIRNATY Original/Omicron BA.4-5 (tozinameran/famtozinameran)

##### *Relative vaccine immunogenicity in participants 12 years of age and older – after bivalent Omicron BA.4-5 (second booster dose)*

In an analysis of a subset from Study C4591044, 105 participants 12 to 17 years of age, 297 participants 18 to 55 years of age, and 286 participants 56 years of age and older who had previously received a 2-dose primary series and booster dose with COMIRNATY (tozinameran) received COMIRNATY Original/Omicron BA.4-5 (15/15 micrograms) as a second booster. In participants 12 to 17 years of age, 18 to 55 years of age, and 56 years of age and older, 75.2%, 71.7% and 61.5% were positive for SARS-CoV-2 at baseline, respectively.

Analyses of 50% neutralising antibody titres (NT50) against Omicron BA.4-5 and against reference strain among participants 56 years of age and older who received COMIRNATY Original/Omicron BA.4-5 as a second booster in Study C4591044 compared to a subset of participants from Study C4591031 who received a second booster of COMIRNATY (tozinameran) demonstrated superiority of COMIRNATY Original/Omicron BA.4-5 to COMIRNATY (tozinameran) based on geometric mean ratio (GMR) and noninferiority based on difference in seroresponse rates with respect to anti-Omicron BA.4-5 response, and noninferiority of anti-reference strain immune response based on GMR (Table 11 and Table 12).

Analyses of NT50 against Omicron BA.4-5 among participants 18 to 55 years of age compared to participants 56 years of age and older who received COMIRNATY Original/Omicron BA.4-5 as a booster dose in Study C4591044 demonstrated noninferiority of anti-Omicron BA.4-5 response among participants 18 to 55 years of age compared to participants 56 years of age and older for both GMR and difference in seroresponse rates (Table 11 and Table 12).

The study also assessed the level of NT50 of the anti-Omicron BA.4-5 and original SARS-CoV-2 strains pre-vaccination and 1 month after vaccination in participants who received COMIRNATY Original/Omicron BA.4-5 as a second booster dose (Table 13).

**Table 11: Geometric Mean Ratios – Study C4591044 – Participants With or Without Evidence of Infection - Evaluable Immunogenicity Population**

SARS-CoV-2 neutralisation assay	Sampling time point <sup>a</sup>	COMIRNATY Original/Omicron BA.4-5 C4591044		COMIRNATY (tozinameran) Subset of C4591031		Age group comparison		Vaccine group comparison $\geq 56$ years
		18 - 55 years of age		$\geq 56$ years of age		$\geq 56$ years of age		COMIRNATY Original/Omicron BA.4-5 18 - 55 years / $\geq 56$ years of age
		n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>e</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>e</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>e</sup> )	GMR <sup>d</sup> (95% CI <sup>d</sup> )
Omicron BA.4-5 - NT50 (titre) <sup>e</sup>	1 month	297	4455.9 (3851.7, 5154.8)	284	4158.1 (3554.8, 4863.8)	282	938.9 (802.3, 1098.8)	0.98 (0.83, 1.16) <sup>f</sup>
Reference strain - NT50 (titre) <sup>e</sup>	1 month	-	-	286	16250.1 (14499.2, 18212.4)	289	10415.5 (9366.7, 11581.8)	1.38 (1.22, 1.56) <sup>h</sup>

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titre; LLOQ = lower limit of quantitation; LS = least square; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

a. Protocol-specified timing for blood sample collection.

b. n = Number of participants with valid and determinate assay results for the specified assay at the given sampling time point.

c. GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .

d. GMRs and 2-sided 95% CIs were calculated by exponentiating the difference of LS means and corresponding CIs based on analysis of logarithmically transformed neutralising titres using a linear regression model with terms of baseline neutralising titre (log scale) and vaccine group or age group.

e. SARS-CoV-2 NT50 were determined using a validated 384-well assay platform (original strain [USA-WA1/2020, isolated in January 2020] and Omicron B.1.1.529 subvariant BA.4/BA.5).

f. Noninferiority is declared if the lower bound of the 2-sided 95% CI for the GMR is greater than 0.67.

g. Superiority is declared if the lower bound of the 2-sided 95% CI for the GMR is greater than 1.

h. Noninferiority is declared if the lower bound of the 2-sided 95% CI for the GMR is greater than 0.67 and the point estimate of the GMR is  $\geq 0.8$ .

**Table 12: Difference in Percentages of Participants with Seroresponse of COMIRNATY Original/Omicron BA.4-5 from Study C4591044 and COMIRNATY (tozinameran) from Subset of Study C4591031 – Participants With or Without Evidence of Infection – Evaluable Immunogenicity Population**

SARS-CoV-2 neutralisation assay	Sampling time point <sup>a</sup>	COMIRNATY Original/Omicron BA.4-5 C4591044		COMIRNATY (tozinameran) Subset of C4591031		Age group comparison		Vaccine group comparison $\geq 56$ years
		18 - 55 years of age		$\geq 56$ years of age		$\geq 56$ years of age		COMIRNATY Original/Omicron BA.4-5 18 - 55 years / $\geq 56$ years of age
		n <sup>b</sup>	N <sup>c</sup> (%) (95% CI <sup>d</sup> )	n <sup>b</sup>	N <sup>c</sup> (%) (95% CI <sup>d</sup> )	n <sup>b</sup>	N <sup>c</sup> (%) (95% CI <sup>d</sup> )	Difference <sup>e</sup> (95% CI <sup>f</sup> )
Omicron BA.4-5 - NT50 (titre) <sup>g</sup>	1 month	294	180 (61.2) (55.4, 66.8)	282	188 (66.7) (60.8, 72.1)	273	127 (46.5) (40.5, 52.6)	-3.03 (-9.68, 3.63) <sup>h</sup>

Abbreviations: CI = confidence interval; LLOQ = lower limit of quantitation; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Seroresponse is defined as achieving a  $\geq 4$ -fold rise from baseline. If the baseline measurement is below the LLOQ, a postvaccination assay result  $\geq 4 \times \text{LLOQ}$  is considered a seroresponse.

a. Protocol-specified timing for blood sample collection.

b. N = Number of participants with valid and determinate assay results for the specified assay at both the prevaccination time point and the given sampling time point. This value is the denominator for the percentage calculation.

c. n = Number of participants with seroresponse for the given assay at the given sampling time point.

- d. Exact 2-sided CI, based on the Clopper and Pearson method.
- e. Difference in proportions, expressed as a percentage.
- f. 2-sided CI based on the Miettinen and Nurminen method stratified by baseline neutralising titre category (< median, ≥ median) for the difference in proportions. The median of baseline neutralising titres was calculated based on the pooled data in 2 comparator groups.
- g. SARS-CoV-2 NT50 were determined using a validated 384-well assay platform (Omicron B.1.1.529 subvariant BA.4/BA.5).
- h. Noninferiority is declared if the lower bound of the 2-sided 95% CI for the difference in percentages of participants with seroresponse is > -10%.
- i. Noninferiority is declared if the lower bound of the 2-sided 95% CI for the difference in percentages of participants with seroresponse is > -5%.

**Table 13: Geometric Mean Titres by Baseline SARS-CoV-2 Status – Subsets of Study C4591044 – Prior to and 1 month after COMIRNATY Original/Omicron BA.4-5 as a Second Booster – Participants 12 years of age and older – Evaluable Immunogenicity Population**

SARS-CoV-2 neutralisation assay	Baseline SARS-CoV-2 Status	Sampling time point <sup>a</sup>	COMIRNATY Original/Omicron BA.4-5					
			12 - 17 years of age		18 - 55 years of age		≥ 56 years of age	
			n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )
Omicron BA.4-5 - NT50 (titre) <sup>f</sup>	All	Pre-vaccination	104	1105.8 (835.1, 1464.3)	294	569.6 (471.4, 688.2)	284	458.2 (365.2, 574.8)
		1 month	105	8212.8 (6807.3, 9908.7)	297	4455.9 (3851.7, 5154.8)	284	4158.1 (3554.8, 4863.8)
	Positive <sup>d</sup>	Pre-vaccination	78	1791.1 (1379.6, 2325.3)	210	1181.4 (1005.3, 1388.3)	174	1291.7 (1027.5, 1623.8)
		1 month	79	9892.5 (8114.6, 12059.8)	213	6031.6 (5203.9, 6991.0)	176	6688.9 (5664.4, 7898.8)
	Negative <sup>e</sup>	Pre-vaccination	26	260.2 (157.1, 430.9)	84	91.9 (71.5, 118.1)	110	88.9 (69.8, 113.4)
		1 month	26	4666.1 (3096.1, 7032.2)	84	2067.7 (1530.2, 2793.9)	108	1916.2 (1489.5, 2465.1)
Reference strain - NT50 (titre) <sup>f</sup>	All	Pre-vaccination	105	6863.3 (5587.8, 8430.1)	296	4017.3 (3430.7, 4704.1)	284	3690.6 (3082.2, 4419.0)
		1 month	105	23641.3 (20473.1, 27299.8)	296	16323.3 (14686.5, 18142.6)	286	16250.1 (14499.2, 18212.4)
	Positive <sup>d</sup>	Pre-vaccination	79	8685.4 (7062.7, 10680.9)	213	7068.6 (6251.9, 7992.0)	174	8082.1 (6843.6, 9544.8)
		1 month	79	25991.8 (22377.5, 30189.8)	212	19076.6 (17056.5, 21336.0)	176	21273.3 (18604.2, 24325.3)
	Negative <sup>e</sup>	Pre-vaccination	26	3356.2 (2106.9, 5346.2)	83	942.3 (705.6, 1258.3)	110	1068.0 (835.9, 1364.6)
		1 month	26	17725.2 (12376.4, 25385.7)	84	11014.6 (8793.9, 13796.0)	110	10560.6 (8827.1, 12634.5)

Abbreviations: CI = confidence interval; GMT = geometric mean titre; LLOQ = lower limit of quantitation; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

a. Protocol-specified timing for blood sample collection.

b. n = Number of participants with valid and determinate assay results for the specified assay at the given sampling time point.

c. GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to 0.5 × LLOQ.

d. Positive N-binding antibody result at baseline, positive NAAT result at baseline, or medical history of COVID-19.

e. Negative N-binding antibody result at baseline, negative NAAT result at baseline, and no medical history of COVID-19.

f. SARS-CoV-2 NT50 were determined using a validated 384-well assay platform (original strain [USA-WA1/2020, isolated in January 2020] and Omicron B.1.1.529 subvariant BA.4/BA.5).

### ***Relative vaccine immunogenicity in participants 5 to <12 years of age– after bivalent Omicron BA.4-5 (second booster dose)***

In an analysis of a subset from Study C4591048, 103 participants 5 to <12 years of age who had previously received a 2-dose primary series and a booster dose with COMIRNATY (tozinameran) received COMIRNATY Original/Omicron BA.4-5 (5/5 micrograms) as a second booster. Results include immunogenicity data from a comparator subset of participants

5 to < 12 years of age in Study C4591007 who received 3 doses of COMIRNATY (tozinameran).

The immune response 1 month after a booster dose, COMIRNATY Original/Omicron BA.4-5 elicited generally similar Omicron BA.4-5-specific neutralising titres compared with the titres in the comparator group who received 3 doses of COMIRNATY (tozinameran). COMIRNATY Original/Omicron BA.4-5 also elicited similar reference strain-specific titres compared with the titres in the comparator group.

The vaccine immunogenicity results after a booster dose in participants 5 to < 12 years of age are presented in Table 14.

**Table 14: Study C4591048 SSD – Geometric Mean Titres, by Baseline (Dose 4 Study C4591048/Dose 3 Study C4591007) SARS-CoV-2 Status – Participants With or Without Evidence of Infection – 5 to < 12 Years of Age – Evaluable Immunogenicity Population**

			Vaccine Group (as Assigned/Randomised)			
			C4591048 SSD COMIRNATY Original/Omicron BA.4-5 5/5 mcg Dose 4 and 1 Month After Dose 4		C4591007 COMIRNATY (tozinameran) 10 mcg Dose 3 and 1 Month After Dose 3	
SARS-CoV-2 neutralisation Assay	Baseline SARS-CoV-2 Status	Sampling Time Point <sup>a</sup>	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )
Omicron BA.4-5 - NT50 (titre) <sup>f</sup>	Overall	Pre- vaccination	102	488.3 (361.9, 658.8)	112	248.3 (187.2, 329.5)
		1 Month	102	2189.9 (1742.8, 2751.7)	112	1393.6 (1175.8, 1651.7)
	Positive <sup>d</sup>	Pre- vaccination	58	1069.2 (782.4, 1461.1)	65	695.0 (538.4, 897.3)
		1 Month	58	3465.6 (2682.8, 4476.7)	65	1893.9 (1547.6, 2317.7)
	Negative <sup>e</sup>	Pre- vaccination	44	173.8 (117.3, 257.4)	47	59.8 (49.0, 73.1)
		1 Month	44	1195.8 (850.2, 1681.9)	47	905.8 (703.0, 1167.2)
Reference strain - NT50 (titre) <sup>f</sup>	Overall	Pre- vaccination	102	2904.0 (2372.6, 3554.5)	113	1323.1 (1055.7, 1658.2)
		1 Month	102	8245.9 (7108.9, 9564.9)	113	7235.1 (6331.5, 8267.8)
	Positive <sup>d</sup>	Pre- vaccination	58	4198.4 (3342.9, 5272.8)	66	2672.7 (2122.4, 3365.6)
		1 Month	58	9228.4 (7707.0, 11050)	66	7632.5 (6471.6, 9001.5)
	Negative <sup>e</sup>	Pre- vaccination	44	1786.4 (1305.0, 2445.5)	47	492.9 (390.9, 621.6)
		1 Month	44	7108.8 (5534.0, 9131.8)	47	6711.9 (5345.4, 8427.7)

Abbreviations: GMT = geometric mean titre; LLOQ = lower limit of quantitation; NAAT = nucleic acid amplification test; N-binding = SARS-CoV-2 nucleoprotein-binding; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

a. Protocol-specified timing for blood sample collection.

b. n = Number of participants with valid and determinate assay results for the specified assay at the given sampling time point.

c. GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .

d. For Study 6: positive N-binding antibody result at the Dose 4 visit, positive NAAT result at the Dose 4 visit, or medical history of COVID-19. For Study 3: positive N-binding antibody result at the Dose 1, 1-month post-Dose 2 (if available), or Dose 3 visit, positive NAAT result at the Dose 1, Dose 2, Dose 3, or any unscheduled illness visit up to the Dose 3 visit, or medical history of COVID-19.

e. For Study 6: negative N-binding antibody result at the Dose 4 visit, negative NAAT result at the Dose 4 visit, and no medical



history of COVID-19. For Study 3: negative N-binding antibody result at the Dose 1, 1-month post-Dose 2 (if available), and Dose 3 visits, negative NAAT result at the Dose 1, Dose 2, Dose 3, and any unscheduled illness visits up to the Dose 3 visit, and no medical history of COVID-19.

f. SARS-CoV-2 NT50 were determined using a validated 384-well assay platform (original strain [USA-WA1/2020, isolated in January 2020] and Omicron B.1.1.529 subvariant BA.4/BA.5).

### ***Immunogenicity in participants 6 months to <5 years of age – after bivalent Omicron BA.4-5 (booster dose)***

In an interim analysis of C4591048, 310 participants (92 participants 6 months to <2 years of age and 218 participants 2 years to <5 years of age) received COMIRNATY Original/Omicron BA.4-5 (1.5/1.5 micrograms) as a booster dose after receiving 3 prior doses of COMIRNATY (tozinameran), data cut-off date 3 March 2023. The average number of days between the Dose 3 to Dose 4 was approximately 174 days (range 59 – 241 days). Results include immunogenicity data from a comparator subset of participants 6 months to <5 years of age in C4591007 who received 3 doses of COMIRNATY (tozinameran).

At 1 month after a booster dose, COMIRNATY Original/Omicron BA.4-5 elicited higher Omicron BA.4-5-specific neutralising titres compared with the titres in the comparator group who received 3 doses of COMIRNATY (tozinameran). COMIRNATY Original/Omicron BA.4-5 also elicited similar reference strain-specific titres compared with the titres in the comparator group.

The vaccine immunogenicity results after a booster dose in participants 6 months to <5 years of age are presented in Table 15.

**Table 15. Study C4591048 SSB Group 2 - Geometric Mean Titres & Geometric Mean Fold Rises, by Baseline (Dose 4) SARS-CoV-2 Status – Participants With or Without Evidence of Infection – 6 Months to <5 Years of Age – Evaluable Immunogenicity Population**

SARS-CoV-2 Neutralisation Assay	Age Group	Baseline SARS-CoV-2 Status	Sampling Time Point <sup>a</sup>	Vaccine Group (as Assigned/Randomised)			
				C4591048 SSB COMIRNATY Original/Omicron BA.4-5 1.5/1.5 mcg Dose 4 and 1 Month After Dose 4			
				n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMFR <sup>c</sup> (95% CI <sup>c</sup> )
Omicron BA.4/BA.5 NT50 (titre) <sup>f</sup>	6 months to <5 years	Overall	Pre- vaccination	266	241.7 (197.3, 296.1)	266	9.3 (8.0, 10.7)
			1 month	274	2237.2 (1884.8, 2655.5)		
		Positive <sup>d</sup>	Pre- vaccination	110	1107.7 (893.5, 1373.3)	110	6.0 (4.9, 7.4)
			1 month	112	6624.5 (5587.5, 7854.0)		
		Negative <sup>e</sup>	Pre- vaccination	153	81.4 (69.2, 95.8)	153	12.5 (10.4, 15.1)
			1 month	155	1040.9 (853.4, 1269.5)		
	6 to 23 months	Overall	Pre- vaccination	74	293.9 (195.4, 441.9)	74	6.7 (5.1, 8.8)
			1 month	78	1905.1 (1328.9, 2731.2)		
		Positive <sup>d</sup>	Pre- vaccination	36	1129.2 (741.6, 1719.2)	36	5.3 (3.6, 7.6)
			1 month	36	5948.3 (4272.9, 8280.5)		

SARS-CoV-2 Neutralisation Assay	Age Group	Baseline SARS-CoV-2 Status	Sampling Time Point <sup>a</sup>	Vaccine Group (as Assigned/Randomised)			
				C4591048 SSB COMIRNATY Original/Omicron BA.4-5 1.5/1.5 mcg Dose 4 and 1 Month After Dose 4			
				n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMFR <sup>c</sup> (95% CI <sup>c</sup> )
		Negative <sup>e</sup>	Pre- vaccination	36	75.3 (55.4, 102.5)	36	8.4 (5.6, 12.6)
			1 month	38	697.0 (446.5, 1087.9)		
	2 to <5 years	Overall	Pre- vaccination	192	224.1 (177.2, 283.5)	192	10.5 (8.9, 12.5)
			1 month	196	2384.9 (1965.4, 2893.9)		
		Positive <sup>d</sup>	Pre- vaccination	74	1097.4 (852.3, 1413.1)	74	6.4 (5.0, 8.3)
			1 month	76	6971.2 (5705.2, 8518.2)		
		Negative <sup>e</sup>	Pre- vaccination	117	83.4 (68.8, 101.1)	117	14.2 (11.6, 17.4)
			1 month	117	1185.7 (953.0, 1475.2)		
Reference strain - NT50 (titre) <sup>f</sup>	6 months to <5 years	Overall	Pre- vaccination	266	1721.9 (1491.0, 1988.4)	266	4.3 (3.8, 4.8)
			1 month	274	7409.3 (6649.4, 8256.1)		
		Positive <sup>d</sup>	Pre- vaccination	110	3549.5 (2922.2, 4311.5)	110	2.8 (2.5, 3.2)
			1 month	112	10080.7 (8720.9, 11652.5)		
		Negative <sup>e</sup>	Pre- vaccination	153	1046.7 (888.7, 1232.7)	153	5.6 (4.8, 6.6)
			1 month	155	5910.6 (5082.7, 6873.4)		
	6 to 23 months	Overall	Pre- vaccination	74	1688.3 (1271.6, 2241.6)	74	3.7 (2.9, 4.7)
			1 month	78	6312.0 (5143.6, 7746.0)		
		Positive <sup>d</sup>	Pre- vaccination	36	3050.0 (2123.6, 4380.5)	36	2.6 (2.0, 3.4)
			1 month	36	7979.2 (5949.9, 10700.7)		
		Negative <sup>e</sup>	Pre- vaccination	36	981.3 (676.5, 1423.2)	36	4.9 (3.4, 7.1)
			1 month	38	5064.3 (3722.6, 6889.5)		
	2 to <5 years	Overall	Pre- vaccination	192	1734.9 (1466.0, 2053.3)	192	4.5 (3.9, 5.2)
			1 month	196	7897.3 (6952.0, 8971.2)		
		Positive <sup>d</sup>	Pre- vaccination	74	3821.3 (3025.6, 4826.3)	74	2.9 (2.5, 3.5)
			1 month	76	11261.1 (9586.2, 13228.7)		
		Negative <sup>e</sup>	Pre- vaccination	117	1067.7 (888.4, 1283.2)	117	5.8 (4.9, 6.9)
			1 month	117	6214.9 (5218.5, 7401.6)		

SARS-CoV-2 Neutralisation Assay	Age Group	Baseline SARS-CoV-2 Status	Sampling Time Point <sup>a</sup>	Vaccine Group (as Assigned/Randomised)			
				C4591048 SSB COMIRNATY Original/Omicron BA.4-5 1.5/1.5 mcg Dose 4 and 1 Month After Dose 4			
				n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMFR <sup>c</sup> (95% CI <sup>c</sup> )

Abbreviations: GMT = geometric mean titre; GMFR = geometric mean fold rise; LLOQ = lower limit of quantitation; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Substudy B Group 2 includes participants  $\geq 6$  months to  $< 5$  years of age who received 3 doses of BNT162b2 3  $\mu$ g 60 to 240 days prior to enrolment.

- Protocol-specified timing for blood sample collection.
- n = Number of participants with valid and determinate assay results for the specified assay at the given sampling time point.
- GMTs/GMFRs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres/fold rises and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times$  LLOQ.
- For C4591048 Substudy B Group 2: positive N-binding antibody result at Dose 4 visit, positive NAAT result at Dose 4 visit, or medical history of COVID-19.
- For C4591048 Substudy B Group 2: negative N-binding antibody result at Dose 4 visit, negative NAAT result at Dose 4 visit, and no medical history of COVID-19.
- SARS-CoV-2 NT50 were determined using a validated 384-well assay platform (original strain [USA-WA1/2020, isolated in January 2020] and Omicron B.1.1.529 subvariant BA.4/BA.5).

Overall, 22 cases of COVID-19 were reported in participants  $\geq 6$  months to  $< 5$  years (9 cases in  $\geq 6$  months to  $< 2$  years, 13 cases in  $\geq 2$  to  $< 5$  years) who received a fourth dose of bivalent BNT162b2 (Original/Omi BA.4/BA.5). Lineage for the cases was determined as follows:

Among the determinate and quantifiable sequence, lineage for the cases was identified as XBB.1.5 (Omicron) (n=4), BQ.1 (Omicron) (n= 2), BA.5.1.3 (Omicron) (n=1), BL.1 (Omicron) (n= 1), BQ.1.1 (Omicron) (n=1), BQ.1.1.18 (Omicron) (n=1), BQ.1.1.35 (Omicron) (n=1), BQ.1.25 (Omicron) (n=1), CQ.1.1 (Omicron) (n=1), FD.2 (Omicron) (n=1), XBB.1.5.14 (Omicron) (n=1), XBB.1.5.34 (Omicron) (n=1), and unknown due to insufficient quantity (n=5) or not sequenced (n=1).

### ***Immunogenicity in participants 6 months to $< 5$ years of age – bivalent Omicron BA.4-5 3-dose primary course***

C4591048 Substudy A (SSA), a phase I, randomised, single-blinded, dose-finding study included evaluation of 1-month post-dose 3 safety and immunogenicity following 3 doses with COMIRNATY Original/Omicron BA.4-5 (1.5/1.5 micrograms) administered on a 0-, 3-, and 11-week schedule in participants 6 months to  $< 5$  years of age.

Descriptive immunogenicity analyses were performed to characterise Omicron BA.4/BA.5 and reference strain neutralisation responses following 3 doses with COMIRNATY Original/Omicron BA.4-5.

The vaccine immunogenicity results after a 3-dose series in participants 6 months to  $< 5$  years of age are presented in Table 16.

**Table 16. Study C4591048 SSA - Geometric Mean Titres & Geometric Mean Fold Rises at 1 month after vaccination course (3-doses), by Baseline SARS-CoV-2 Status – Participants With or Without Evidence of Infection – 6 Months to <5 Years of Age – Evaluable Immunogenicity Population**

SARS-CoV-2 Neutralisation Assay	Age Group	Baseline SARS-CoV-2 Status	Sampling Time Point <sup>a</sup>	Vaccine Group (as Randomised)			
				C4591048 SSA COMIRNATY Original/Omicron BA.4-5 1.5/1.5 micrograms 3-doses and 1 Month After Dose 3			
				n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMFR <sup>c</sup> (95% CI <sup>c</sup> )
Omicron BA.4/BA.5 NT50 (titre)	6 months to <5 years	Overall	Pre-vaccination	35	492.6 (247.2, 981.4)	25	15.5 (8.1, 29.5)
			1 month	29	10452.6 (7091.3, 15407.2)		
		Positive <sup>d</sup>	Pre-vaccination	28	950.6 (495.1, 1825.3)	21	10.0 (5.7, 17.3)
			1 month	21	12700.1 (7861.5, 20516.7)		
		Negative <sup>e</sup>	Pre-vaccination	7	35.5 (35.5, 35.5)	4	158.7 (31.2, 808.4)
			1 month	5	6659.9 (2017.3, 21986.4)		
Reference strain - NT50 (titre)	6 months to <5 years	Overall	Pre-vaccination	35	234.4 (147.3, 373.1)	25	22.7 (12.8, 40.3)
			1 month	29	6674.9 (4222.2, 10552.2)		
		Positive <sup>d</sup>	Pre-vaccination	28	322.3 (195.8, 530.5)	21	18.7 (10.2, 34.3)
			1 month	21	8370.1 (4725.8, 14824.8)		
		Negative <sup>e</sup>	Pre-vaccination	7	65.6 (32.7, 131.8)	4	64.1 (8.1, 507.3)
			1 month	5	4860.4 (2073.5, 11393.2)		

Abbreviations: GMT = geometric mean titre; GMFR = geometric mean fold rise; LLOQ = lower limit of quantitation; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

a. Protocol-specified timing for blood sample collection.

b. n = Number of participants with valid and determinate assay results for the specified assay at the given sampling time point.

c. GMTs/GMFRs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres/fold rises and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to 0.5 × LLOQ.

d. Positive N-binding antibody result at Dose 1, positive NAAT result at Dose 1, or medical history of COVID-19.

e. Negative N-binding antibody result at Dose 1, negative NAAT result at Dose 1, and no medical history of COVID-19.

For C4591048 SSA: Evaluable immunogenicity (3-Dose) population.

## COMIRNATY (tozinameran)

Study C4591001 is a multicentre, multinational, Phase 1/2/3 randomised, placebo-controlled, observer-blind dose-finding, vaccine candidate selection and efficacy study in participants 12 years of age and older. Randomisation was stratified by age: 12 to 15 years of age, 16 to 55 years of age, or 56 years of age and older, with a minimum of 40% of participants in the ≥56-year stratum. The study excluded participants who were immunocompromised and those who had previous clinical or microbiological diagnosis of COVID-19. Participants with pre-existing stable disease, defined as disease not requiring significant change in therapy or hospitalisation for worsening disease during the 6 weeks before enrolment, were included as were participants with known stable infection with HIV, hepatitis C virus (HCV) or hepatitis B virus (HBV).

### ***Efficacy in participants 16 years of age and older – after 2 doses***

In the Phase 2/3 portion of Study C4591001, based on data accrued through 14 November 2020, approximately 44,000 participants were randomised equally and were to receive 2 doses of COMIRNATY (tozinameran) or placebo. The efficacy analyses included participants that received their second vaccination within 19 to 42 days after their first vaccination. The majority (93.1%) of vaccine recipients received the second dose 19 days to 23 days after Dose 1. Participants are planned to be followed for up to 24 months after Dose 2, for assessments of safety and efficacy against COVID-19. In the clinical study, participants were required to observe a minimum interval of 14 days before and after administration of an influenza vaccine in order to receive either placebo or COMIRNATY (tozinameran). In the clinical study, participants were required to observe a minimum interval of 60 days before or after receipt of blood/plasma products or immunoglobulins through to conclusion of the study in order to receive either placebo or COMIRNATY (tozinameran).

The population for the analysis of the primary efficacy endpoint included, 36,621 participants 12 years of age and older (18,242 in the COMIRNATY (tozinameran) group and 18,379 in the placebo group) who did not have evidence of prior infection with SARS-CoV-2 through 7 days after the second dose. In addition, 134 participants were between the ages of 16 to 17 years of age (66 in the COMIRNATY (tozinameran) group and 68 in the placebo group) and 1616 participants 75 years of age and older (804 in the COMIRNATY (tozinameran) group and 812 in the placebo group).

At the time of the primary efficacy analysis, participants had been followed for symptomatic COVID-19 for in total 2,214 person-years for the COMIRNATY (tozinameran) group and in total 2,222 person-years for the placebo group.

There were no meaningful clinical differences in overall vaccine efficacy in participants who were at risk of severe COVID-19 including those with 1 or more comorbidities that increase the risk of severe COVID-19 (e.g. asthma, body mass index (BMI)  $\geq 30$  kg/m<sup>2</sup>, chronic pulmonary disease, diabetes mellitus, hypertension).

COMIRNATY (tozinameran) efficacy information is presented in Table 17.

**Table 17: Vaccine efficacy – First COVID-19 occurrence from 7 days after Dose 2, by age subgroup – participants without evidence of infection prior to 7 days after Dose 2 – evaluable efficacy (7 days) population**

<b>First COVID-19 occurrence from 7 days after Dose 2 in participants without evidence of prior SARS-CoV-2 infection*</b>			
<b>Subgroup</b>	<b>COMIRNATY (tozinameran) N<sup>a</sup> = 18,198 Cases n1<sup>b</sup> Surveillance time<sup>c</sup> (n2<sup>d</sup>)</b>	<b>Placebo N<sup>a</sup> = 18,325 Cases n1<sup>b</sup> Surveillance time<sup>c</sup> (n2<sup>d</sup>)</b>	<b>Vaccine efficacy % (95% CI)<sup>f</sup></b>
All participants <sup>e</sup>	8 2.214 (17,411)	162 2.222 (17,511)	95.0 (90.0, 97.9)
16 to 64 years	7 1.706 (13,549)	143 1.710 (13,618)	95.1 (89.6, 98.1)
65 years and older	1 0.508 (3848)	19 0.511 (3880)	94.7 (66.7, 99.9)
65 to 74 years	1 0.406 (3074)	14 0.406 (3095)	92.9 (53.1, 99.8)
75 years and older	0 0.102 (774)	5 0.106 (785)	100.0 (-13.1, 100.0)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 [\*Case definition: (at least 1 of) fever, new or increased cough, new or increased shortness of breath, chills, new or increased muscle pain, new loss of taste or smell, sore throat, diarrhoea or vomiting.]

\* Participants who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by nucleic acid amplification tests (NAAT) [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of participants in the specified group.

b. n1 = Number of participants meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of participants at risk for the endpoint.

e. No confirmed cases were identified in adolescents 12 to 15 years of age.

f. Two-sided confidence interval (CI) for vaccine efficacy (VE) is derived based on the Clopper and Pearson method adjusted to the surveillance time. CI not adjusted for multiplicity.

In the second primary analysis, efficacy of COMIRNATY (tozinameran) in preventing first COVID-19 occurrence from 7 days after Dose 2 compared to placebo was 94.6% (95% credible interval of 89.9% to 97.3%) in participants 16 years of age and older with or without evidence of prior infection with SARS-CoV-2.

Additionally, subgroup analyses of the primary efficacy endpoint showed similar efficacy point estimates across genders, ethnic groups, and participants with medical comorbidities associated with high risk of severe COVID-19.

Updated efficacy analyses were performed with additional confirmed COVID-19 cases accrued during blinded placebo-controlled follow-up through 13 March 2021, representing up to 6 months of follow-up after Dose 2 for participants in the efficacy population.

The updated vaccine efficacy information is presented in Table 18.

**Table 18: Vaccine efficacy – First COVID-19 occurrence from 7 days after Dose 2, by age subgroup – participants without evidence of infection prior to 7 days after Dose 2 – evaluable efficacy (7 days) population during the placebo-controlled follow-up period**

<b>First COVID-19 occurrence from 7 days after Dose 2 in participants without evidence of prior SARS-CoV-2 infection*</b>			
<b>Subgroup</b>	<b>COMIRNATY (tozinameran) N<sup>a</sup>=20,998 Cases n1<sup>b</sup> Surveillance Time<sup>c</sup> (n2<sup>d</sup>)</b>	<b>Placebo N<sup>a</sup>=21,096 Cases n1<sup>b</sup> Surveillance Time<sup>c</sup> (n2<sup>d</sup>)</b>	<b>Vaccine efficacy % (95% CI)<sup>e</sup></b>
All participants <sup>f</sup>	77 6.247 (20,712)	850 6.003 (20,713)	91.3 (89.0, 93.2)
16 to 64 years	70 4.859 (15,519)	710 4.654 (15,515)	90.6 (87.9, 92.7)
65 years and older	7 1.233 (4192)	124 1.202 (4226)	94.5 (88.3, 97.8)
65 to 74 years	6 0.994 (3350)	98 0.966 (3379)	94.1 (86.6, 97.9)
75 years and older	1 0.239 (842)	26 0.237 (847)	96.2 (76.9, 99.9)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhoea; vomiting).

\* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = Number of participants in the specified group.

b. n1 = Number of participants meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of participants at risk for the endpoint.

e. Two-sided confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted to the surveillance time.

- f. Included confirmed cases in participants 12 to 15 years of age: 0 in the COMIRNATY (tozinameran) group (both without and with or without evidence of prior SARS-CoV-2 infection); 16 and 18 in the placebo group (without and with or without evidence of prior SARS-CoV-2 infection, respectively).

### ***Efficacy against severe COVID-19 in participants 12 years of age and older – after 2 doses***

As of 13 March 2021, vaccine efficacy against severe COVID-19 is presented only for participants with or without prior SARS-CoV-2 infection (Table 19) as the COVID-19 case counts in participants without prior SARS-CoV-2 infection were the same as those in participants with or without prior SARS-CoV-2 infection in both the COMIRNATY (tozinameran) and placebo groups.

**Table 19. Vaccine Efficacy – First Severe COVID-19 Occurrence in Participants With or Without\* Prior SARS-CoV-2 Infection Based on Food and Drug Administration (FDA)<sup>†</sup> Definition After Dose 1 or From 7 Days After Dose 2 in the Placebo-Controlled Follow-up**

	<b>COMIRNATY (tozinameran) Cases n1<sup>a</sup> Surveillance Time (n2<sup>b</sup>)</b>	<b>Placebo Cases n1<sup>a</sup> Surveillance Time (n2<sup>b</sup>)</b>	<b>Vaccine Efficacy % (95% CI)<sup>c</sup></b>
After Dose 1 <sup>d</sup>	1 8.439 <sup>e</sup> (22,505)	30 8.288 <sup>e</sup> (22,435)	96.7 (80.3, 99.9)
7 days after Dose 2 <sup>f</sup>	1 6.522 <sup>g</sup> (21,649)	21 6.404 <sup>g</sup> (21,730)	95.3 (70.9, 99.9)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhoea; vomiting).

\* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

<sup>†</sup> Severe illness from COVID-19 as defined by FDA is confirmed COVID-19 and presence of at least 1 of the following:

- Clinical signs at rest indicative of severe systemic illness (respiratory rate  $\geq 30$  breaths per minute, heart rate  $\geq 125$  beats per minute, saturation of oxygen  $\leq 93\%$  on room air at sea level, or ratio of arterial oxygen partial pressure to fractional inspired oxygen  $< 300$  mm Hg);
- Respiratory failure [defined as needing high-flow oxygen, noninvasive ventilation, mechanical ventilation or extracorporeal membrane oxygenation (ECMO)];
- Evidence of shock (systolic blood pressure  $< 90$  mm Hg, diastolic blood pressure  $< 60$  mm Hg, or requiring vasopressors);
- Significant acute renal, hepatic, or neurologic dysfunction;
- Admission to an Intensive Care Unit;
- Death.

a. n1 = Number of participants meeting the endpoint definition.

b. n2 = Number of participants at risk for the endpoint.

c. Two-side confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted to the surveillance time.

d. Efficacy assessed based on the Dose 1 all available efficacy (modified intention-to-treat) population that included all randomised participants who received at least 1 dose of study intervention.

e. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from Dose 1 to the end of the surveillance period.

f. Efficacy assessed based on the evaluable efficacy (7 Days) population that included all eligible randomised participants who receive all dose(s) of study intervention as randomised within the predefined window, have no other important protocol deviations as determined by the clinician

g. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

### ***Efficacy and immunogenicity in adolescents 12 to 15 years of age – after 2 doses***

An analysis of Study C4591001 has been performed in adolescents 12 to 15 years of age up to a data cut-off date of 13 March 2021.

In an analysis of Study C4591001 in adolescents 12 to 15 years of age without evidence of prior infection, there were no cases in 1005 participants who received the vaccine and 16 cases out of 978 who received placebo. The point estimate for efficacy is 100% (95% confidence interval 75.3, 100.0). In participants with or without evidence of prior infection there were 0 cases in the 1119 who received vaccine and 18 cases in 1110 participants who received placebo. This also indicates the point estimate for efficacy is 100% (95% confidence interval 78.1, 100.0). No cases of severe disease occurred in adolescents.

In Study C4591001, an analysis of SARS-CoV-2 neutralising titres in a randomly selected subset of participants was performed to demonstrate non-inferior immune responses (within 1.5-fold) comparing adolescents 12 to 15 years of age to participants 16 to 25 years of age who had no serological or virological evidence of past SARS-CoV-2 infection. The immune response to COMIRNATY (tozinameran) in adolescents 12 to 15 years of age (n = 190) was non-inferior to the immune response in participants 16 to 25 years of age (n = 170), based on results for SARS-CoV-2 neutralising titres at 1 month after Dose 2. The geometric mean titres (GMT) ratio of the adolescents 12 to 15 years of age group to the participants 16 to 25 years of age group was 1.76, with a 2-sided 95% CI of 1.47 to 2.10, meeting the 1.5-fold non-inferiority criterion (the lower bound of the 2-sided 95% CI for the geometric mean ratio [GMR] > 0.67).

An updated efficacy analysis of Study C4591001 has been performed in approximately 2,260 adolescents 12 to 15 years of age evaluating confirmed COVID-19 cases accrued up to a data cut-off date of 2 September 2021, representing up to 6 months of follow-up after Dose 2 for participants in the efficacy population. The dominant SARS-CoV-2 variant at the time of the efficacy study was B.1.1.7 (Alpha).

The updated vaccine efficacy information in adolescents 12 to 15 years of age is presented in Table 20.

**Table 20: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2: Without Evidence of Infection and With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Blinded Placebo-Controlled Follow-up Period, Adolescents 12 To 15 Years of Age Evaluable Efficacy (7 Days) Population**

First COVID-19 occurrence from 7 days after Dose 2 in adolescents 12 to 15 years of age without evidence of prior SARS-CoV-2 infection*			
	COMIRNATY (tozinameran) N <sup>a</sup> =1057 Cases n <sup>1b</sup> Surveillance Time <sup>c</sup> (n <sup>2d</sup> )	Placebo N <sup>a</sup> =1030 Cases n <sup>1b</sup> Surveillance Time <sup>c</sup> (n <sup>2d</sup> )	Vaccine Efficacy % (95% CI <sup>e</sup> )
Adolescents 12 to 15 years of age	0 0.343 (1043)	28 0.322 (1019)	100.0 (86.8, 100.0)
First COVID-19 occurrence from 7 days after Dose 2 in adolescents 12 to 15 years of age with or without evidence of prior SARS-CoV-2 infection			
	COMIRNATY (tozinameran) N <sup>a</sup> =1119 Cases n <sup>1b</sup> Surveillance Time <sup>c</sup> (n <sup>2d</sup> )	Placebo N <sup>a</sup> =1109 Cases n <sup>1b</sup> Surveillance Time <sup>c</sup> (n <sup>2d</sup> )	Vaccine Efficacy % (95% CI <sup>e</sup> )
Adolescents 12 to 15 years of age	0 0.362 (1098)	30 0.345 (1088)	100.0 (87.5, 100.0)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhoea; vomiting).

\* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had



negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

- a. N = Number of participants in the specified group.
- b. n1 = Number of participants meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.
- d. n2 = Number of participants at risk for the endpoint.
- e. Two-sided confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted for surveillance time.

### ***Efficacy in children 5 to <12 years of age – after 2 doses***

A initial descriptive efficacy analysis of Study C4591007 has been performed in 1,968 children 5 to 11 years of age without evidence of infection prior to 7 days after Dose 2. This analysis evaluated confirmed symptomatic COVID-19 cases accrued up to a data cut-off date of 8 October 2021.

The initial descriptive vaccine efficacy results in children 5 to 11 years of age without evidence of prior SARS-CoV-2 infection are presented in Table 21. None of the cases accrued met criteria for severe COVID-19 or multisystem inflammatory syndrome in children (MIS-C). No cases of COVID-19 were observed in either the vaccine group or the placebo group in participants with evidence of prior SARS-CoV-2 infection.

**Table 21: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2: Without Evidence of Infection Prior to 7 Days After Dose 2 – Phase 2/3 – Children 5 To 11 Years of Age Evaluable Efficacy Population**

First COVID-19 occurrence from 7 days after Dose 2 in children 5 to 11 years of age without evidence of prior SARS-CoV-2 infection*			
	COMIRNATY <sup>±</sup> (tozinameran) 10 microgram/dose N <sup>a</sup> =1305 Cases n1 <sup>b</sup> Surveillance Time <sup>c</sup> (n2 <sup>d</sup> )	Placebo N <sup>a</sup> =663 Cases n1 <sup>b</sup> Surveillance Time <sup>c</sup> (n2 <sup>d</sup> )	Vaccine Efficacy % (95% CI)
Children 5 to 11 years of age	3 0.322 (1273)	16 0.159 (637)	90.7 (67.7, 98.3)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhoea; vomiting).

\* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

± Pfizer-BioNTech COVID-19 Vaccine (10 micrograms modRNA).

- a. N = Number of participants in the specified group.
- b. n1 = Number of participants meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.
- d. n2 = Number of participants at risk for the endpoint.

Prespecified hypothesis-driven efficacy analysis was performed with additional confirmed COVID-19 cases accrued during blinded placebo-controlled follow-up, representing up to 6 months after Dose 2 in the efficacy population.

In the efficacy analysis of Study C4591007 in children 5 to 11 years of age without evidence of prior infection, there were 10 cases out of 2,703 participants who received the vaccine and 42 cases out of 1,348 participants who received placebo. The point estimate for efficacy is 88.2% (95% CI: 76.2, 94.7). In participants with or without evidence of prior infection there were 12 cases in the 3,018 who received vaccine and 42 cases in 1,511 participants who received placebo. The point estimate for efficacy is 85.7% (95% CI: 72.4, 93.2).

### ***Immunogenicity in children 5 to <12 years of age – after 2 doses***

Study C4591007 is a Phase 1/2/3 study comprised of an open-label vaccine dose-finding portion (Phase 1) and a multicentre, multinational, randomised, saline placebo-controlled, observer-blind efficacy portion (Phase 2/3) that has enrolled participants 5 to <12 years of age.

In C4591007, an analysis of SARS-CoV-2 50% neutralising titres (NT50) 1 month after Dose 2 in a randomly selected subset of participants demonstrated effectiveness by immunobridging of immune responses comparing children 5 to <12 years of age in the Phase 2/3 part of Study C4591007 to participants 16 to 25 years of age in the Phase 2/3 part of Study C4591001 who had no serological or virological evidence of past SARS-CoV-2 infection up to 1 month after Dose 2, meeting the prespecified immunobridging criteria for both the geometric mean ratio (GMR) and the seroresponse difference with seroresponse defined as achieving at least 4-fold rise in SARS-CoV-2 NT50 from baseline (before Dose 1).

The ratio of the SARS-CoV-2 NT50 in children 5 to <12 years of age to that of young adults 16 to 25 years of age was 1.04 (2-sided 95% CI: 0.93, 1.18), as presented in Table 22.

**Table 22: Summary of geometric mean ratio for 50% neutralising titre – Comparison of children 5 to <12 years of age (Study C4591007) to participants 16 to 25 years of age (Study C4591001) – participants without\* evidence of infection up to 1 month after Dose 2 – evaluable immunogenicity population**

		COMIRNATY (tozinameran)		5 to <12 years/ 16 to 25 years	
		10 microgram/dose 5 to <12 years n <sup>a</sup> =264	30 microgram/dose 16 to 25 years n <sup>a</sup> =253		
Assay	Time point <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	GMT <sup>c</sup> (95% CI <sup>c</sup> )	GMR <sup>d</sup> (95% CI <sup>d</sup> )	Met immunobridging objective <sup>e</sup> (Y/N)
SARS-CoV-2 neutralisation assay - NT50 (titre) <sup>f</sup>	1 month after Dose 2	1197.6 (1106.1, 1296.6)	1146.5 (1045.5, 1257.2)	1.04 (0.93, 1.18)	Y

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titre; LLOQ = lower limit of quantitation; NAAT = nucleic acid amplification test; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

\*Participants who had no serological or virological evidence (up to 1 month post-Dose 2 blood sample collection) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and 1 month after Dose 2, SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2, and negative NAAT (nasal swab) at any unscheduled visit up to 1 month after Dose 2 blood collection) and had no medical history of COVID-19 were included in the analysis.

- n = Number of participants with valid and determinate assay results for the specified assay at the given dose/sampling time point.
- Protocol-specified timing for blood sample collection.
- GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .

- d. GMRs and 2-sided 95% CIs were calculated by exponentiating the mean difference of the logarithms of the titres (Group 1 [5 to < 12 years of age] - Group 2 [16 to 25 years of age]) and the corresponding CI (based on the Student t distribution).
- e. Immunobridging is declared if the lower bound of the 2-sided 95% CI for the GMR is greater than 0.67 and the point estimate of the GMR is  $\geq 0.8$ .
- f. SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralisation Assay. The assay uses a fluorescent reporter virus derived from the USA\_WA1/2020 strain and virus neutralisation is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralised.

Among participants without prior evidence of SARS-CoV-2 infection up to 1 month after Dose 2, 99.2% of children 5 to <12 years of age and 99.2% of participants 16 to 25 years of age had a seroresponse from before vaccination to 1 month after Dose 2. The difference in proportions of participants who had seroresponse between the 2 age groups (children – young adult) was 0.0% (2-sided 95% CI: -2.0%, 2.2%) as presented in Table 23.

**Table 23: Difference in percentages of participants with seroresponse – participants without evidence of infection up to 1 month after Dose 2 – immunobridging subset – Phase 2/3 – comparison of 5 to <12 years of age to Study C4591001 Phase 2/3 16 to 25 years of age – evaluable immunogenicity population**

		COMIRNATY (tozinameran)		5 to <12 years/ 16 to 25 years	
		10 microgram/dose 5 to <12 years N <sup>a</sup> =264	30 microgram/dose 16 to 25 years N <sup>a</sup> =253		
Assay	Time point <sup>b</sup>	n <sup>c</sup> (%) (95% CI <sup>d</sup> )	n <sup>c</sup> (%) (95% CI <sup>d</sup> )	Difference % <sup>e</sup> (95% CI <sup>f</sup> )	Met immunobridging objective <sup>g</sup> (Y/N)
SARS-CoV-2 neutralisation assay – NT50 (titre) <sup>h</sup>	1 month after Dose 2	262 (99.2) (97.3, 99.9)	251 (99.2) (97.2, 99.9)	0.0 (-2.0, 2.2)	Y

Abbreviations: LLOQ = lower limit of quantitation; NAAT = nucleic acid amplification test; N-binding = SARS-CoV-2 nucleoprotein-binding; NT50 = 50% neutralising titre 50; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Seroresponse is defined as achieving a  $\geq 4$ -fold rise from baseline (before Dose 1). If the baseline measurement is below the LLOQ, a postvaccination assay result  $\geq 4 \times$  LLOQ is considered a seroresponse.

Note: Participants who had no serological or virological evidence (up to 1 month post-Dose 2 blood sample collection) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and 1 month after Dose 2, SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2, and negative NAAT (nasal swab) at any unscheduled visit up to 1 month after Dose 2 blood collection) and had no medical history of COVID-19 were included in the analysis.

- a. N = number of participants with valid and determinate assay results both before vaccination and at 1 month after Dose 2. These values are the denominators for the percentage calculations.
- b. Protocol-specified timing for blood sample collection.
- c. n = Number of participants with seroresponse for the given assay at the given dose/sampling time point.
- d. Exact 2-sided CI based on the Clopper and Pearson method.
- e. Difference in proportions, expressed as a percentage (Group 1 [5 to < 12 years of age] – Group 2 [16 to 25 years of age]).
- f. 2-Sided CI, based on the Miettinen and Nurminen method for the difference in proportions, expressed as a percentage.
- g. Immunobridging is declared if the lower bound of the 2-sided 95% CI for the difference in proportions is greater than -10.0%.
- h. SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralisation Assay. The assay uses a fluorescent reporter virus derived from the USA\_WA1/2020 strain and virus neutralisation is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralised.

## ***Efficacy and immunogenicity in infants and children 6 months to <5 years of age – 3-dose primary course***

Effectiveness in individuals 6 months to <5 years of age is based on a comparison of efficacy against symptomatic COVID-19 comparing to placebo and immune responses in this age group to individuals 16 through 25 years of age.

### ***Efficacy in infants and children 6 months to <5 years of age – after 3 doses***

The efficacy analysis of Study C4591007 was performed across the combined population of participants 6 months to <5 years of age based on cases confirmed among 873 participants in the COMIRNATY (tozinameran) group and 381 participants in the placebo group (2:1 randomisation ratio) who received all 3 doses of study intervention during the blinded follow-up period when the Omicron variant of SARS-CoV-2 (BA.2) was the predominant variant in circulation (data cutoff date of 17 June 2022).

The vaccine efficacy results after Dose 3 in participants 6 months to <5 years of age are presented in Table 24.

**Table 24: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 3 – Blinded Follow-Up Period – Participants Without Evidence of Infection and Participants With or Without Evidence of Infection Prior to 7 Days After Dose 3 – Phase 2/3 – 6 Months to <5 Years of Age – Evaluable Efficacy (3-Dose) Population**

<b>First COVID-19 occurrence from 7 days after Dose 3 in participants without evidence of prior SARS-CoV-2 infection*</b>			
<b>Subgroup</b>	<b>COMIRNATY (tozinameran) 3 micrograms/Dose N<sup>a</sup>=873 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Placebo N<sup>a</sup>=381 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Vaccine Efficacy % (95% CI<sup>e</sup>)</b>
6 months to <5 years <sup>e</sup>	13 0.124 (794)	21 0.054 (351)	73.2 (43.8, 87.6)
2 to <5 years	9 0.081 (498)	13 0.033 (204)	71.8 (28.6, 89.4)
6 months to <2 years	4 0.042 (296)	8 0.020 (147)	75.8 (9.7, 94.7)
<b>First COVID-19 occurrence from 7 days after Dose 3 in participants with or without evidence of prior SARS-CoV-2 infection</b>			
<b>Subgroup</b>	<b>COMIRNATY (tozinameran) 3 micrograms/Dose N<sup>a</sup>=1294 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Placebo N<sup>a</sup>=612 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Vaccine Efficacy % (95% CI<sup>e</sup>)</b>
6 months to <5 years <sup>e</sup>	14 0.149 (981)	23 0.067 (459)	72.5 (44.3, 86.9)
2 to <5 years	10 0.100 (639)	15 0.044 (286)	70.7 (30.3, 88.2)
6 months to <2 years	4 0.048 (342)	8 0.023 (173)	76.2 (11.1, 94.8)

---

Abbreviations: NAAT = nucleic acid amplification test; N-binding = SARS-CoV-2 nucleoprotein-binding; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

- \* Participants who had no serological or virological evidence (prior to 7 days after receipt of Dose 3) of past SARS-CoV-2 infection (i.e., negative N-binding antibody [serum] result at Dose 1, 1 month post-Dose 2 (if available), Dose 3 (if available) visits, SARS-CoV-2 not detected by NAAT [nasal swab] at Dose 1, Dose 2, and Dose 3 study visits, and a negative NAAT [nasal swab] result at any unscheduled visit prior to 7 days after receipt of Dose 3) and had no medical history of COVID-19 were included in the analysis.
- a. N = number of participants in the specified group.
- b. n1 = Number of participants meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 3 to the end of the surveillance period.
- d. n2 = Number of participants at risk for the endpoint.
- e. Two-sided 95% confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.

Analysis of COVID-19 cases that excluded those involving coinfection with other respiratory pathogens did not meaningfully impact the estimated vaccine efficacy in this population.

Among participants 2 to <5 years of age, severe COVID-19 criteria (as described in the protocol, based on FDA definition and modified for children) were fulfilled for 9 cases [6 COMIRNATY (tozinameran) and 3 placebo] of which 5 of the 6 cases in the COMIRNATY (tozinameran) group fulfilled a single criterion of increased heart rate or respiratory rate and all 3 cases in the placebo group fulfilled a single criterion of increased heart rate or decreased peripheral oxygen saturation. None of the cases accrued met criteria for multisystem inflammatory syndrome in children (MIS-C).

Among participants 6 months to <2 years of age, severe COVID-19 criteria were fulfilled for 3 cases [2 COMIRNATY (tozinameran) and 1 placebo] of which 1 of the 2 cases in the COMIRNATY (tozinameran) group fulfilled a single criterion of increased heart rate (152 bpm) and 1 case in the placebo group fulfilled a single criterion of increased heart rate (172 bpm). None of the cases accrued met criteria for MIS-C.

Vaccine efficacy analyses were associated with wide confidence intervals. In addition, the preliminary nature of the data (prespecified number of cases not yet reached in Study C4591007) may preclude any definitive vaccine efficacy conclusions.

Dosing intervals: In the evaluable efficacy population, there was a wide dosing interval range between COMIRNATY Dose 2 and Dose 3, for participants 2 to <5 years of age was 6.0 to 34.1 weeks with a median interval of 11.0 weeks and for participants 6 months to <2 years of age was 8.0 to 31.9 weeks with a median interval of 16.0 weeks.

### ***Immunogenicity in children 2 to <5 years of age – after 3 doses***

Immunogenicity analyses have been performed in the immunobridging subset of 143 C4591007 participants 2 to <5 years of age without evidence of infection up to 1 month after Dose 3 based on a data cutoff date of 29 April 2022.

SARS-CoV-2 50% neutralising antibody titres (NT50) were compared between an immunogenicity subset of Phase 2/3 participants 2 to <5 years of age from C4591007 at 1 month after the 3-dose primary course and a randomly selected subset from C4591001 Phase 2/3 participants 16 to 25 years of age at 1 month after the 2-dose primary course, using a microneutralisation assay against the reference strain (USA\_WA1/2020). The primary immunobridging analyses compared the geometric mean titres (using a geometric mean ratio

[GMR]) and the seroresponse (defined as achieving at least 4-fold rise in SARS-CoV-2 NT50 from before Dose 1) rates in the evaluable immunogenicity population of participants without evidence of prior SARS-CoV-2 infection up to 1 month after Dose 3 in participants 2 to <5 years of age and up to 1 month after Dose 2 in participants 16 to 25 years of age. The prespecified immunobridging criteria were met for both the GMR and the seroresponse difference (Table 25 and Table 26, respectively).

**Table 25: SARS-CoV-2 GMTs (NT50) at 1 month after vaccination course – immunobridging subset - participants 2 to <5 years of age (C4591007) 1 month after Dose 3 and participants 16 to 25 years of age (C4591001) 1 month after Dose 2 – without evidence of SARS-CoV-2 infection – evaluable immunogenicity population**

	COMIRNATY (tozinameran)		GMR (95%CI) (2 to <5 years of age/16 to 25 years of age) <sup>c,d</sup>
	3 micrograms/dose 2 to <5 years of age (1 month after Dose 3) n <sup>a</sup> =143	30 micrograms/dose 16 to 25 years of age (1 month after Dose 2) n <sup>a</sup> =170	
Assay	GMT <sup>b</sup> (95% CI <sup>b</sup> )	GMT <sup>b</sup> (95% CI <sup>b</sup> )	
SARS-CoV-2 neutralisation assay - NT50 (titre) <sup>e</sup>	1535.2 (1388.2, 1697.8)	1180.0 (1066.6, 1305.4)	1.30 (1.13, 1.50)

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titre; LLOQ = lower limit of quantitation; NAAT = nucleic-acid amplification test; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Participants who had no serological or virological evidence [(up to 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007) blood sample collection)] of past SARS-CoV-2 infection [(i.e., N-binding antibody [serum] negative at Dose 1, Dose 3 (C4591007) and 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007), SARS-CoV-2 not detected by NAAT [nasal swab] at Dose 1, Dose 2, and Dose 3 (C4591007) study visits, and negative NAAT (nasal swab) at any unscheduled visit up to 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007) blood collection)] and had no medical history of COVID-19 were included in the analysis.

- n = Number of participants with valid and determinate assay results for the specified assay at the given dose/sampling time point.
- GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .
- GMRs and 2-sided 95% CIs were calculated by exponentiating the mean difference of the logarithms of the titres (2 to <5 years of age minus 16 to 25 years of age) and the corresponding CI (based on the Student t distribution).
- Immunobridging is declared if the lower bound of the 2-sided 95% CI for the GMR ratio is greater than 0.67 and the point estimate of the GMR is  $\geq 0.8$ .
- SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralisation Assay. The assay uses a fluorescent reporter virus derived from the USA\_WA1/2020 strain and virus neutralisation is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralised.

**Table 26: Difference in percentages of participants with seroresponse at 1 month after vaccination course – immunobridging subset –participants 2 to <5 years of age (C4591007) 1 month after Dose 3 and participants 16 to 25 years of age (C4591001) 1 month after Dose 2 without evidence of infection – evaluable immunogenicity population**

	COMIRNATY (tozinameran)		Difference in seroresponse rates % <sup>d</sup> (95% CI) <sup>e</sup> (2 to <5 years of age minus 16 to 25 years of age) <sup>f</sup>
	3 micrograms/dose 2 to <5 years of age (1 month after Dose 3) N <sup>a</sup> =141	30 micrograms/dose 16 to 25 Years of age (1 month after Dose 2) N <sup>a</sup> =170	
Assay	n <sup>b</sup> (%) (95% CI) <sup>c</sup>	n <sup>b</sup> (%) (95% CI) <sup>c</sup>	
SARS-CoV-2 neutralisation assay - NT50 (titre) <sup>g</sup>	141 (100.0) (97.4, 100.0)	168 (98.8) (95.8, 99.9)	1.2 (-1.5, 4.2)

Abbreviations: LLOQ = lower limit of quantitation; NAAT = nucleic acid amplification test; N-binding = SARS-CoV-2 nucleoprotein-binding; NT50 = 50% neutralising titre 50; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Seroresponse is defined as achieving a  $\geq 4$ -fold rise from baseline (before Dose 1). If the baseline measurement is below the LLOQ, a postvaccination assay result  $\geq 4 \times$  LLOQ is considered a seroresponse.

Note: Participants who had no serological or virological evidence (up to 1 month after Dose 2 [(C4591001) or 1 month after Dose 3 (C4591007) blood sample collection]) of past SARS-CoV-2 infection [(i.e., N-binding antibody [serum] negative at pre-Dose 1, pre-Dose 3 (C4591007) and 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007), SARS-CoV-2 not detected by NAAT [nasal swab] at pre-Dose 1, pre-Dose 2, and pre-Dose 3 (C4591007) study visits, and negative NAAT (nasal swab) at any unscheduled visit up to 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007) blood collection)] and had no medical history of COVID-19 were included in the analysis.

- N = number of participants with valid and determinate assay results both before vaccination and at 1 month after Dose 2. These values are the denominators for the percentage calculations.
- n = Number of participants with seroresponse for the given assay at the given dose/sampling time point.
- Exact 2-sided CI based on the Clopper and Pearson method.
- Difference in proportions, expressed as a percentage (2 to <5 years of age minus 16 to 25 years of age).
- 2-sided CI, based on the Miettinen and Nurminen method for the difference in proportions, expressed as a percentage.
- Immunobridging is declared if the lower bound of the 2-sided 95% CI for the difference in proportions is greater than -10.0% provided that the immunobridging criteria based on GMR were met.
- SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralisation Assay. The assay uses a fluorescent reporter virus derived from the USA\_WA1/2020 strain and virus neutralisation is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralised.

### Omicron and Delta variants

Using a non-validated fluorescence focus reduction neutralisation test assay against the Omicron variant of SARS-CoV-2 (BA.1), the NT50 GMT at 1 month after Dose 3 among a subset of 34 study participants without evidence of prior SARS-CoV-2 infection (82.5 [2-sided 95% CI: 55.4, 122.9]) was increased compared to the NT50 GMT before Dose 3 (14.0 [2-sided 95% CI: 10.6, 18.5]).

By comparison, in the same subset of 34 study participants without evidence of prior SARS-CoV-2 infection, there were notable higher NT50 GMTs at 1 month after Dose 3 against the Delta variant and wildtype SARS-CoV-2 (471.4 [2-sided 95% CI: 341.2, 651.1] and 471.4 [2-sided 95% CI: 344.6, 644.8], respectively). The NT50 GMTs before Dose 3 against the Delta variant and wildtype SARS-CoV-2 were 68 [2-sided 95% CI: 49.5, 93.3] and 70.1 [2-sided 95% CI: 51.1, 96], respectively.

An additional descriptive immunogenicity analysis was performed for participants 2 to <5 years of age who received a 3-dose course of COMIRNATY (tozinameran) in Phase 2/3 C4591007, compared with a subset of participants 18 to 50 years of age in Phase 3 Study C4591017 who had received a 2-dose primary course followed by a booster dose of COMIRNATY 30 micrograms. The comparator group (participants 18 to 50 years of age) in this analysis had a similar interval between COMIRNATY (tozinameran) Dose 2 and Dose 3 (median 13.0 weeks) as the participants 2 to <5 years of age (median 10.6 weeks). Among 34 participants 2 to <5 years of age without evidence of prior SARS-CoV-2 infection who received 3 doses of COMIRNATY 3 micrograms, neutralising GMTs were 114.3 at 1-month post-Dose 3. Among 27 participants 18 to 50 years of age without evidence of prior SARS-CoV-2 infection who received 3 doses of COMIRNATY 30 micrograms, Omicron neutralising GMTs were 164.2 at 1-month post Dose 3.

### ***Immunogenicity in infants 6 months to <2 years of age – after 3 doses***

Immunogenicity analyses have been performed in the immunobridging subset of 82 C4591007 participants 6 months to <2 years of age without evidence of infection up to 1 month after Dose 3 based on a data cutoff date of 29 April 2022.

SARS-CoV-2 50% neutralising antibody titres (NT50) 1 month after the vaccination course were compared between an immunogenicity subset of Phase 2/3 participants 6 months to <2 years of age from C4591007 and a randomly selected subset from C4591001 Phase 2/3 participants 16 to 25 years of age, using a microneutralisation assay against the reference strain (USA\_WA1/2020). The primary immunobridging analyses compared the geometric mean titres (using a GMR) and the seroresponse (defined as achieving at least 4-fold rise in SARS-CoV-2 NT50 from before Dose 1) rates in the evaluable immunogenicity population of participants without evidence of prior SARS-CoV-2 infection up to 1 month after Dose 3 in participants 6 months to <2 years of age and up to 1 month after Dose 2 in participants 16 to 25 years of age. The prespecified immunobridging criteria were met for both the GMR and the seroresponse difference (Table 27 and Table 28, respectively).

**Table 27: SARS-CoV-2 GMTs (NT50) at 1 month after vaccination course – immunobridging subset - participants 6 months to <2 years of age (C4591007) 1 month after Dose 3 and participants 16 to 25 years of age (C4591001) 1 month after Dose 2 – without evidence of SARS-CoV-2 – evaluable immunogenicity population**

	COMIRNATY (tozinameran)		GMR (95%CI) (6 months to <2 years of age/16 to 25 years of age) <sup>c,d</sup>
	3 micrograms/dose 6 months to <2 years of age (1 month after Dose 3) n <sup>a</sup> =82	30 micrograms/dose 16 to 25 years of age (1 month after Dose 2) n <sup>a</sup> =170	
Assay	GMT <sup>b</sup> (95% CI <sup>b</sup> )	GMT <sup>b</sup> (95% CI <sup>b</sup> )	
SARS-CoV-2 neutralisation assay - NT50 (titre) <sup>c</sup>	1406.5 (1211.3, 1633.1)	1180.0 (1066.6, 1305.4)	1.19 (1.00, 1.42)

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titre; LLOQ = lower limit of quantitation; NAAT = nucleic-acid amplification test; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Participants who had no serological or virological evidence [(up to 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007) blood sample collection)] of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Dose 1, Dose 3 (C4591007) and 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007), SARS-CoV-2 not detected by NAAT [nasal swab] at Dose 1, Dose 2, and Dose 3 (C4591007) study visits, and negative NAAT (nasal swab) at any unscheduled visit up to 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007) blood collection)] and had no medical history of COVID-19 were included in the analysis.



- n = Number of participants with valid and determinate assay results for the specified assay at the given dose/sampling time point.
- GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titre titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .
- GMRs and 2-sided 95% CIs were calculated by exponentiating the mean difference of the logarithms of the titres (6 months to <2 years of age minus 16 to 25 years of age) and the corresponding CI (based on the Student t distribution).
- Immunobridging is declared if the lower bound of the 2-sided 95% CI for the GMR ratio is greater than 0.67 and the point estimate of the GMR is  $\geq 0.8$ .
- SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralisation Assay. The assay uses a fluorescent reporter virus derived from the USA\_WA1/2020 strain and virus neutralisation is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralised.

**Table 28: Difference in percentages of participants with seroresponse at 1 month after vaccination course – immunobridging subset – participants 6 months to <2 years of age (C4591007) 1 month after Dose 3 and participants 16 to 25 years of age (C4591001) to 1 month after Dose 2 without evidence of infection – evaluable immunogenicity population**

	COMIRNATY (tozinameran)		Difference in seroresponse rates % <sup>d</sup> (95% CI) <sup>e</sup> (6 months to <2 years of age minus 16 to 25 years of age) <sup>f</sup>
	3 micrograms/dose 6 to 23 months of age (1 month after Dose 3) N <sup>a</sup> =80	30 micrograms/dose 16 to 25 years of age (1 month after Dose 2) N <sup>a</sup> =170	
Assay	n <sup>b</sup> (%) (95% CI) <sup>c</sup>	n <sup>b</sup> (%) (95% CI) <sup>c</sup>	
SARS-CoV-2 neutralisation assay - NT50 (titre) <sup>g</sup>	80 (100.0) (95.5, 100.0)	168 (98.8) (95.8, 99.9)	1.2 (-3.4, 4.2,)

Abbreviations: LLOQ = lower limit of quantitation; NAAT = nucleic acid amplification test; N-binding = SARS-CoV-2 nucleoprotein-binding; NT50 = 50% neutralising titre 50; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Seroresponse is defined as achieving a  $\geq 4$ -fold rise from baseline (before Dose 1). If the baseline measurement is below the LLOQ, a postvaccination assay result  $\geq 4 \times \text{LLOQ}$  is considered a seroresponse.

Note: Participants who had no serological or virological evidence [(up to 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007) blood sample collection) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at pre-Dose 1, Dose 3 (C4591007) and 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007), SARS-CoV-2 not detected by NAAT [nasal swab] at pre-Dose 1, pre-Dose 2, and pre-Dose 3 (C4591007) study visits, and negative NAAT (nasal swab) at any unscheduled visit up to 1 month after Dose 2 (C4591001) or 1 month after Dose 3 (C4591007) blood collection)] and had no medical history of COVID-19 were included in the analysis.

- N = number of participants with valid and determinate assay results both before vaccination and at 1 month after Dose 2. These values are the denominators for the percentage calculations.
- n = Number of participants with seroresponse for the given assay at the given dose/sampling time point.
- Exact 2-sided CI based on the Clopper and Pearson method.
- Difference in proportions, expressed as a percentage (6 months to <2 years of age minus 16 to 25 years of age).
- 2-sided CI, based on the Miettinen and Nurminen method for the difference in proportions, expressed as a percentage.
- Immunobridging is declared if the lower bound of the 2-sided 95% CI for the difference in proportions is greater than -10.0% provided that the immunobridging criteria based on GMR were met.
- SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralisation Assay. The assay uses a fluorescent reporter virus derived from the USA\_WA1/2020 strain and virus neutralisation is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralised.

## Omicron and Delta variants

Using a non-validated fluorescence focus reduction neutralisation test assay against the Omicron variant of SARS-CoV-2 (BA.1), the NT50 GMT at 1 month after Dose 3 among a subset of 32 study participants without evidence of prior SARS-CoV-2 infection (127.5 [2-sided 95% CI: 90.2, 180.1]) was increased compared to the NT50 GMT before Dose 3 (16.3 [2-sided 95% CI: 12.8, 20.8]).

By comparison, in the same subset of 32 study participants without evidence of prior SARS-CoV-2 infection, there were notable higher NT50 GMTs at 1 month after Dose 3 against the Delta variant and wildtype SARS-CoV-2 (606.3 [2-sided 95% CI: 455.5, 806.9] and 640.0 [2-sided 95% CI: 502.6, 815.0], respectively). The NT50 GMTs before Dose 3 against the Delta variant and wildtype SARS-CoV-2 were 94.1 [2-sided 95% CI: 67.9, 130.5] and 103.7 [2-sided 95% CI: 78.4, 137.3], respectively.

An additional descriptive immunogenicity analysis was performed for participants 6 months to <2 years of age who received a 3-dose course of COMIRNATY (tozinameran) in Phase 2/3 C4591007, compared with a subset of participants 18 to 50 years of age in Phase 3 Study C4591017 who had received a 2-dose primary course followed by a booster dose of COMIRNATY 30 micrograms. The comparator group (participants 18 to 50 years of age) in this analysis had a similar interval between COMIRNATY (tozinameran) Dose 2 and Dose 3 (median 13.0 weeks) as the participants 6 months to <2 years of age (median 12.9 weeks). Among 32 participants 6 months to <2 years of age without evidence of prior SARS-CoV-2 infection who received 3 doses of COMIRNATY 3 micrograms, Omicron neutralising GMTs were 128.8 at 1-month post-Dose 3. Among 27 participants 18 to 50 years of age without evidence of prior SARS-CoV-2 infection who received 3 doses of COMIRNATY (tozinameran) 30 micrograms, Omicron neutralising GMTs were 164.2 at 1-month post Dose 3.

### ***Immunogenicity in participants 18 years of age and older – after booster dose***

Effectiveness of a booster dose of COMIRNATY (tozinameran) was based on an assessment of 50% neutralising titres (NT50) against SARS-CoV-2 (USA\_WA1/2020). In Study C4591001, analyses of NT50 1 month after the booster dose compared to 1 month after the primary series in individuals 18 to 55 years of age who had no serological or virological evidence of past SARS-CoV-2 infection up to 1 month after the booster vaccination demonstrated noninferiority for both GMR and difference in seroresponse rates. Seroresponse for a participant was defined as achieving a  $\geq 4$ -fold rise in NT50 from baseline (before Dose 1). These analyses are summarised in Table 29.

**Table 29. SARS-CoV-2 Neutralisation Assay - NT50 (titre)<sup>†</sup> (SARS-CoV-2 USA\_WA1/2020) – GMT and Seroresponse rate comparison of 1 month after booster dose to 1 month after primary series – Participants 18 to 55 years of age without evidence of infection up to 1 month after booster dose\* – Booster dose Evaluable Immunogenicity Population<sup>‡</sup>**

	n	1 month after booster dose (95% CI)	1 month after primary series (95% CI)	1 month after booster dose/- 1 month after primary series (97.5% CI)	Met noninferiority objective (Y/N)
Geometric mean 50% neutralising titre (GMT) <sup>b</sup>	212 <sup>a</sup>	2466.0 <sup>b</sup> (2202.6, 2760.8)	750.6 <sup>b</sup> (656.2, 858.6)	3.29 <sup>c</sup> (2.77, 3.90)	Y <sup>d</sup>

<b>Seroresponse rate (%) for 50% neutralising titre<sup>†</sup></b>	200 <sup>c</sup>	199 <sup>f</sup> 99.5% (97.2%, 100.0%)	196 <sup>f</sup> 98.0% (95.0%, 99.5%)	1.5% <sup>g</sup> (-0.7%, 3.7% <sup>h</sup> )	Y <sup>i</sup>
---	------------------	--	---	--	----------------

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titre; LLOQ = lower limit of quantitation; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; Y/N = yes/no.

<sup>†</sup> SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralisation Assay. The assay uses a fluorescent reporter virus derived from the USA\_WA1/2020 strain and virus neutralisation is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralised.

\* Participants who had no serological or virological evidence (up to 1 month after receipt of a booster dose of COMIRNATY (tozinameran)) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative and SARS-CoV-2 not detected by NAAT [nasal swab]) and had a negative NAAT (nasal swab) at any unscheduled visit up to 1 month after the booster dose were included in the analysis.

± All eligible participants who had received 2 doses of COMIRNATY (tozinameran) as initially randomised, with Dose 2 received within the predefined window (within 19 to 42 days after Dose 1), received a booster dose of COMIRNATY (tozinameran), had at least 1 valid and determinate immunogenicity result after booster dose from a blood collection within an appropriate window (within 28 to 42 days after the booster dose), and had no other important protocol deviations as determined by the clinician.

- n = Number of participants with valid and determinate assay results at both sampling time points within specified window.
- GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to 0.5 × LLOQ.
- GMRs and 2-sided 97.5% CIs were calculated by exponentiating the mean differences in the logarithms of the assay and the corresponding CIs (based on the Student t distribution).
- Noninferiority is declared if the lower bound of the 2-sided 97.5% CI for the GMR is > 0.67 and the point estimate of the GMR is ≥ 0.80.
- n = Number of participants with valid and determinate assay results for the specified assay at baseline, 1 month after Dose 2 and 1 month after the booster dose within specified window. These values are the denominators for the percentage calculations.
- Number of participants with seroresponse for the given assay at the given dose/sampling time point. Exact 2-sided CI based on the Clopper and Pearson method.
- Difference in proportions, expressed as a percentage (1 month after booster dose – 1 month after Dose 2).
- Adjusted Wald 2-sided CI for the difference in proportions, expressed as a percentage.
- Noninferiority is declared if the lower bound of the 2-sided 97.5% CI for the percentage difference is > -10%.

### ***Relative vaccine efficacy in participants 16 years of age and older – after booster dose***

An interim efficacy analysis of Study C4591031, a placebo-controlled booster study, was performed in approximately 10,000 participants 16 years of age and older who were recruited from Study C4591001, evaluated confirmed COVID-19 cases accrued from at least 7 days after booster vaccination up to a data cut-off date of 8 February 2022 (a period when Delta and then Omicron was the predominant variant), which represents a median of 2.8 months (range 0.3 to 7.5 months) post-booster follow-up. Vaccine efficacy of the COMIRNATY (tozinameran) booster dose after the primary series relative to the placebo booster group who only received the primary series dose was assessed. The relative vaccine efficacy information for participants 16 years of age and older is presented in Table 30.

**Table 30: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Booster Vaccination – Participants 16 Years of Age and Older Without Evidence of Infection and Participants With or Without Evidence of Infection Prior to 7 Days After Booster Vaccination – Evaluable Efficacy Population**

<b>First COVID-19 occurrence from 7 days after booster dose in participants without evidence of prior SARS-CoV-2 infection*</b>			
	<b>COMIRNATY (tozinameran) N<sup>a</sup>=4689 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Placebo N<sup>a</sup>=4664 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Relative Vaccine Efficacy<sup>e</sup> % (95% CI<sup>f</sup>)</b>
First COVID-19 occurrence from 7 days after booster vaccination	63 1.098 (4639)	148 0.932 (4601)	63.9 (51.1, 73.5)

<b>First COVID-19 occurrence from 7 days after booster dose in participants with or without evidence of prior SARS-CoV-2 infection</b>			
	<b>COMIRNATY (tozinameran) N<sup>a</sup>=4977 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Placebo N<sup>a</sup>=4942 Cases n<sup>1b</sup> Surveillance Time<sup>c</sup> (n<sup>2d</sup>)</b>	<b>Relative Vaccine Efficacy<sup>e</sup> % (95% CI<sup>f</sup>)</b>
First COVID-19 occurrence from 7 days after booster vaccination	67 1.173 (4903)	150 0.989 (4846)	62.4 (49.5, 72.2)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhea; vomiting).

\* Participants who had no serological or virological evidence (prior to 7 days after receipt of the booster vaccination) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visit 1, and had a negative NAAT [nasal swab] at any unscheduled visit prior to 7 days after booster vaccination) were included in the analysis.

- a. N = Number of participants in the specified group.
- b. n<sup>1</sup> = Number of participants meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after the booster vaccination to the end of the surveillance period.
- d. n<sup>2</sup> = Number of participants at risk for the endpoint.
- e. Relative vaccine efficacy of the COMIRNATY booster group relative to the placebo group (non-booster).
- f. Two-sided confidence interval (CI) for relative vaccine efficacy is derived based on the Clopper and Pearson method adjusted for surveillance time.

### ***Immunogenicity in children 5 to <12 years of age – after booster dose***

In a subset from C4591007, a total of 123 children 5 to <12 years of age received a booster dose of COMIRNATY (tozinameran) 10 micrograms after completing the primary series. All participants in the 3-Dose immunogenicity subset, received the booster dose 7 - < 9 months after Dose 2, (n = 37 [30.1%] at 7 - < 8 months and n = 86 [69.9%] at 8 - < 9 months).

Effectiveness of a booster dose of COMIRNATY (tozinameran) was based on an assessment of NT50 against the reference strain of SARS-CoV-2 (USA\_WA1/2020). Analyses of NT50 1 month after the booster dose compared to before the booster dose demonstrated an increase in GMTs in individuals 5 to <12 years of age who had no serological or virological evidence of past SARS-CoV-2 infection up to 1 month after the booster dose. This analysis is summarised in Table 31.

**Table 31: Summary of Geometric Mean Titres – NT50 – Participants Without Evidence of Infection – Phase 2/3 – Immunogenicity Set – 5 to <12 Years of Age – Evaluable Immunogenicity Population**

		COMIRNATY (tozinameran) 10 micrograms/Dose					
		3-Dose Set		2-Dose Set		Total	
Assay	Dose/ Sampling Time Point <sup>a</sup>	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )
SARS-CoV-2 neutralisation assay - NT50 (titre)	Dose 1 Prevax	79	20.5 (20.5, 20.5)	67	20.5 (20.5, 20.5)	146	20.5 (20.5, 20.5)
	1 month after Dose 2	29	1659.4 (1385.1, 1988.0)	67	1110.7 (965.3, 1278.1)	96	1253.9 (1116.0, 1408.9)
	Dose 3 Prevax	67	271.0 (229.1, 320.6)	-	-	67	271.0 (229.1, 320.6)

	1 month after Dose 3	67	2720.9 (2280.1, 3247.0)	-	-	67	2720.9 (2280.1, 3247.0)
--	----------------------	----	----------------------------	---	---	----	----------------------------

Abbreviations: CI = confidence interval; GMT = geometric mean titre; LLOQ = lower limit of quantitation; NAAT = nucleic acid amplification test; N-binding = SARS-CoV-2 nucleoprotein-binding; NT50 = 50% neutralising titre; Prevax = before vaccination; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Three-dose immunogenicity set included the first 130 participants who received Dose 3 and completed 1-month post-Dose 3 visit prior to March 15, 2022. Among those, 30 had blood sample collection at 1-month post-Dose 2. Two-dose immunogenicity set included an extra 67 participants randomly selected from previous Dose-2 evaluable immunogenicity population and without evidence of infection up to 1-month post-Dose 2 subset used for 2-dose immunobridging analysis.

Note: Participants included in this analysis had no serological or virological evidence of past SARS-CoV-2 infection up to the 1-month post-Dose 2 (for 1-month post-Dose 2 time point) or 1-month post-Dose 3 (for pre-Dose 3 and 1-month post-Dose 3 time point) study blood sample collection. Having no evidence of past SARS-CoV-2 infection up to 1-month post-Dose 2 was defined as having a negative N-binding antibody (serum) result at the Dose 1 and 1-month post-Dose 2 study visits; a negative NAAT (nasal swab) result at the Dose 1 and Dose 2 study visits and any unscheduled visit prior to the 1-month post-Dose 2 blood sample collection; and no medical history of COVID-19. Having no evidence of past SARS-CoV-2 infection up to 1-month post-Dose 3 was defined as having a negative N-binding antibody (serum) result at the Dose 1, 1-month post-Dose 2 (if available), Dose 3, and 1-month post-Dose 3 study visits; a negative NAAT (nasal swab) result at the Dose 1, Dose 2, and Dose 3 study visits and any unscheduled visit prior to the 1-month post-Dose 3 blood sample collection; and no medical history of COVID-19.

- Protocol-specified timing for blood sample collection.
- n = Number of participants with valid and determinate assay results for the specified assay at the given dose/sampling time point.
- GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .

### ***Immunogenicity in pregnant women and infants born to maternal participants – after 2 doses with COMIRNATY (tozinameran)***

Study C4591015 was a Phase 2/3 multinational, placebo-controlled, observer-blind study that enrolled pregnant women 18 years of age and older to receive 2 doses of COMIRNATY (tozinameran) (n=173) or placebo (n=173). Pregnant women received Dose 1 of COMIRNATY (tozinameran) at 24 to 34 weeks gestation and the majority (90.2%) received the second dose 19 to 23 days after Dose 1.

Descriptive immunogenicity analysis was performed in pregnant participants receiving COMIRNATY (tozinameran) in Study C4591015 compared to a historical subset of age-matched nonpregnant participants randomly selected from Study C4591001 evaluating the ratio of the neutralising GMT (GMR) 1 month after Dose 2.

The evaluable immunogenicity population who received COMIRNATY (tozinameran) in the maternal participants group in Study C4591015 (n=111) and in nonpregnant participants in Study C4591001 (n=114) comprised of 69.4% vs. 82.5% White, 27.0% vs. 5.3% Black or African American, 1.8% vs. 6.1% Asian, 0 vs 4.4% multiracial participants, 37.8% vs 34.2% Hispanic/Latino, 37.8% vs 3.5% had a positive baseline SARS-CoV-2 status, and 38.7% vs 27.2% were obese [BMI  $\geq 30$  kg/m<sup>2</sup> (pre-pregnancy weight in participants in Study C4591015)], respectively. The median age was 30 years (range 18 to 44 years of age) in both groups.

The immunogenicity results after 2 doses of COMIRNATY (tozinameran) in pregnant participants 18 years of age and older without evidence of infection up to 1 month after Dose 2 are presented in Table 32.

Post hoc immunogenicity results after 2 doses of COMIRNATY (tozinameran) in pregnant participants 18 years of age and older with or without evidence of infection are presented in Table 33. The model-adjusted GMT and GMR were calculated based on a regression model adjusting for age and baseline neutralising titres.

**Table 32. Geometric Mean Ratios – Participants Without Evidence of Infection up to 1 Month After Dose 2 – Maternal Participants (Study C4591015) and Nonpregnant Female Participants (Study C4591001) – Evaluable Immunogenicity Population**

		Vaccine Group (as Randomised) Tozinameran (30 micrograms)				Pregnant/ Nonpregnant
		C4591015 (Maternal)		C4591001 (Nonpregnant Women)		
Assay	Dose/ Sampling Time Point <sup>a</sup>	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	GMR <sup>d</sup> (95% CI <sup>d</sup> )
SARS-CoV-2 neutralisation assay - NT50 (titre)	2/1 Month	58	1109.2 (849.2, 1448.9)	107	1663.7 (1411.5, 1960.8)	0.67 (0.50, 0.90)

Abbreviations: GMR = geometric mean ratio; GMT = geometric mean titre; LLOQ = lower limit of quantitation; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Blood samples for immunogenicity assessment drawn at the delivery visit but within 1 month after Dose 2 visit window were also included in this analysis.

Note: Human immunodeficiency virus (HIV)-positive participants are not included in this summary.

Note: Participants from C4591001 are a selected subset of age matched nonpregnant female Phase 3 participants.

Note: Participants who had no serological or virological evidence (prior to the 1 month after Dose 2 blood sample collection) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Dose 1 and 1 month after Dose 2 and no positive result between visits, negative NAAT [nasal swab] at Dose 1, Dose 2, and any unscheduled visit prior to the 1 month after Dose 2 blood sample collection) were included in the analysis.

a. Protocol-specified timing for blood sample collection.

b. n = Number of participants with valid and determinate assay results for the specified assay at the given dose/sampling time point.

c. GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .

d. GMRs and 2-sided 95% CIs were calculated by exponentiating the mean difference of the logarithms of the assay and the corresponding CIs (based on the Student t distribution).

**Table 33. Model Adjusted Geometric Mean Ratios – Participants With or Without Evidence of Infection – Maternal Participants (Study C4591015) and Nonpregnant Female Participants (Study C4591001) – Evaluable Immunogenicity Population**

		Vaccine Group (as Randomised) Tozinameran (30 micrograms)				Pregnant/ Nonpregnant
		C4591015 (Maternal)		C4591001 (Nonpregnant Women)		
Assay	Dose/ Sampling Time Point <sup>a</sup>	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	n <sup>b</sup>	GMT <sup>c</sup> (95% CI <sup>c</sup> )	GMR <sup>d</sup> (95% CI) <sup>d</sup>
SARS-CoV-2 neutralisation assay - NT50 (titre) <sup>a</sup>	2/1 month	99	1900.0 (1518.2, 2377.7)	113	2005.7 (1627.0, 2472.6)	0.95 (0.69, 1.30)

Abbreviations: GMR = geometric mean ratio; GMT = geometric mean titre; LLOQ = lower limit of quantitation; NT50 = 50% neutralising titre; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Blood samples for immunogenicity assessment drawn at the delivery visit but within 1 month after Dose 2 visit window were also included in this analysis.

Note: Human immunodeficiency virus (HIV)-positive participants are not included in this summary.

Note: Participants from C4591001 are a selected subset of age matched nonpregnant female Phase 3 participants.

a. Protocol-specified timing for blood sample collection.

b. n = Number of participants with valid and determinate assay results for the specified assay at both baseline and the given dose/sampling time point.

c. GMTs and 2-sided CIs were calculated by exponentiating the LS means and the corresponding CIs based on analysis of log-transformed NT50 titres using a regression model with group, age at Dose 1 in years (continuous), and baseline log transformed NT50 titres.

d. GMRs (ratio of GMTs of pregnant women to nonpregnant women) and 2-sided CIs were calculated by exponentiating the difference of LS means and the corresponding CIs based on the same regression model as above.

In an additional descriptive immunogenicity analysis, infants born to maternal participants who received COMIRNATY (tozinameran) had higher geometric mean concentrations (GMCs) of full length S-binding immunoglobulin G (IgG) concentrations at birth and at 6 months after delivery [5576.4 (95% CI: 4246.2, 7323.2); n=91 and 311.1 (95% CI: 235.8, 410.5); n=83], respectively, compared to infants born to maternal participants from the placebo group [19.4 (95% CI: 10.2, 37.0); n=92 and 22.0 (95% CI: 11.4, 42.7); n=69].

### ***Immunogenicity in immunocompromised participants (adults and children)***

Study C4591024 is a Phase 2b, open-label study (n=124) that enrolled 117 immunocompromised participants 2 to <18 years of age receiving immunomodulator therapy (n = 35) or who have undergone solid organ transplant (within the previous 3 months) and are on immunosuppression (n = 40) or who have undergone bone marrow or stem cell transplant at least 6 months prior to enrolment (n = 42). Study C4591024 also enrolled 7 immunocompromised participants 18 years of age and older treated for NSCLC (n = 1) or receiving haemodialysis for secondary to end-stage renal disease (n = 1), or receiving immunomodulator therapy for an autoimmune inflammatory disorder (n = 5). Participants received 4 age-appropriate doses of COMIRNATY (tozinameran) (3 micrograms, 10 micrograms, or 30 micrograms); the first 2 doses separated by 21 days, with the third dose occurring 28 days after the second dose, followed by a fourth dose, 3 to 6 months after Dose 3.

Exploratory immunogenicity results pre-vaccination and after 3 and 4 doses of COMIRNATY (tozinameran) in immunocompromised participants 2 years of age and older are presented in Table 34.

**Table 34. Summary of Geometric Mean Titres – Participants With or Without Evidence of Infection by Age Group – Dose 3 or Dose 4 Evaluable Immunogenicity Population**

		COMIRNATY (tozinameran)							
		3 micrograms Age Group: 2 to <5 Years		10 micrograms Age Group: 5 to <12 Years		30 micrograms Age Group: 12 to <18 Years		30 micrograms Age Group: ≥18 Years	
Assay	Dose/ Sampling Time Point <sup>b</sup>	n <sup>c</sup>	GMT <sup>d</sup> (95% CI <sup>d</sup> )	n <sup>c</sup>	GMT <sup>d</sup> (95% CI <sup>d</sup> )	n <sup>c</sup>	GMT <sup>d</sup> (95% CI <sup>d</sup> )	n <sup>c</sup>	GMT <sup>d</sup> (95% CI <sup>d</sup> )
SARS-CoV-2 neutralisation assay – reference strain – NT50 (titre) <sup>a</sup>	1/Prevax <sup>c</sup>	27	43.5 (43.5, 43.5)	59	44.5 (42.5, 46.7)	13	55.1 (32.9, 92.1)	5	93.4 (11.2, 779.0)
	3/1 Month <sup>f</sup>	26	899.7 (484.2, 1671.6)	56	1701.2 (1094.8, 2643.3)	11	2837.2 (888.9, 9056.4)	4	205.4 (21.8, 1936.1)
	4/Pre-Dose 4 <sup>c</sup>	24	492.0 (246.9, 980.2)	55	924.3 (580.1, 1472.7)	10	3745.6 (1805.3, 7771.3)	3	606.2 (5.3, 68756.0)
	4/1 Month <sup>g</sup>	16	3893.7 (1738.4, 8721.4)	31	6236.1 (3711.3, 10478.6)	6	10255.4 (3643.7, 28864.2)	4	1031.3 (56.9, 18681.7)
	4/6 Months <sup>g</sup>	15	1503.9 (685.9, 3297.4)	28	2224.6 (1144.1, 4325.5)	5	5020.6 (1773.6, 14212.3)	3	1605.6 (28.5, 90614.9)

Abbreviations: CI = confidence interval; GMT = geometric mean titre; LLOQ = lower limit of quantitation; NT50 = 50% neutralising titre; Prevax = before vaccination; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

a. SARS-CoV-2 NT50 were determined using a validated 384-well assay platform (original strain [USA-WA1/2020, isolated in January 2020]).

b. Protocol-specified timing for blood sample collection.

- c. n = Number of participants with valid and determinate assay results for the specified assay at the given dose/sampling time point.
- d. GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .
- e. Dose 3 or Dose 4 evaluable immunogenicity population.
- f. Dose 3 evaluable immunogenicity population.
- g. Dose 4 evaluable immunogenicity population.

### ***Concomitant vaccine administration with influenza vaccine***

In Study C4591030, a Phase 3 multicentre, randomised, observer-blind study, 1,134 participants 18 to 64 years of age who had received 3 doses of COMIRNATY (tozinameran) at least 3 months prior were randomised in a 1:1 ratio to receive either COMIRNATY (tozinameran) coadministered with a SIIV, quadrivalent (Afluria Quad) followed 1 month later by placebo (Group 1, n=568) or an inactivated influenza vaccine with placebo followed 1 month later with COMIRNATY (tozinameran) (Group 2, n=566).

The immune responses to COMIRNATY (tozinameran) and SIIV were similar after COMIRNATY (tozinameran) administered concomitantly with SIIV compared with those elicited by either vaccine administered alone. The non-inferiority criterion was achieved for both full-length S-binding immunoglobulin G (IgG) and all 4 influenza strain-specific hemagglutination inhibition (HAI) titres.

The immunogenicity results are presented in Table 35 and Table 36.

**Table 35. Geometric Mean Ratio for Full-Length S-Binding IgG Levels (U/mL) at 1 Month After COMIRNATY (tozinameran) Vaccination – Evaluable Immunogenicity Population**

Assay	Vaccine Group (as Randomised)				Coadministration Group/Separate Administration Group
	Coadministration Group		Separate-Administration Group		
	n <sup>a</sup>	GMC <sup>b</sup> (95% CI <sup>b</sup> )	n <sup>a</sup>	GMC <sup>b</sup> (95% CI <sup>b</sup> )	GMR <sup>c</sup> (95% CI <sup>c</sup> )
Full-length S-binding IgG (U/mL)	499	13806.5 (12838.9, 14847.0)	413	16254.6 (15035.5, 17572.5)	0.83 (0.77, 0.89)

Abbreviations: CI = confidence interval; GMC = geometric mean concentration; GMR = geometric mean ratio; IgG = immunoglobulin G; LLOQ = lower limit of quantitation; LS Means = least squares means; S = spike protein.

Note: The baseline was defined as Visit 1 for the coadministration group and Visit 2 for the separate-administration group.

- a. n = Number of participants with valid and determinate assay results for the specified assay at the given sampling time point.
- b. GMC and the 2-sided 95% CI were calculated by exponentiating the concentrations and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ .
- c. GMR and the corresponding 2-sided 95% CI were calculated by exponentiating the difference in LS Means and the corresponding CIs based on analysis of logarithmically transformed assay results using a linear regression model with terms of vaccine group, age group, and the corresponding baseline assay results (log scale). Noninferiority is declared if the lower bound of the 2-sided 95% CI for the GMR is greater than 0.67.



**Table 36. Geometric Mean Ratio for Strain-Specific HAI Titres at 1 Month After SIIV Vaccination – Evaluable SIIV Immunogenicity Population**

Strain	Vaccine Group (as Randomised)				Coadministration Group/Separate Administration Group
	Coadministration Group		Separate Administration Group		
	n <sup>a</sup>	GMT <sup>b</sup> (95% CI <sup>b</sup> )	n <sup>a</sup>	GMT <sup>b</sup> (95% CI <sup>b</sup> )	GMR <sup>c</sup> (95% CI <sup>c</sup> )
B/Austria	514	72.4 (64.2, 81.7)	491	78.3 (69.3, 88.5)	0.89 (0.77, 1.04)
B/Phuket	520	87.4 (79.7, 95.7)	496	86.3 (78.4, 94.9)	1.00 (0.89, 1.13)
H1N1 A/Victoria	516	344.3 (312.4, 379.3)	492	362.3 (326.3, 401.6)	0.95 (0.83, 1.09)
H3N2 A/Darwin	519	230.6 (209.5, 253.8)	491	242.2 (221.2, 265.2)	0.96 (0.85, 1.09)

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titre; HAI = hemagglutination inhibition; LLOQ = lower limit of quantitation; LS Means = least squares means; SIIV = seasonal inactivated influenza vaccine; ULOQ = upper limit of quantitation.

Note: The baseline for the SIIV assay was defined at Visit 1.

- n = Number of participants with valid and determinate assay results for the specified assay at the given sampling time point.
- GMTs and the 2-sided 95% CIs were calculated by exponentiating the titres and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to  $0.5 \times \text{LLOQ}$ , and results above the ULOQ were set to  $\text{ULOQ} + 1$ .
- GMRs and the corresponding 2-sided 95% CI were calculated by exponentiating the difference in LS Means and the corresponding CIs based on analysis of logarithmically transformed assay results using a linear regression model with terms of vaccine group, age group, and the corresponding baseline assay results (log scale). Noninferiority is declared if the lower bound of the 2-sided 95% CI for the GMR is greater than 0.67.

## 5.2 Pharmacokinetic properties

Not applicable.

## 5.3 Preclinical safety data

### Genotoxicity/Carcinogenicity

Neither genotoxicity nor carcinogenicity studies were performed. The components of the vaccine (lipids and mRNA) are not expected to have genotoxic potential.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

((4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl)bis(2-hexyldecanoate) (ALC-0315)

2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide (ALC-0159)

Distearoylphosphatidylcholine (DSPC)

Cholesterol

Trometamol

Trometamol hydrochloride

Sucrose

Water for injections

## 6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned in Section 4.2 Dose and method of administration.

## 6.3 Shelf life

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

### *Unopened vial*

COMIRNATY Original/Omicron BA.4-5 may be received frozen at -90°C to -60°C or at -25°C to -15°C. Frozen vaccine can be stored either at -90°C to -60°C or 2°C to 8°C upon receipt.

Once removed from frozen storage, the unopened vial may be stored refrigerated at 2°C to 8°C for a single period of up to 10 weeks within the approved shelf life.

Upon moving the product to 2°C to 8°C storage, the updated expiry date must be written on the outer carton and the vaccine should be used or discarded by the updated expiry date. The original expiry date should be crossed out.

If the vaccine is received at 2°C to 8°C it should be stored at 2°C to 8°C. Check that the expiry date on the outer carton has been updated to reflect the refrigerated expiry date and that the original expiry date has been crossed out.

When stored frozen at -90°C to -60°C, the vaccine can be thawed at either 2°C to 8°C or at temperatures up to 30°C.

Thawed vials can be handled in room light conditions.

Once thawed, the vaccine should not be re-frozen.

### ***COMIRNATY Original/Omicron BA.4-5 – Suspension for Injection (Grey or Blue Cap)***

#### *Opened vial*

Chemical and physical in-use stability has been demonstrated for 12 hours at 2°C to 30°C. From a microbiological point of view, unless the method of opening precludes the risks of microbial contamination, the product should be used immediately after the first puncture. If not used immediately, in-use storage times and conditions cannot be longer than 12 hours at 2°C to 30°C.

### ***COMIRNATY Original/Omicron BA.4-5 - Concentrated Suspension for Injection (Orange or Maroon Cap)***

#### *Diluted medicinal product*

Chemical and physical in-use stability has been demonstrated for 12 hours at 2°C to 30°C, after dilution with sodium chloride 9 mg/mL (0.9%) solution for injection. From a microbiological point of view, unless the method of opening precludes the risks of microbial contamination, the product should be used immediately. If not used immediately, in-use storage times and conditions cannot be longer than 12 hours at 2°C to 30°C.

## 6.4 Special precautions for storage

Store in the original package in order to protect from light.

During storage, minimise exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.

When stored frozen at -90°C to -60°C, the vaccine can be thawed at either 2°C to 8°C or at room temperature (up to 30°C). For detailed instructions see Section 4.2 Dose and method of administration, Handling instructions (Handling prior to use).

Once thawed, the vaccine cannot be re-frozen.

Thawed vials can be handled in room light conditions.

For storage conditions after thawing and dilution of the medicinal product, see Section 6.3 Shelf life.

For additional advice on storing COMIRNATY Original/Omicron BA.4-5, contact Pfizer Australia on 1800 675 229.

## 6.5 Nature and contents of container

COMIRNATY Original/Omicron BA.4-5 – Suspension for injection (Grey or Blue cap): 2 mL clear vial (Type I glass) with a stopper (synthetic bromobutyl rubber) and a Grey or Blue flip-off plastic cap with aluminium seal. Each vial contains either 1 or 6 doses, see Section 4.2 Dose and method of administration.

- Light Grey or Light Blue cap: single dose vial
- Dark Grey or Dark Blue cap: 6 dose multidose vial

COMIRNATY Original/Omicron BA.4-5 – Concentrated Suspension for injection (Orange or Maroon cap): 2 mL clear vial (Type I glass) with a stopper (synthetic bromobutyl rubber) and an Orange or Maroon flip-off plastic cap with aluminium seal. Each vial contains 10 doses after dilution, see Section 4.2 Dose and method of administration.

- Orange or Maroon cap: 10 dose multidose vial after dilution

Pack size: 10 vials, 195 vials

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal

In Australia, any unused medicine or waste material should be disposed of in accordance with local requirements.

## 6.7 Physicochemical properties

### CAS number

2417899-77-3

## 7. MEDICINE SCHEDULE (POISONS STANDARD)

S4 – Prescription Only Medicine.

## 8. SPONSOR

Pfizer Australia Pty Ltd  
Level 17, 151 Clarence Street  
Sydney NSW 2000  
Toll Free Number: 1800 675 229  
[www.pfizermedinfo.com.au](http://www.pfizermedinfo.com.au)

## 9. DATE OF FIRST APPROVAL

Aust R 400874: 23 January 2023

Aust R 413718, 412350, 413720 & 413719: 21 December 2023

Aust R 417266: 14 May 2024

## 10. DATE OF REVISION

02 December 2025

COMIRNATY<sup>®</sup> is a registered trademark of BioNTech SE. Used under license.

### Summary Table of Changes

Section changed	Summary of new information
4.4	Addition of Study C4591024 data (immunocompromised)
4.6	Addition of Study C4591015 data (maternal study)
4.8	Addition of safety data for Study C4591024, C4591015 and C4591048
4.9	Inclusion of post-authorisation experience
5.1	Addition of study C4591024, C4591015 and C4591048 clinical data