

AUSTRALIAN PRODUCT INFORMATION

FASENRA® (benralizumab) solution for injection prefilled syringe and prefilled pen (FASENRA PEN™)

1 NAME OF THE MEDICINE

Benralizumab

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each FASENRA prefilled syringe contains 30 mg benralizumab in 1 mL (30 mg/mL).

Each FASENRA PEN prefilled pen contains 30 mg benralizumab in 1 mL (30 mg/mL).

For the full list of excipients, see Section 6.1 List of excipients.

3 PHARMACEUTICAL FORM

Solution for injection

Clear to opalescent, colourless to yellow solution for injection in a prefilled syringe (FASENRA) or prefilled pen (FASENRA PEN).

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Asthma

FASENRA is indicated as add-on therapy in patients aged 12 years and over with severe eosinophilic asthma (blood eosinophil count ≥ 300 cells/ μ L or ≥ 150 cells/ μ L if on oral corticosteroid treatment) (see Section 5.1 Pharmacodynamic properties [*Clinical Trials*]).

Relapsed or refractory Eosinophilic granulomatosis with polyangiitis (EGPA)

FASENRA is indicated as an add-on treatment for relapsing or refractory Eosinophilic Granulomatosis with Polyangiitis (EGPA) in adult patients aged 18 years and over (see Section 5.1 Pharmacodynamic properties [*Clinical Trials*]).

Hypereosinophilic Syndrome (HES)

FASENRA is indicated as add-on therapy for patients aged 12 years and older with inadequately controlled hypereosinophilic syndrome without an identifiable non-haematologic secondary cause.

4.2 Dose and method of administration

FASENRA should be prescribed by a health care professional in consultation with a specialist physician experienced in the diagnosis and treatment of conditions for which FASENRA is indicated (see Section 4.1 Therapeutic indications).

FASENRA is intended for long-term treatment. A decision to continue therapy should be made at least annually based on disease severity and level of disease control.

Asthma

Treatment with high-dose inhaled corticosteroids (ICS) and long-acting β -agonists (LABA) should be optimised prior to commencement of treatment with FASENRA.

Adults and adolescents (12 years and over)

The recommended dose is 30 mg of FASENRA by subcutaneous injection every 4 weeks for the first 3 doses, and then every 8 weeks thereafter.

EGPA

The recommended dosage is 30 mg of FASENRA administered by subcutaneous injection every 4 weeks.

HES

The recommended dosage is 30 mg of FASENRA administered by subcutaneous injection every 4 weeks.

Missed Dose

If an injection is missed on a planned date, dosing should resume as soon as possible on the indicated regimen; a double dose must not be administered.

Special patient populations

Renal impairment

No dose adjustment is required for patients with renal impairment (see Section 5.2 Pharmacokinetic properties [*Special patient populations/Renal impairment*]).

Hepatic impairment

No dose adjustment is required for patients with hepatic impairment (see Section 5.2 Pharmacokinetic properties [*Special patient populations/Hepatic impairment*]).

Use in the elderly

No dose adjustment is required for elderly patients (see Section 5.2 Pharmacokinetic properties [*Special patient populations/Elderly (≥ 65 years old)*]).

Paediatric use

The safety and efficacy of FASENRA in children with asthma below 12 years of age have not been established.

The safety and efficacy of FASENRA in children and adolescents (below 18 years of age) with EGPA have not been established.

The safety and efficacy of FASENRA in children with HES below 12 years of age have not been established.

Method of administration

FASENRA is for single use in one patient only. Discard any residue.

FASENRA is administered as a subcutaneous injection.

Administer FASENRA into the thigh or abdomen. If somebody else administers the injection, the upper arm can also be used. Do not administer into areas where the skin is tender, bruised, erythematous, or hardened.

Instructions for administration

Do not shake. Do not use if frozen.

Prior to administration, if the FASENRA prefilled syringe/FASENRA PEN has been in the refrigerator allow FASENRA to reach room temperature (approximately 30 minutes).

Visually inspect FASENRA for particulate matter and discolouration prior to administration. FASENRA is clear, colourless to yellow, and may contain translucent or white to off white particles. Do not use FASENRA if liquid is cloudy, discoloured, or if it contains large particles or foreign particulate matter.

Additional information and instructions for the preparation and administration of FASENRA using the prefilled syringe or prefilled pen (FASENRA PEN) are given in the *Instructions for Use* booklet provided in each pack.

4.3 Contraindications

FASENRA is contraindicated in patients who have known hypersensitivity to benralizumab or any of its excipients (see Section 4.4 Special warnings and precautions for use [*Hypersensitivity reactions*]).

4.4 Special warnings and precautions for use

FASENRA should not be used to treat acute asthma exacerbations.

Patients should be instructed to seek medical advice if their asthma remains uncontrolled or worsens after initiation of treatment.

Reduction in oral corticosteroids (OCS) dose, if appropriate, should be gradual and performed under the supervision of a physician. Abrupt discontinuation of OCS after initiation of FASENRA therapy is not recommended.

Hypersensitivity reactions

Hypersensitivity reactions (eg anaphylaxis, angioedema, urticaria, urticaria papular, rash) have occurred following administration of FASENRA (see Section 4.8 Adverse effects (Undesirable effects)). These reactions may occur within hours of administration, but in some instances have a delayed onset (ie days). A history of serious hypersensitivity/anaphylaxis unrelated to benralizumab may be a risk factor for anaphylaxis following FASENRA administration.

In the event of a hypersensitivity reaction, FASENRA should be discontinued and appropriate therapy initiated.

Parasitic (helminth) infection

Eosinophils may be involved in the immunological response to some helminth infections. Patients with known helminth infections were excluded from participation in clinical trials. It is unknown if FASENRA may influence a patient's response against helminth infections.

Treat patients with pre-existing helminth infections before initiating therapy with FASENRA. If patients become infected while receiving treatment with FASENRA and do not respond to anti-helminth treatment, discontinue treatment with FASENRA until infection resolves.

Use in the elderly

No dose adjustment is required for elderly patients (see Section 4.2 Dose and method of administration [*Special patient populations/Use in the elderly*] and Section 5.2 Pharmacokinetic properties [*Special patient populations/ Elderly (≥65 years old)*]).

Paediatric use

The safety and efficacy of FASENRA in children below 12 years of age have not been established.

In Phase 3 studies SIROCCO/CALIMA, the treatment responses in adolescent patients (12 to 17 years of age) were less than that observed in adults, however they were not powered to detect a response in this sub-group. The adverse event profile in adolescents was generally similar to the overall population in these studies of up to 56 weeks duration, as well as in a 108-week extension study (BORA).

Effects on laboratory tests

As expected based on the mechanism of action of benralizumab, in the pivotal Phase 3 trials, following subcutaneous administration of benralizumab at the recommended dose blood eosinophils were reduced to a median absolute blood eosinophil count of 0 cells/ μ L. This magnitude of reduction was seen at the first observed time point, 4 weeks of treatment, and was maintained throughout the treatment period. Basophils also express IL-5R α , and were also reduced.

4.5 Interactions with other medicines and other forms of interactions

In a randomised, double-blind parallel-group study of 103 patients aged between 12 and 21 years with severe asthma, the humoral antibody responses induced by seasonal influenza virus vaccination were similar between benralizumab 30 mg and placebo. Vaccine efficacy was not directly assessed in this study.

Cytochrome P450 enzymes, efflux pumps and protein-binding mechanisms are not involved in the clearance of benralizumab. There is no evidence of IL-5R α expression on hepatocytes. Eosinophil depletion does not produce chronic systemic alterations of proinflammatory cytokines.

An effect of benralizumab on the pharmacokinetics of co-administered medications is not expected. Based on population pharmacokinetic analysis, commonly co-administered medications had no effect on the benralizumab clearance in patients with asthma.

4.6 Fertility, pregnancy and lactation

Effects on fertility

No fertility studies have been conducted with benralizumab in humans or animals.

Examination of surrogate fertility parameters (including organ weights and histopathology of reproductive tissues) in male and female cynomolgus monkeys treated with benralizumab at intravenous doses up to 25 mg/kg or at subcutaneous doses of up to 30 mg/kg once every 2 weeks for 9 months suggested no impairment of fertility. Systemic exposure (serum AUC) at the no adverse effect dose in monkeys was more than 400 times that in patients at the maximum recommended human dose.

Use in pregnancy – Category B1

It is preferable to avoid the use of FASENRA during pregnancy, especially during the third trimester due to the potential for eosinophil depletion in the newborn. Administration of FASENRA

to pregnant women should only be considered if the expected benefit to the mother is greater than any possible risk to the fetus.

IgG antibodies such as benralizumab are increasingly transported across the placenta as pregnancy progresses; therefore, greater fetal exposure occurs in the third trimester of pregnancy.

The effect of FASENRA on human pregnancy is unknown.

No adverse effects on pre- and postnatal survival, growth or development were observed in cynomolgus monkeys with maternal administration of intravenous doses of benralizumab (10 or 30 mg/kg) once every 2 weeks from early pregnancy (gestation day 20–22) to 1-month postpartum. Serum benralizumab levels in infants were 66% of maternal levels 7 days post-partum and declined over time (eg 10% of maternal levels 3 months post-partum). While there was no effect observed on the primary and secondary humoral immune responses to immunisation or levels of serum IgM, IgG and IgA in the offspring of the treated monkeys, there was a marked eosinophil depletion, consistent with significant placental transfer of benralizumab that resulted in pharmacologically active drug levels in infants. The immunological response to parasitic infection was not examined in the offspring.

Use in lactation

It is unknown whether benralizumab is excreted in human milk. Since antibodies can be secreted in human milk a risk to the breastfed child cannot be excluded.

A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from benralizumab therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the mother.

4.7 Effects on ability to drive and use machines

FASENRA has no or negligible influence on the ability to drive and use machines.

4.8 Adverse effects (Undesirable effects)

Clinical trials experience - asthma

A total of 895 patients received the recommended dose of FASENRA (30 mg every 4 weeks for the first 3 doses, and then every 8 weeks thereafter) within the 3 placebo-controlled Phase 3 clinical trials – 822 patients (including 38 adolescents) in SIROCCO and CALIMA (48 and 56-week duration respectively) and 73 patients (adults only) in a 28-week OCS sparing trial (ZONDA). Note - all 3 trials assessed 2 dosing regimens of FASENRA compared to placebo, however only the safety data for the recommended dosing regimen (Q8w; see Section 4.2 Dose and method of administration) has been presented below.

In clinical trials in patients with severe asthma with eosinophilic phenotype the most commonly reported adverse drug reactions during treatment were headache and pharyngitis.

Phase 3 exacerbation trials (SIROCCO and CALIMA)

Table 1 presents the most common ($\geq 3\%$ frequency) adverse events regardless of causality from the two placebo-controlled trials (SIROCCO/CALIMA) in patients receiving the recommended dose (Q8w) of benralizumab. Table 2 presents the SIROCCO/CALIMA adverse drug reactions reported at a frequency less than 3%. All patients (including placebo) were on a background of high-dose ICS/LABA.

Table 1 The most frequent ($\geq 3\%$) adverse events regardless of causality reported during the on-treatment period and more common than placebo (SIROCCO/CALIMA - safety analyses set)

Preferred term	FASENRA Q8w (%) (N=822)	Placebo (%) (N=847)
Headache ^a	8.6	6.3
Pharyngitis ^b	4.0	2.5
Arthralgia	3.9	2.2
Cough	3.3	2.1

^a Headache was also identified as an adverse drug reaction.

^b Pharyngitis was also identified as an adverse drug reaction which was defined by the following grouped preferred terms 'Pharyngitis', 'Pharyngitis bacterial', 'Viral pharyngitis', 'Pharyngitis streptococcal' (grouped adverse drug reaction rate of 5.0% FASENRA and 3.4% placebo).

Table 2 Adverse drug reactions reported at a frequency less than 3% (SIROCCO/CALIMA - safety analyses set)

Frequency	System Organ Class	Event
Common ($\geq 1\%$ - $< 10\%$ *)	<i>Immune system disorders</i>	Hypersensitivity reactions ^a
	<i>General disorders & administration site conditions</i>	Pyrexia ^b ; Injection site reactions ^c

* Definition of common, however only adverse reactions less than 3% included within this table. See above for those $\geq 3\%$.

^a In placebo-controlled trials (SIROCCO and CALIMA), hypersensitivity reactions (defined as urticaria, urticaria papular and rash) occurred at a rate of 3.2% in both patients treated with the indicated benralizumab dose and those treated with placebo. See Section 4.4 Special warnings and precautions for use [*Hypersensitivity reactions*].

^b Pyrexia was also reported as an adverse drug reaction in SIROCCO and CALIMA with a rate of 2.9% and 1.7% in patients treated with benralizumab and placebo respectively.

^c In placebo-controlled trials (SIROCCO and CALIMA), injection site reactions (eg pain, erythema, pruritus, papule) occurred at a rate of 2.2% in patients treated with the indicated benralizumab dose compared with 1.9% in patients treated with placebo.

Adolescents (12 to 17 years of age)

The frequency, type and severity of adverse drug reactions in the adolescent population were observed to be similar to those seen in adults.

Phase 3 OCS sparing trial (ZONDA)

In ZONDA all patients were taking daily OCS (7.5 to 40 mg per day) in addition to regular use of high-dose ICS/LABA. In general, the safety results in ZONDA were similar to those observed in SIROCCO and CALIMA.

Long-term safety

In a 56-week double blind, randomized, parallel-group extension trial (BORA) in patients with asthma from SIROCCO, CALIMA and ZONDA trials, 842 patients were treated with FASENRA at the recommended dose and remained in the trial. The overall adverse event profile was similar to the asthma trials described above. Additionally, in an open-label safety extension trial (MELTEMI) in patients with asthma from previous trials, 226 adult patients were treated with FASENRA at the recommended dose for up to 42.3 months. Combined with the treatment period in previous studies, this corresponds to median follow-up of 3.4 years (range 8.5 months – 5.3 years). The safety profile during this follow-up period was consistent with the known safety profile of FASENRA.

Adolescent patients aged 12 to 17 (n=61) from SIROCCO and CALIMA continued treatment with FASENRA (at the recommended dose) in BORA for up to 108 weeks. Safety was consistent with the predecessor trials.

Clinical trials experience – EGPA

The safety profile of a total of 70 patients with EGPA, who received FASENRA 30 mg every 4 weeks during an active-controlled Phase 3 clinical study (MANDARA) of 52 weeks duration, was similar to the established safety profile of FASENRA. This incidence of adverse reactions were similar to those reported in asthma, with the exception of headache, which occurred in 17% of FASENRA-treated patients compared to 8% in asthma. No additional adverse reactions were identified.

Clinical trials experience – HES

The safety of FASENRA is based on 67 adult and adolescent patients, who received at least 1 dose of FASENRA 30 mg every 4 weeks in a placebo-controlled study of 24 weeks duration (NATRON). The safety profile of FASENRA for HES was consistent with the established safety profile from asthma studies, with the exception of headache, which occurred in 16% of FASENRA-treated patients with HES. No new adverse reactions were identified.

Immunogenicity

Treatment with benralizumab, like other monoclonal antibodies, may result in an anti-drug antibody (ADA) response (see Section 5.1 Pharmacodynamic properties [*Immunogenicity*]). However, there is no apparent correlation of ADA development to efficacy or adverse events.

Post marketing experience

In addition to the above, Table 3 describes adverse reactions that have been identified during post-approval use of FASENRA.

Table 3 Post marketing adverse reactions

Frequency	System Organ Class	Event
Not known ^a	<i>Immune system disorders</i>	Anaphylaxis ^b

^a It is generally not possible to reliably determine the frequency because such reactions have been reported spontaneously from a population of uncertain size and therefore represent reporting rates. The frequency of these adverse reactions is therefore 'not known' (cannot be estimated from available data).

^b Anaphylaxis was defined by the following grouped preferred terms: 'Anaphylactic reaction', 'Angioedema'.

Reporting suspected adverse effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at www.tga.gov.au/reporting-problems.

4.9 Overdose

Doses of up to 200 mg were administered subcutaneously in clinical trials to patients with eosinophilic disease without evidence of dose-related toxicities.

There is no specific treatment for an overdose with benralizumab. If overdose occurs, the patient should be treated supportively with appropriate monitoring as necessary.

For information on the management of overdose, contact the Poison Information Centre on 131126 (Australia).

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Mechanism of action

Benralizumab is an antibody that binds to the alpha subunit of the human interleukin-5 receptor (IL-5R α) with high affinity (16 pM) and specificity. The IL-5 receptor is specifically expressed on the surface of eosinophils and basophils. The absence of fucose sugar units in the Fc domain of benralizumab results in high affinity (45.5 nM) for Fc γ RIII receptors on immune effectors cells such as natural killer (NK) cells leading to apoptosis of eosinophils and basophils through enhanced antibody-dependent cell-mediated cytotoxicity (ADCC).

Eosinophilic inflammation is an important component in the pathogenesis of asthma, EGPA and HES. Eosinophils are a rich source of proinflammatory mediators (eg eicosanoids, leukotrienes, cytokines) and granule proteins (eg eosinophil cationic protein, eosinophil peroxidase, eosinophil neurotoxin and major basic protein). Benralizumab, by enhanced ADCC, reduces eosinophilic inflammation.

Pharmacodynamic effects

The pharmacodynamic response (blood eosinophil depletion) following repeat subcutaneous dosing was evaluated in asthma patients in a 12-week Phase 2 trial. Patients with mild-moderate asthma received 1 of 3 doses of benralizumab [25 mg (n=7), 100 mg (n=6) or 200 mg (n=6) subcutaneous] or placebo (n=6) every 4 weeks for a total of 3 doses. Median blood eosinophil levels at baseline were 400, 200, 120 and 200 cells/ μ L in the 25, 100, and 200 mg benralizumab and placebo groups, respectively. Blood eosinophil depletion was observed following subcutaneous administration of benralizumab at all dose levels and no depletion was observed in the placebo group. Twenty-four hours post dosing, all benralizumab dosage groups demonstrated complete or near complete depletion of median blood eosinophil levels (0, 0 and 5 cells/ μ L respectively). There were no changes in median blood eosinophils in the placebo group. The effect on blood eosinophil depletion was maintained throughout the dosing period.

In a Phase 1 trial, the effect of benralizumab on eosinophils in airway mucosa was evaluated in asthmatic patients with 2.5% or more eosinophils in sputum. Patients received 100 or 200 mg subcutaneous benralizumab once every 4 weeks for 8 weeks (total benralizumab subcutaneous group n=9) or matching placebo (n=5). At the end of the 12-week treatment period, there was a median reduction from baseline in eosinophils in the airway mucosa of 96% in the total benralizumab subcutaneous group compared to a 47% reduction from baseline in the placebo group which was statistically significant (p=0.039).

In the Phase 1 trial, treatment with benralizumab was also associated with reductions in blood basophils, and in both Phase 1 and 2 trials eosinophil granule products such as serum eosinophil derived neurotoxin (EDN) and eosinophil cationic protein (ECP).

In the two pivotal Phase 3 asthma trials (SIROCCO and CALIMA), following subcutaneous administration of benralizumab at the recommended dose blood eosinophils were reduced to a median absolute blood eosinophil count of 0 cells/ μ L, which corresponds to a median reduction of 100% (see Section 5.1 Pharmacodynamic properties [*Clinical trials*] below). This magnitude of reduction was seen at the first observed time point, 4 weeks of treatment, and was maintained throughout the treatment period. Maintenance of eosinophil depletion was observed throughout the 56-week extension phase (BORA extension trial) consistent with previous trials.

In patients with EGPA, depletion of blood eosinophils was consistent with the effect observed in the asthma trials. Blood eosinophil depletion was seen at the first observed time point, 1 week of treatment, and was maintained throughout the 52-week treatment period.

In patients with HES, depletion of blood eosinophils was consistent with the effect observed in asthma and EGPA trials. Blood eosinophil depletion was seen at the first observed timepoint (Week 4) and was maintained over the 24-week treatment period.

Immunogenicity

Overall, treatment-emergent anti-drug antibody (ADA) response developed in 107 out of 809 (13%) of asthma patients treated with FASENRA at the recommended dosing regimen during the 48 to 56-week treatment period. In a majority of the ADA positive patients, in vitro neutralizing antibodies were detected. Anti-benralizumab antibodies were associated with increased clearance of benralizumab and increased blood eosinophil levels in patients with high ADA titers compared to antibody negative patients. No evidence of an association of ADA with efficacy or safety was observed.

Following a second year of treatment of these asthma patients from the Phase 3 placebo-controlled trials, an additional 18 out of 510 (4%) had newly developed treatment-emergent antibodies. Overall, in patients who were ADA positive in the predecessor trials, titers remained stable or declined in the second year of treatment. Consistent with predecessor trials, no evidence of an association of ADA with efficacy or safety was observed.

In patients with EGPA, treatment-emergent ADA response developed in 6 out of 67 (9%) patients treated with FASENRA during the Phase 3 active-controlled 52-weeks treatment period. Neutralising antibody activity was detected in one of the ADA positive patients.

In patients with HES, treatment-emergent ADA response developed in 7 out of 66 (11%) patients treated with benralizumab during the Phase 3 placebo-controlled 24-week treatment period. A total of 2 patients that developed treatment-emergent ADA following benralizumab administration, also developed neutralising antibodies.

The data reflect the percentage of patients whose test results were positive for antibodies to benralizumab in specific assays. The observed treatment-emergent ADA response is highly dependent on several factors, including assay sensitivity and specificity, assay methodology, sample handling, timing of sample collection, concomitant medication, and underlying disease. For these reasons, comparison of the treatment-emergent ADA response to benralizumab with the treatment-emergent ADA response to other products may be misleading.

Clinical trials

Severe asthma

The safety and efficacy of FASENRA as an add-on therapy in patients with severe asthma were evaluated in 3 randomised, double-blind, parallel-group, placebo-controlled Phase 3 clinical trials:

- Two replicate long-term exacerbation trials in adults and adolescents (12 years and older) with 48 and 56 weeks duration (SIROCCO and CALIMA respectively); and
- One 28-week OCS reduction trial in adults (18 years and over (ZONDA)).

While all 3 trials assessed 2 dosing regimens compared to placebo, the recommended dosing regimen is FASENRA administered every 4 weeks for the first 3 doses, then every 8 weeks thereafter (referred herein as Q8w; see Section 4.2 Dose and method of administration) as no

additional benefit was observed with the more frequent dosing regimen (administered every 4 weeks). Only the results for the recommended Q8w dosing regimen have been presented below.

A total of 805, 881 and 148 patients were randomised to treatment with FASENRA Q8w and placebo in SIROCCO, CALIMA and ZONDA respectively. This included 84 adolescents (SIROCCO/CALIMA combined; 38 within the FASENRA Q8w arms and 4 within the placebo arms). See Table 4 for further details.

Table 4 Randomised patient numbers (N) for the pivotal phase III studies (FASENRA Q8w and placebo arms only)

	SIROCCO	CALIMA	ZONDA
Total randomised population*	805	881	148
FASENRA Q8w	398	441	73
Placebo	407	440	75
High dose ICS/LABA & ≥ 300 cells/ μ L eosinophil count**	534	487	-
FASENRA Q8w	267	239	-
Placebo	267	248	-
High dose ICS/LABA & < 300 cells/ μ L eosinophil count^	271	247	-
FASENRA Q8w	131	125	-
Placebo	140	122	-

ICS – inhaled corticosteroids; LABA – long acting β -agonists; *all high-dose ICS/LABA patients regardless of baseline blood eosinophil count = Full analysis set (FAS) for SIROCCO/CALIMA and ITT population for ZONDA; **SIROCCO/CALIMA enriched ITT population - high dose ICS/LABA and ≥ 300 cells/ μ L baseline blood eosinophil count; ^prespecified analysis population (Note: patients aged 12–17 years could have received medium-dose or high-dose ICS).

Table 5 Key demographic and baseline characteristics of the pivotal phase III placebo-controlled asthma studies (ITT populations only)

	CALIMA		SIROCCO		ZONDA	
	FASENRA	Placebo	FASENRA	Placebo	FASENRA	Placebo
Mean age (years) [range]	49.6 [12, 74]	48.5 [12, 75]	47.6 [12, 74]	48.6 [12, 75]	52.9 [27, 75]	49.9 [21, 74]
Mean baseline eosinophil count (cells/ μ L) [range]	621 [300, 2600]	642 [300, 4494]	620 [300, 3100]	621 [300, 2690]	509 [154, 2140]	656 [160, 4550]
Mean pre-bronchodilator FEV ₁ (L) [range]	1.8 [0.6, 3.8]	1.8 [0.6, 3.8]	1.7 [0.5, 3.5]	1.7 [0.5, 3.5]	1.8 [0.6, 3.6]	1.9 [0.7, 3.8]
Mean number of exacerbations in previous year [range]	2.7 [2, 11]	2.8 [2, 18]	2.8 [2, 11]	3.1 [2, 15]	3.1 [1, 15]	2.5 [1, 10]
Mean ACQ-6 score [range]	2.8 [0.2, 5.7]	2.8 [0.2, 5.8]	2.8 [0.2, 5.7]	2.9 [0.5, 5.5]	2.4 [0.0, 5.2]	2.7 [0.8, 5.0]
Mean ICS dose (μ g) [range]	1002.3 [250, 4750]	929.5 [250, 3000]	913.4 [500, 2500]	901.6 [125, 3000]	1191.6 [100, 3250]	1232.3 [100, 5000]

ICS – inhaled corticosteroids (dose converted to fluticasone propionate dry powder equivalent); FEV₁ – forced expiratory volume in 1 second.

SIROCCO and CALIMA

Patients enrolled into SIROCCO and CALIMA were required to have a history of ≥ 2 asthma exacerbations requiring oral or systemic corticosteroid treatment in the past 12 months, an Asthma Control Questionnaire-6 (ACQ-6) score of ≥ 1.5 at screening and reduced lung function at baseline [pre-bronchodilator forced expiratory volume in 1 second (FEV₁) $< 80\%$ in adults and $< 90\%$ in adolescents] despite treatment with high-dose ICS (SIROCCO) or with medium- or high-dose ICS

(CALIMA) and their current standard of care. Medium- and high-dose ICS were defined as ≥ 250 μg and ≥ 500 $\mu\text{g/day}$ fluticasone propionate dry powder formulation or equivalent respectively. The medium-dose ICS arm in CALIMA was assessed as a descriptive analysis only and has not been discussed further.

Patients were stratified 2:1 according to baseline blood eosinophils count (≥ 300 or < 300 cells/ μL). The primary efficacy population (intent-to-treat (ITT)) in both studies was patients with a high-dose ICS/LABA and a baseline blood eosinophil count ≥ 300 cells/ μL . Prespecified analyses were also conducted on the high-dose ICS/LABA populations with a baseline blood eosinophil count < 300 cells/ μL (patients not included within the ITT population) and the full analysis set (FAS) for predefined eosinophil ranges. Table 4 provides a summary of the randomised patient numbers for the different analysis groups.

ITT population (high-dose ICS/LABA and baseline blood eosinophil count ≥ 300 cells/ μL)

The primary endpoint for both trials was the annual asthma exacerbation rate ratio versus placebo within the primary efficacy population (ITT). Key secondary endpoints were FEV₁ and total asthma symptom score. Other lung function, symptom control and quality of life measures were also assessed.

Demographic, key respiratory and other baseline disease characteristics were balanced across the ITT treatment groups (FASENRA Q8w and placebo) in both trials. The demographic and patient characteristics in the ITT population were generally similar to the overall population for the two treatment arms, as well as the < 300 cells/ μL subgroup.

Treatment with FASENRA Q8w significantly reduced the annual rate of exacerbations (primary endpoint) by 51% (SIROCCO) and 28% (CALIMA) compared to placebo (see Table 6). Similar reductions were observed compared to placebo for exacerbations requiring hospitalisations only (52%) and emergency room visits only (77%) in SIROCCO. In CALIMA, there were too few events in the placebo treatment arm to draw conclusions for exacerbations requiring hospitalisation or emergency room visits.

Clinically and statistically significant improvements in lung function were also observed with a 0.159 L (SIROCCO) and 0.116 L (CALIMA) increase in pre-bronchodilator FEV₁ relative to placebo (see Table 6). Compared with placebo, FASENRA Q8w provided consistent improvements over time in the mean change from baseline in FEV₁. Benefits of FASENRA Q8w on lung function were further supported by improvements in the mean change for morning and evening peak expiratory flow (PEF) from baseline compared to placebo.

Significant improvements were also observed for total asthma symptoms score and the Asthma Control Questionnaire (ACQ-6) after treatment with FASENRA Q8w relative to placebo (see Table 6). Significant improvements in quality of life, as measured by the Standardised Asthma Quality of Life Questionnaire for 12 years and older (AQLQ(S)+12), were also demonstrated for FASENRA Q8w compared to placebo (see Table 6).

Table 6 Overview of key efficacy results (ITT population – high-dose ICS/LABA & baseline blood eosinophils ≥ 300 cells/ μ L) – SIROCCO and CALIMA

FASENRA Q8w vs Placebo					
SIROCCO			CALIMA		
N(n) v N(n)	Comparison (95% CI)	p-value	N(n) v N(n)	Comparison (95% CI)	p-value
Primary endpoint: Annual asthma exacerbation^a rate (rate ratio)^a					
267 (267) v 267 (267)	0.49 (0.37, 0.64)	<0.001*	239 (239) v 248 (248)	0.72 (0.54, 0.95)	0.019*
Pre-bronchodilator FEV₁ (L) change from baseline at EOT (difference in LS means)^b					
264 (235) v 261 (233)	0.159 (0.068, 0.249)	0.001*	238 (211) v 244 (221)	0.116 (0.028, 0.204)	0.010*
Total asthma symptom score change from baseline at EOT (difference in LS means)^c					
263 (178) v 267 (180)	-0.25 (-0.45, -0.06)	0.012*	237 (185) v 247 (187)	-0.23 (-0.43, -0.04)	0.019*
ACQ-6 score change from baseline at EOT (difference in LS means)^d					
267 (263) v 267 (267)	-0.29 (-0.48, -0.10)	0.003	239 (239) v 248 (247)	-0.25 (-0.44, -0.07)	0.008
Overall AQLQ(S)+12 score change from baseline at EOT (difference in LS means)^e					
267 (252) v 267 (254)	0.30 (0.10, 0.50)	0.004	239 (230) v 248 (240)	0.24 (0.04, 0.45)	0.019

^a Exacerbations were defined as worsening of asthma requiring use of oral/systemic corticosteroids for ≥ 3 days and/or emergency department visits requiring use of oral/systemic corticosteroids and/or hospitalisation. For patients on maintenance OCS, an exacerbation requiring OCS was defined as a temporary increase in stable oral/systemic corticosteroids for at ≥ 3 days or a single depo-injectable dose of corticosteroids.

* Statistically significant under the multiple testing procedure.

^a The crude placebo rate was 1.53 in SIROCCO and 1.03 in CALIMA

^b The LS mean change from baseline at EOT for placebo was 0.239 L in SIROCCO and 0.215 L in CALIMA

^c The LS mean change from baseline at EOT for placebo was -1.04 units in SIROCCO and -1.16 units in CALIMA

^d The LS mean change from baseline at EOT for placebo was -1.17 units in SIROCCO and -1.19 units in CALIMA

^e The LS mean change from baseline at EOT for placebo was 1.26 units in SIROCCO and 1.31 units in CALIMA


End of treatment (EOT) was Week 48 in Trial 1 and Week 56 in Trial 2.

CI – Confidence interval; EOT – End of treatment; FEV₁ – Forced expiratory volume in 1 second; LS – Least squares; N – Number of patients in the analysis; n – Number of patients with data at EOT.

Baseline blood eosinophil subgroup analyses

While efficacy benefits were observed with FASENRA compared to placebo irrespective of baseline eosinophil count, increasing baseline eosinophil counts were identified as a potential predictor of improved treatment response (see Figure 1).

Figure 1 Annual asthma exacerbation (primary endpoint) rate ratio comparison by baseline blood eosinophil count category, negative binomial model - forest plot (Integrated SIROCCO/CALIMA - Full analysis set, high-dose ICS)



Prior exacerbation history

A subgroup analysis on prior exacerbation history within the ITT population indicated that a higher prior exacerbation history may also be a potential predictor of an improved treatment response. When considered alone, or in combination with a higher baseline eosinophil count, these factors may further identify patients who may achieve a greater response from treatment with FASENRA.

OCS dose reduction trials (ZONDA and PONENTE)

ZONDA, a placebo-controlled study, and PONENTE, an open-label study, evaluated the effect of FASENRA on reducing the use of maintenance OCS.

ZONDA

ZONDA included patients who were treated with daily OCS (7.5 to 40 mg/day) in addition to regular use of high-dose ICS/LABA with or without additional controller(s) to maintain asthma control. The definition of high-dose ICS was as per SIROCCO/CALIMA. There was an 8-week run-in period during which a patient's OCS dose was titrated to the minimum effective dose while maintaining asthma control. The baseline median OCS dose was 10 mg (range: 8-40 mg) for both treatment groups. Patients were also required to have blood eosinophil counts $\geq$ 150 cells/ μ L and a history of at least one exacerbation in the past 12 months. All patients were included in the analysis, including 20 patients with a baseline blood eosinophil count of $\geq$ 150-299 cells/ μ L (12 in FASENRA Q8w arm and 11 in the placebo arm).

Demographic, key respiratory and other baseline disease characteristics were balanced across the treatment groups (FASENRA Q8w and placebo).

The primary endpoint was percent (%) reduction from baseline of the final OCS dose during Weeks 24 to 28, while maintaining asthma control. Compared to placebo, patients receiving FASENRA Q8w achieved greater reductions in daily maintenance OCS dose while still maintaining asthma control. Reductions of $\geq 50\%$ in the OCS dose were observed in 66% of patients receiving FASENRA Q8w compared to 37% for the placebo arm (see Table 7). The proportion of patients with a mean final OCS dose ≤ 5 mg at Weeks 24 to 28 were 59% for FASENRA Q8w and 33% for placebo (odds ratio 2.74 (95% CI: 1.41, 5.31), $p=0.002$). Patients with an optimised baseline OCS dose (12.5 mg or less) were eligible to achieve a 100% reduction in OCS dose during the study. A significant difference was observed in the percentage of eligible patients who achieved 100% reduction with FASENRA Q8w compared to placebo (52.4% vs 19.0%; Odds ratio 4.19 (95% CI: 1.58, 11.12), $p=0.002$).

Table 7 Effect of FASENRA Q8w on OCS dose reduction - ZONDA

	FASENRA (n=73)	Placebo (n=75)
Wilcoxon rank sum test (primary analysis method)		
Median % reduction in daily OCS dose from baseline (95% CI)	75 (60, 88)	25 (0, 33)
Wilcoxon rank sum test p-value	<0.001	
Proportional odds model (sensitivity analysis)		
Percent reduction in OCS from baseline at Week 28		
≥90% reduction	27 (37%)	9 (12%)
≥75% reduction	37 (51%)	15 (20%)
≥50% reduction	48 (66%)	28 (37%)
>0% reduction	58 (79%)	40 (53%)
No change or no decrease in OCS	15 (21%)	35 (47%)
Odds ratio (95% CI), p-value	4.12 (2.22, 7.63), p<0.001	

OCS – oral corticosteroids; CI – confidence intervals; n – Number of patients in the analysis.

FASENRA Q8w demonstrated a 70% reduction in the annual asthma exacerbation rate over 28 weeks compared with placebo (Rate ratio: 0.30 (95% CI: 0.17, 0.53), $p<0.001$), and a 93% reduction in the annual asthma exacerbation rate associated with an emergency room visit or hospitalisations (Rate ratio: 0.07 (95% CI: 0.01, 0.63), $p=0.018$) over 28 weeks compared with placebo.

PONENTE

PONENTE enrolled 598 adult patients with severe asthma (blood eosinophil count ≥ 150 cells/ μ L at entry or ≥ 300 cells/ μ L in the past 12 months if study entry count was < 150 cells/ μ L) who were OCS-dependent. The primary endpoints included the proportion of patients who achieved a final OCS dose less than or equal to 5 mg while maintaining asthma control and taking into account adrenal function. The proportion of patients who achieved an OCS dose less than or equal to 5 mg (while maintaining asthma control and not limited by adrenal function) was 81.9%. Effects on OCS reduction were similar irrespective of blood eosinophil count at study entry (including patients with blood eosinophils < 150 cells/ μ L) and maintained over an additional period of 24 to 32 weeks.

Long term data

The long-term safety and efficacy of FASENRA was evaluated in a double-blind, randomised, parallel group, Phase 3, 56-week extension trial (BORA) in adult and adolescent (aged 12 years and older) patients from the predecessor studies - SIROCCO, CALIMA and ZONDA. The long-term safety of FASENRA was evaluated in an open-label safety extension trial MELTEMI (see Section 4.8 Adverse effects (Undesirable effects)).


BORA assessed the long-term effect of FASENRA on annual asthma exacerbation rate, lung function and ACQ-6, AQLQ(S)+12 at the 2 dosing regimens studied in the predecessor studies.

A total of 543 adult and adolescent patients were randomised to receive FASENRA Q8w in the predecessor trials and were treated and remained in the BORA extension trial – 512 from SIROCCO/CALIMA (full analysis set) and 31 from ZONDA.

The reduction in annual rate of asthma exacerbations observed in the placebo-controlled predecessor SIROCCO and CALIMA trials in patients with baseline blood eosinophil counts of ≥ 300 cells/ μ L who were taking high-dose ICS (ITT population) was maintained over the second year of treatment. In patients who received FASENRA in predecessor SIROCCO and CALIMA trials, 73% were exacerbation free in the extension BORA trial. Similar maintenance of effect was observed throughout BORA in lung function, ACQ-6 and AQLQ(S)+12.

In a pre-specified pooled analysis, maintenance of the reduction in daily OCS dose was also observed over the extension trial in patients enrolled from ZONDA (Figure 2).

Figure 2 Median percent reduction in daily OCS over time (ZONDA and BORA)^a



^a Predecessor ZONDA trial patients who continued FASENRA treatment into BORA. Patients were permitted to enter a second extension trial after a minimum of 8 weeks in BORA without completing the 56-week extension period.

MELTEMI was designed as an open-label safety extension study, enrolling adult patients who completed at least 16 weeks in ZONDA (see Section 4.8 Adverse effects (Undesirable effects)). The safety of the 2 dosing regimens studied in the predecessor studies was evaluated in 446 patients who received at least 1 dose of FASENRA.

Eosinophilic granulomatosis with polyangiitis (EGPA)

MANDARA

The efficacy and safety of FASENRA was evaluated in a randomised, double-blind, active-controlled, non-inferiority clinical trial of 52-weeks treatment duration, in patients aged 18 years and older with EGPA. A total of 140 patients who has a history of relapsing or refractory disease were randomised. FASENRA 30 mg was evaluated compared to mepolizumab 300 mg administered subcutaneously every 4 weeks in addition to background prednisolone/prednisone

with or without immunosuppressive therapy. The OCS dose was tapered at the discretion of the investigator.

The demographics and baseline characteristics are provided in Table 8.

Table 8 Demographics and Baseline Characteristics of EGPA Trial (MANDARA)

	Total Population (N=140)
Mean age (years)	52
Female (%)	60
White (%)	79
Time since diagnosis of EGPA, years, mean (SD)	5.2 (5.64)
History of ≥ 1 confirmed relapse in past 2 years (%)	79
Refractory disease (%)	60
Baseline oral corticosteroid ^a daily dose, mg, median (range)	10 (5–40)
Receiving immunosuppressive therapy ^b (%)	36
ANCA positive ^c (%)	29

SD – standard deviation.

^a Prednisone or prednisolone equivalent.

^b Azathioprine, methotrexate, mycophenolic acid.

^c Anti-neutrophil cytoplasmic antibody (ANCA) positive historically or at screening.

Remission

The primary endpoint was the proportion of patients in remission, defined as Birmingham Vasculitis Activity Score (BVAS)=0 (no active vasculitis) plus prednisolone/prednisone dose ≤ 4 mg/day, at both Week 36 and Week 48. As shown in Table 9, FASENRA demonstrated non-inferiority to mepolizumab for the primary endpoint. Results for accrued duration of remission and the components of remission are also shown in Table 9.

Table 9 Remission and Components of Remission in EGPA

	Remission (OCS≤ 4 mg/day + BVAS=0)		OCS≤ 4 mg/day		BVAS=0	
	FASENRA^a N=70	Mepo^b N=70	FASENRA^a N=70	Mepo^b N=70	FASENRA^a N=70	Mepo^b N=70
Patients in remission at both Weeks 36 and 48						
Patients, n (%) ^c	40 (58)	40 (57)	42 (61)	41 (58)	58 (83)	59 (84)
Differences in remission rate, (%) ^c (95% CI)	1.21 (-14.11, 16.53)	-- --	2.64 (-12.67, 17.95)	-- --	-1.17 (-13.27, 10.94)	-- --
Accrued duration over 52 weeks, n (%)						
0 weeks ^d	9 (13)	15 (21)	9 (13)	12 (17)	0	0
>0 to <12 weeks	13 (19)	10 (14)	11 (16)	12 (17)	0	2 (3)
12 to <24 weeks	8 (11)	8 (11)	9 (13)	8 (11)	2 (3)	2 (3)

	Remission (OCS≤4 mg/day + BVAS=0)		OCS≤4 mg/day		BVAS=0	
	FASENRA^a N=70	Mepo^b N=70	FASENRA^a N=70	Mepo^b N=70	FASENRA^a N=70	Mepo^b N=70
24 to <36 weeks	20 (29)	19 (27)	19 (27)	18 (26)	6 (9)	7 (10)
≥36 weeks	20 (29)	18 (26)	22 (31)	20 (29)	62 (89)	59 (84)
Odds ratio ^c (95% CI)	1.32 (0.72, 2.40)	-- --	1.27 (0.70, 2.31)	-- --	1.50 (0.54, 4.15)	-- --

N – number of patients in analysis.

^a FASENRA 30 mg administered every 4 weeks.

^b Mepolizumab (Mepo) 300 mg administered every 4 weeks.

^c Model adjusted percentages.

^d Did not achieve remission at any point.

^e An odds ratio >1 favours FASENRA.

The proportion of patients achieving remission within the first 24 weeks of treatment and remaining in remission through Week 52 was 42% for FASENRA and 37% for mepolizumab (difference in responder rate 5.54%, 95% CI: -9.30, 20.37).

Using an alternative remission definition of BVAS=0 plus prednisolone/prednisone ≤7.5 mg/day, a consistent efficacy between groups for these endpoints was observed.

Patients achieved the primary remission endpoint across the prespecified demographic and baseline characteristic subgroups.

Relapse

The hazard ratio for time to first relapse (defined as worsening related to vasculitis, asthma, or sino-nasal symptoms requiring an increase in dose of corticosteroids or immunosuppressive therapy or hospitalisation) was 0.98 (95% CI: 0.53, 1.82). Relapse was observed in 30% of patients on FASENRA and 30% of patients on mepolizumab. The annualised relapse rate was 0.50 for patients receiving FASENRA versus 0.49 for patients receiving mepolizumab (rate ratio 1.03, 95% CI: 0.56, 1.90). The types of relapse were consistent for patients receiving FASENRA or mepolizumab.

Oral corticosteroid reduction

The average daily OCS dose during Weeks 48 to 52 is presented in Table 10. A 100% reduction in the OCS dose was observed in 41% of patients receiving FASENRA compared to 26% of those receiving mepolizumab (difference 15.69%, 95% CI: 0.67, 30.71). Reductions of 50% or higher were observed in 85% of patients receiving FASENRA compared to 74% of those receiving mepolizumab (difference 10.79%, 95% CI: -2.25, 23.83).

Table 10 **Average Daily Oral Corticosteroid Dosing during Weeks 48 to 52 in EGPA**

	Number (%) of Patients	
	FASENRA ^a (N=70)	Mepolizumab ^b (N=70)
0 mg	29 (41)	19 (27)
>0 to ≤4.0 mg	19 (27)	30 (43)
>4.0 to ≤7.5 mg	15 (21)	13 (19)
>7.5 mg	7 (10)	8 (11)
Odds ratio ^c (95% CI)	1.38 (0.75, 2.54)	--

N – number of patients in analysis.

^a FASENRA 30 mg administered every 4 weeks.

^b Mepolizumab 300 mg administered every 4 weeks.

^c An odds ratio >1 favours FASENRA.

Asthma Control Questionnaire-6 (ACQ-6)

The ACQ-6 responder rate during Weeks 48 to 52 (defined as a decrease in score of 0.5 or more compared with baseline) was 42% for FASENRA and 48% for mepolizumab (difference -6.16%, 95% CI: -18.52, 6.21).

Hypereosinophilic syndrome (HES)

NATRON

The efficacy and safety of FASENRA was evaluated in a randomised, double-blind, parallel-group, placebo-controlled clinical trial of 24-week treatment duration, in patients aged 12 years and older with HES.

A total of 133 patients (129 adults and 4 adolescents) who had signs or symptoms of HES flare or had experienced at least two HES flares within the past 12 months received randomised treatment. At screening, patients had a blood eosinophil count ≥1,000 cells/μL and stable HES therapy for at least 4 weeks. HES therapy included OCS, immunosuppressive/cytotoxic therapy or other symptomatic therapies. Patients who were FIP1L1-PDGFRα kinase-positive were excluded from the study. Patients were randomised to FASENRA 30 mg or placebo administered subcutaneously every 4 weeks while continuing their stable HES therapy.


Table 11 **Demographics and Baseline Characteristics of the HES Trial**

	Total Population (N=133)
Mean age (years)	48
Female (%)	62
White (%)	76
Time since diagnosis of HES, years, mean (SD)	4.3 (6.0)
Eosinophil count at screening (local laboratory) (cells/μL), median	1600

The primary endpoint was time to first HES flare, defined as a HES clinical manifestation or laboratory abnormality that resulted in an increase or addition of OCS of 10 mg/day or more for at

least 2 days or, an increase or addition of new cytotoxic and/or immunosuppressive therapy or hospitalisation. Treatment with FASENRA significantly delayed the time to first HES flare (Figure 3) and resulted in a 65% reduction in the risk of first flare over the treatment period versus placebo (HR: 0.35, 95% CI: 0.18, 0.69, p=0.0024).

Figure 3 **Kaplan Meier curve for time to first HES flare versus placebo**



FASENRA significantly improved all key secondary endpoints in the HES trial (Table 12).

Table 12 Key Secondary Efficacy Variables, HES Trial (ITT Population)

	FASENRA^a (N=67)	Placebo (N=66)
Proportion of patients who experienced a HES flare		
Patients with ≥ 1 HES flare or who withdrew from study ^b , n (%)	15 (22)	30 (45)
Odds ratio ^c (95% CI)	0.31 (0.14, 0.69)	--
CMH p-value	0.003	--
Frequency of HES flares		
Patients with 0 flares, n (%)	54 (81)	38 (58)
Patients with 1 flare, n (%)	13 (19)	20 (30)
Patients with 2 flares, n (%)	0 (0)	8 (12)
Patients with ≥ 3 flares, n (%)	0 (0)	0 (0)
<i>Comparison (FASENRA versus placebo)</i>		
Rate/year	0.41	1.23
Rate ratio ^c (95% CI)	0.34 (0.18, 0.63)	--
p-value	0.0008	--
Time to first haematologic relapse^d		
Patients with haematologic relapse, n (%)	5 (7)	39 (59)
Hazard ratio ^c (95% CI)	0.08 (0.03, 0.20)	--
p-value	<0.0001	--
Change from baseline in fatigue severity based on PROMIS Fatigue^e		
Absolute change from baseline in standardised T-score at Week 24, LS mean (95% CI)	-8.6 (-10.6, -6.6)	-3.9 (-6.0, -1.8)
Difference versus placebo (95% CI)	-4.7 (-7.6, -1.8)	--
p-value	0.0017	--

CMH – Cochran-Mantel-Haenszel; ITT – intention to treat; n – number of patients in analysis; PROMIS – Patient-Reported Outcomes Measurement Information System.

^a FASENRA 30 mg administered every 4 weeks.

^b Any patients who withdrew from the study without having flared (placebo, n=2; FASENRA, n=2) are included in the analysis as if they had flared.

^c An odds, rate or hazard ratio <1 favours FASENRA.

^d Haematologic relapse is defined as eosinophil count ≥ 1000 cells/ μ L.

^e PROMIS Fatigue Short Form 7a scale: standardised T-score (range 29.4 to 83.2). A reduction in score indicates improvement. A greater improvement in fatigue was observed with FASENRA compared to placebo at Week 4 (difference -3.70, 95% CI -6.5, -0.9, nominal p=0.0092) and this was maintained to Week 24.

Over the 24-week treatment period, 22% of patients treated with FASENRA experienced a HES flare compared to 45% of patients treated with placebo (52% relative reduction).

Oral Corticosteroid Use

Fewer patients experienced HES flares resulting in an increase of OCS with FASENRA (12 [18%]) compared to placebo (28 [42%]) over the 24-week treatment period. The total cumulative OCS dose over the 24-week period was lower in the FASENRA group (LS mean 876 mg) than the placebo group (LS mean 1,221 mg).

Paediatric population

A total of 4 adolescents aged 12 to 17 years were enrolled in the Phase 3 trial, 3 received FASENRA and 1 placebo. One patient in each treatment group had a HES flare during the 24-week treatment period. No new adverse reactions were identified in adolescents treated with FASENRA.

5.2 Pharmacokinetic properties

The pharmacokinetic properties of benralizumab below are based on the population pharmacokinetic analyses from the asthma trials in adults and adolescents. Findings in EGPA and HES were generally similar with those in asthma. The pharmacokinetics of benralizumab were dose-proportional in patients with asthma following subcutaneous administration over a dose range of 2 to 200 mg.

Absorption

Following subcutaneous administration to patients with asthma, the absorption half-life was 3.5 days. Based on population pharmacokinetic analysis, the estimated absolute bioavailability was approximately 59% and there was no clinically relevant difference in relative bioavailability in the administration to the abdomen, thigh or upper arm.

Distribution

Based on population pharmacokinetic analysis, central and peripheral volume of distribution of benralizumab was 3.1 L and 2.5 L respectively for a 70 kg individual.

Metabolism

Benralizumab is a humanized IgG1 monoclonal antibody that is degraded by proteolytic enzymes widely distributed in the body and not restricted to hepatic tissue.

Excretion

From population pharmacokinetic analysis, benralizumab exhibited linear pharmacokinetics and no evidence of target receptor-mediated clearance pathway. The estimated systemic clearance (CL) for benralizumab was at 0.29 L/day. In patients with EGPA and HES, the model estimated systemic CL was approximately 0.22 L/d. Following subcutaneous administration, the elimination half-life was approximately 15.5 days.

Special patient populations

Elderly (≥65 years old)

Based on population pharmacokinetic analysis, age did not affect benralizumab clearance.

Gender, race

A population pharmacokinetics analysis indicated that there was no significant effect of gender and race on benralizumab clearance.

Renal impairment

No formal clinical studies have been conducted to investigate the effect of renal impairment on benralizumab. Based on population pharmacokinetic analysis, mild to moderate renal impairment

(eGFR 30-89 mL/min/1.73m²) did not affect benralizumab clearance. There are limited data available in patients with eGFR less than 30 mL/min/1.73m², however benralizumab is not cleared renally.

Hepatic impairment

No formal clinical studies have been conducted to investigate the effect of hepatic impairment on benralizumab. IgG monoclonal antibodies are not primarily cleared via hepatic pathway; change in hepatic function is not expected to influence benralizumab clearance. Based on population pharmacokinetic analysis, baseline hepatic function biomarkers (ALT, AST and bilirubin) had no clinically relevant effect on benralizumab clearance.

Paediatric use

Based on the population pharmacokinetic analysis, the pharmacokinetics of benralizumab in adolescents with asthma and HES aged 12 to 17 years was consistent with adults. Benralizumab has not been studied in children below 12 years of age (see Section 4.2 Dose and method of administration [*Special patient populations/Paediatric Use*]).

5.3 Preclinical safety data

Genotoxicity

No genotoxicity studies have been conducted. As a monoclonal antibody, benralizumab is not expected to interact with DNA or other chromosomal material.

Carcinogenicity

Long-term animal studies have not been performed to evaluate the carcinogenic potential of benralizumab. Published literature using animal models suggests that IL-5 and eosinophils are part of an early inflammatory reaction at the site of tumorigenesis and can promote tumour rejection. However, other reports indicate that eosinophil infiltration into tumours can promote tumour growth. Therefore, the malignancy risk in humans from an antibody that binds to IL-5R α , such as benralizumab, is unknown.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

FASENRA prefilled syringe and FASENRA PEN contain the excipients histidine, histidine hydrochloride monohydrate, trehalose, polysorbate 20 and water for injections.

FASENRA prefilled syringe and FASENRA PEN do not contain latex, lactose, sucrose, gluten, tartrazine or any other azo dyes.

6.2 Incompatibilities

Incompatibilities were not identified as part of the registration of this medicine.

6.3 Shelf life

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

6.4 Special precautions for storage

Store at 2°C to 8°C (Refrigerate. Do not freeze).

FASENRA may be kept at room temperature up to 25°C for a maximum of 14 days. After removal from the refrigerator, FASENRA must be used within 14 days or discarded.

Store FASENRA in the original package in order to protect from light. Do not freeze. Do not expose to heat.

6.5 Nature and contents of container

FASENRA prefilled syringe**

Each prefilled syringe contains 30 mg benralizumab in 1 mL (30 mg/mL). The prefilled syringe is comprised of a type I glass syringe with a staked 29 gauge 12.7 mm stainless steel needle, rigid needle shield and FluroTec-coated plunger stopper. The prefilled syringe is assembled with a needle guard, a finger flange, and a plunger rod.

FASENRA prefilled syringe is available in a pack containing one single-dose, single use, sterile prefilled syringe.

FASENRA PEN (prefilled pen)

Each FASENRA PEN contains 30 mg benralizumab in 1 mL (30 mg/mL). The prefilled pen is comprised of an autoinjector made from type I glass syringe with a staked 29 gauge 12.7 mm stainless steel needle, rigid needle shield, and Flurotec-coated plunger stopper. The autoinjector consists of the syringe and handheld, mechanical (spring-based) injection device.

FASENRA PEN is available in a pack containing one single-dose, single-use, sterile prefilled pen.

***Not all presentations may be available in Australia*

6.6 Special precautions for disposal


Discard used FASENRA prefilled syringes/FASENRA PEN into a sharps disposal container.

In Australia, any unused medicine should be disposed of by taking to your local pharmacy.

6.7 Physicochemical properties

Benralizumab is a humanised, afucosylated, monoclonal antibody selective for the alpha subunit of the human interleukin-5 receptor (IL-5R α). Benralizumab is of the IgG1, kappa-class produced in Chinese hamster ovary (CHO) cells by recombinant DNA technology. Benralizumab has a molecular weight of approximately 150 kDa.

Figure 4 General structure of benralizumab



CAS number: 1044511-01-4

7 MEDICINE SCHEDULE (POISONS STANDARD)

Prescription only medicine (Schedule 4)

8 SPONSOR

AstraZeneca Pty Ltd
ABN 54 009 682 311
66 Talavera Road
MACQUARIE PARK NSW 2113

Telephone: 1800 805 342

9 DATE OF FIRST APPROVAL

2 April 2018

10 DATE OF REVISION

16 September 2026

Summary table of changes

Section changed	Summary of new information
4.1, 4.2	Extension of indication – Hypereosinophilic Syndrome (HES) as per Study NATRON
4.8	Updated with information related to HES indication
5.1	Updated with clinical data from Study NATRON
5.2	Updated with information related to the HES indication
All	Minor editorial changes throughout the PI to improve clarity

FASENRA and FASENRA PEN are trade marks of the AstraZeneca group of companies.

© AstraZeneca, 2026

VV-RIM-01426683 v9