

# IMFINZI®

---

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using IMFINZI, speak to your doctor or pharmacist.

---

### 1. Why am I using IMFINZI?

IMFINZI contains the active ingredient durvalumab. IMFINZI is used to treat several kinds of cancers.

For more information, see Section [1. Why am I using IMFINZI?](#) in the full CMI.

### 2. What should I know before I use IMFINZI?

Do not use if you have ever had an allergic reaction to IMFINZI or any of the ingredients listed at the end of the CMI.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see Section [2. What should I know before I use IMFINZI?](#) in the full CMI.

---

### 3. What if I am taking other medicines?

Some medicines may interfere with IMFINZI and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### 4. How do I use IMFINZI?

- IMFINZI will be given to you as a liquid infusion into your vein and your doctor will decide how many treatments you need.

More instructions can be found in Section [4. How do I use IMFINZI?](#) in the full CMI.

### 5. What should I know while using IMFINZI?

<b>Things you should do</b>	<ul style="list-style-type: none"><li>• Remind any doctor, dentist or pharmacist you visit that you are using IMFINZI.</li><li>• Keep all of your doctor's appointments so that your progress can be checked.</li></ul>
<b>Things you should not do</b>	<ul style="list-style-type: none"><li>• Do not miss a dose of IMFINZI</li></ul>

<b>Driving or using machines</b>	<ul style="list-style-type: none"> <li>• Be careful before you drive or use any machines or tools until you know how IMFINZI affects you.</li> </ul>
----------------------------------	--

For more information, see Section [5. What should I know while using IMFINZI?](#) in the full CMI.

## 6. Are there any side effects?

Like all medicines, IMFINZI can cause side effects, although not everybody gets them. Your doctor will discuss these with you and will explain the risks and benefits of your treatment.

When you take IMFINZI, you can have some serious side effects.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

# IMFINZI®

**Active ingredient(s):** *durvalumab*

---

## Consumer Medicine Information (CMI)

This leaflet provides important information about using IMFINZI. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using IMFINZI.**

**Where to find information in this leaflet:**

- [1. Why am I using IMFINZI?](#)
- [2. What should I know before I use IMFINZI?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use IMFINZI?](#)
- [5. What should I know while using IMFINZI?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

## 1. Why am I using IMFINZI?

**IMFINZI contains the active ingredient durvalumab.** IMFINZI is a monoclonal antibody, a type of protein. It is a type of immunotherapy and belongs to a group of medicines called immune checkpoint inhibitors. It works with your immune system to destroy cancer cells.

**IMFINZI can be used to treat a type of lung cancer called non-small cell lung cancer (NSCLC).** It may be prescribed to you:

- **in combination with chemotherapy that contains platinum prior to surgery (neoadjuvant treatment) and alone after surgery (adjuvant treatment) if:**
  - your cancer has spread within your lung and is able to be removed by surgery
- **alone if:**
  - your cancer has spread within your lung and cannot be removed by surgery, and
  - you have tried radiotherapy and chemotherapy that contains platinum, and your cancer has shrunk or has not worsened.

**IMFINZI can be used to treat a type of lung cancer called small cell lung cancer (SCLC).** It may be prescribed to you:

- **alone if:**
  - you have limited stage SCLC (LS-SCLC) and
  - your cancer has not been removed by surgery and
  - your cancer has responded or stabilised after initial treatment with radiotherapy and chemotherapy that contains platinum.
- **in combination with chemotherapy if:**
  - you have extensive stage SCLC (ES-SCLC) where your cancer has spread within your lungs (or to other parts of the body) and
  - you have not received previous treatment.

**IMFINZI in combination with chemotherapy can be used to treat a type of cancer called biliary tract cancer (BTC), such as cancer of the bile ducts (cholangiocarcinoma) and gallbladder.** It may be prescribed to you if:

- your cancer has spread within these regions (or to other parts of the body).

**IMFINZI in combination with tremelimumab can be used to treat a type of liver cancer, called unresectable hepatocellular carcinoma (uHCC).** It may be prescribed to you if your cancer:

- cannot be removed by surgery (unresectable), and
- has spread within your liver (or to other parts of the body)

**IMFINZI can be used to treat a type of cancer of the lining of the uterus (endometrial cancer) that has spread beyond the original tumour or come back (recurred).** It may be prescribed to you in combination with chemotherapy, followed by IMFINZI alone.

A test may be used to find out the mismatch repair (MMR) status of your endometrial cancer to help your doctor decide which treatment option may be the best for your cancer.

When IMFINZI is given in combination, it is important that you also read the Consumer Medicine Information (CMI) for the specific anti-cancer medicines you may be receiving. If you have any questions about these medicines, ask your doctor.

Your doctor may have prescribed IMFINZI for another reason.

## **2. What should I know before I use IMFINZI?**

### **Warnings**

#### **Do not use IMFINZI if:**

- you are allergic to durvalumab, or any of the ingredients listed at the end of this leaflet.
- Always check the ingredients to make sure you can use IMFINZI.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

#### **Check with your doctor if you:**

- have an autoimmune disease (a condition where the body attacks its own cells) like Crohn's disease, ulcerative colitis or lupus. If you already have an autoimmune disease, your risk of immune-mediated side effects may be higher with immune checkpoint inhibitor medicines (such as IMFINZI) as they may

cause inflammation in parts of your body. You may also experience more frequent flares of your autoimmune disease, which in the majority of cases are mild

- have had an organ transplant
- have lung or breathing problems
- have liver problems
- have any other medical conditions
- take any medicines for any other condition

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

## **Pregnancy and breastfeeding**

Check with your doctor if you are pregnant or intend to become pregnant.

**You should not be given IMFINZI if you are pregnant.**

It may affect your developing baby if you use it during pregnancy.

If you could become pregnant, you must use adequate birth control while you are being treated with IMFINZI and for at least 3 months after your last dose.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

**You should not be given IMFINZI if you are breast-feeding.**



It is not known if the active ingredient in IMFINZI passes into breast milk, but if it does, there is a possibility that your baby may be affected.

Do not breast-feed if you are being given IMFINZI and for at least 3 months after the last dose.

### **3. What if I am taking other medicines?**

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect IMFINZI.**

### **4. How do I use IMFINZI?**

#### **How much and when to take IMFINZI**

- IMFINZI will be given to you as a liquid infusion into your vein (IV). An infusion takes about 1 hour and will normally be given every 2, 3 or 4 weeks. Depending on your type of cancer, IMFINZI may be given in combination with other anti-cancer medicines.
- Your doctor will decide how many treatments you need.

When IMFINZI is given on the same day as chemotherapy, the IMFINZI infusion goes first, followed by chemotherapy.

When IMFINZI is given on the same day as tremelimumab, the tremelimumab infusion goes first, followed by IMFINZI.

Please refer to the CMI of the other anti-cancer medicines in order to understand the use of these other medicines. If you have questions about these medicines, ask your doctor.

## **If you miss an appointment to be given IMFINZI**

**Call your doctor right away to reschedule your appointment. It is very important that you do not miss a dose of IMFINZI.**

## **5. What should I know while using IMFINZI?**

### **Things you should do**

**Keep all of your doctor's appointments so that your progress can be checked.**

Remind any doctor, dentist or pharmacist you visit that you are using IMFINZI.

### **Driving or using machines**

**Be careful before you drive or use any machines or tools until you know how IMFINZI affects you.**

IMFINZI is not expected to affect your ability to drive a car or operate machinery.

## **Looking after your medicine**

- Store unopened vials under refrigeration at 2°C to 8°C in the original carton to protect from light.
- Do not freeze.
- Do not shake.

Follow the instructions in the carton on how to take care of your medicine properly.

**Keep it where young children cannot reach it.**

## **Getting rid of any unwanted medicine**

If you no longer need to use IMFINZI or it is out of date, take it to any pharmacy for safe disposal.

Do not use IMFINZI after the expiry date.

## **6. Are there any side effects?**

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

IMFINZI can cause your immune system to attack normal organs and tissues in many areas of your body and can affect the way they work and this can cause the side effects.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

## Serious side effects

Serious side effects (signs and symptoms)	What to do
<p><b>Problems with your lungs:</b></p> <ul style="list-style-type: none"> <li>• lung inflammation that may include new or worse cough, shortness of breath, chest pain</li> <li>• chronic obstructive pulmonary disease</li> </ul>	<p><b>Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious</b></p>

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>(CO which causes the narrowing of airways and makes it difficult to breathe)</p> <ul style="list-style-type: none"> <li>• lung infection (pneumonia) or influenza that may include cough or phlegm, fever, chills</li> </ul>	<p><b>side effects.</b></p> <p><b>Your doctor may give you other medicines in order to prevent more severe complications and reduce your symptoms.</b></p> <p><b>Your doctor may withhold the next</b></p>

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>and difficult to breathe</p> <p><b>Problems with your liver:</b></p> <ul style="list-style-type: none"> <li>• inflammation of the liver that may include nausea or vomiting, feeling less hungry, pain on the right side</li> </ul>	<p><b>dose of IMFINZI or stop your treatment with IMFINZI.</b></p>

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>of stomach yellow of skin or white of eyes drowsy dark urine or bleeding or bruising more easily than normal</p> <p><b>Problems with your intestines</b></p>	

Serious side effects (signs and symptoms)	What to do
<ul style="list-style-type: none"> <li>inflammation of the intestines may include diarrhea or more frequent bowel movements than usual, black tarry sticky stools or stools with blood or mucus, severe stomach pain</li> </ul>	<p>n</p>



<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>or tend</p> <p><b>Problems with some of your hormone glands</b></p> <ul style="list-style-type: none"> <li>• inflammation of your hormone gland may include headache that will not go away or unusual</li> </ul>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
head extre tired weigh gain or weigh loss, dizzi or faint feelin more hung or thirs than usua hair loss, feelin cold cons char to your	

Serious side effects (signs and symptoms)	What to do
voice urina more often than usual naus or vom stom area (abd pain char in moo or beha such as decr sex drive incre anxie irrita	

Serious side effects (signs and symptoms)	What to do
<p>or  forge  fast  and  deep  brea  conf  or  a  swee  sme  to  your  brea  a  swee  or  meta  taste  in  your  mou  or  a  diffe  odou</p>	<p>s,</p>

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>to your urine or sweat</p> <p><b>Problems with your kidney</b></p> <ul style="list-style-type: none"> <li>• inflammation of your kidney that may include changes in the amount or color of your urine</li> </ul>	<p>n</p>

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>swelling in your ankles or loss of appetite</p> <p><b>Problems with your urinary tract:</b></p> <ul style="list-style-type: none"> <li>• infection that may include the need to urinate urgently and frequently with burning</li> </ul>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p> pain or  sensation  when urinating  bladder  still feels  full after  urinating  pain above  your pubic  bone  blood in  the urine </p> <p> <b>Problems with your skin</b> </p>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<b>or mouth</b> <ul style="list-style-type: none"> <li>• inflammation of the skin or mouth may include rash, itching, skin blisters or ulcers in the mouth or other mucous membranes.</li> <li>• thrush in</li> </ul>	



<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>the mouth</p> <p><b>Problem with your nose and throat</b></p> <ul style="list-style-type: none"> <li>• nose or throat infection (sinus or tonsillitis)</li> </ul> <p><b>Problem with your heart:</b></p> <ul style="list-style-type: none"> <li>• inflammation of your heart may include</li> </ul>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>chest pain shortness of breath or irregular heartbeat</p> <p><b>Problems with your muscles</b></p> <ul style="list-style-type: none"> <li>• inflammation of your muscles may include muscle weakness, tiredness and/or pain and/</li> </ul>	<p>n</p>

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>or rapid fatigue of the muscles in one or more areas of your body</p> <p><b>Problems with your blood:</b></p> <ul style="list-style-type: none"> <li>• low number of blood platelets - may</li> </ul>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>include bleeding (eg nose or gum bleeding and/or bruising)</p> <p><b>Infusion related reactions</b></p> <ul style="list-style-type: none"> <li>• may include chills or shakiness, itching or rash, flushing, shortness of breath</li> </ul>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>or wheezing, dizziness, fever, feeling like you are passing out, back or neck pain, or facial swelling</p> <p><b>Encephalitis or meningitis</b></p> <ul style="list-style-type: none"> <li>• may include seizure, neck stiffness, headache, fever</li> </ul>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>chills vomiting eye sensitivity to light confusion and sleep</p> <p><b>Guillain-Barré syndrome</b></p> <ul style="list-style-type: none"> <li>may include pain, weakness and paraesthesia in the extremities</li> </ul> <p><b>General body:</b></p> <ul style="list-style-type: none"> <li>an inflammatory</li> </ul>	<p>y</p>

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>respiratory distress that may include low or high body temperature, increased heart rate, increased breathing rate and/or abdominal white blood cell count</p> <p><b>Problems with your</b></p>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<b>immune system</b> <ul style="list-style-type: none"> <li>immune system attacks your red blood cells - signs and symptoms may include unusual weakness and fatigue with increased heart rate and breathing</li> </ul>	



<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>           difficult to breathe            yellow or white patches of the skin or white of the eyes (jaundice)            dark urine and/or stool            or an enlarged spleen         </p> <p> <b>Inflammation of the spinal cord:</b> </p> <ul style="list-style-type: none"> <li>• symptoms may include:</li> </ul>	

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
<p>           include            pain            numb            tingli            or            weak            in            the            arms            or            legs            blad            or            bowe            prob            inclu            need            to            urina            more            frequ            urina            incor            diffic            urina         </p>	<p>e,</p>

**Serious side effects (signs and symptoms)**

and  
cons

**Problems with your eyes:**

- inflammation of your eyes (uveitis) - signs and symptoms include eye redness, eye pain, light sensitivity and/or changes in vision

**What to do**

**Contact a medical eye specialist (ophthalmologist) straight away, or go straight to the Emergency Department at your nearest hospital if you notice these side effects**

<b>Serious side effects (signs and symptoms)</b>	<b>What to do</b>
in vision	<b>effects as you may need urgent medical attention.</b>

## **Immune checkpoint inhibitor class effects**

There have been cases of the following serious side effects reported during treatment with other medicines in this class which may also occur during treatment with IMFINZI:

- decreased ability of the pancreas to make digestive enzymes, which may result in symptoms such as diarrhoea with loose and oily stools, weight loss, metabolic bone disease, and vitamin or mineral deficiencies (known as exocrine pancreatic insufficiency)
- decreased production of blood cells, which may result in symptoms such as weakness, shortness of breath, dizziness, fast or irregular heartbeat, frequent infections, unexplained or easy bruising,

or bleeding from gums or other tissues (known as aplastic anaemia)

- a condition where the immune system makes too many infection-fighting cells which may result in symptoms such as an enlarged liver and/or spleen, skin rash, lymph node enlargement, breathing problems, easy bruising, kidney abnormalities and heart problems (known as haemophagocytic lymphohistiocytosis)
- coeliac disease which may result in symptoms such as stomach pain, diarrhoea and bloating after consuming gluten-containing foods)

**Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.**

**Most common side effects reported in clinical trials:**

<b>Most common side effects</b>	<b>What to do</b>
<b>Report in clinical trials with patients receiving</b>	<b>Speak to your doctor if you have any</b>

Most common side effects	What to do to
<b>IMFINZI alone:</b> <ul style="list-style-type: none"> <li>• cough</li> <li>• diarrhea</li> <li>• stomach pain</li> <li>• skin rash or itching</li> <li>• fever</li> <li>• swelling of the legs</li> <li>• upper respiratory tract infection</li> <li>• underactive thyroid gland that can cause tired</li> </ul>	<b>of these side effects and they worry you.</b>

Most common side effects	What to do
<p>or weight gain (hypothyroidism)</p> <ul style="list-style-type: none"> <li>• overactive thyroid gland which may cause many different symptoms including weight loss, hand tremor, rapid or irregular heart</li> </ul>	
Report in a clinical trial with	Speak to your doctor if

Most common side effects	What do to
<p>patients you receive IMFINZI in combination with chemotherapy follow by IMFINZI alone:</p> <ul style="list-style-type: none"> <li>• nausea</li> <li>• constipation</li> <li>• feeling tired or weak (fatigue)</li> <li>• skin rash and itching</li> <li>• decrease in appetite</li> </ul>	<p>you have any of these side effects, and they worry you.</p>



Most common side effects	What do to
<ul style="list-style-type: none"> <li>• hair loss</li> <li>• diarrhea</li> <li>• inflammation of the nerves causing numbness, weakness, tingling or burning pain of the arms and legs</li> <li>• cough</li> <li>• vomiting</li> <li>• underactive thyroid gland that can cause</li> </ul>	<p>n</p>

Most common side effects	What to do
tired or weight gain (hypothyroidism)	
<b>Report in a clinical trial with patients receiving IMFINZI with etoposide and carboplatin or cisplatin</b> <ul style="list-style-type: none"> <li>• nausea</li> <li>• hair loss</li> <li>• decreased appetite</li> <li>• feeling tired</li> </ul>	<b>Speak to your doctor if you have any of these side effects and they worry you.</b>

<b>Most common side effects</b>	<b>What do to</b>
<p>or weak (fatigue)</p> <ul style="list-style-type: none"> <li>• constipation</li> <li>• vomiting</li> <li>• cough</li> </ul>	
<p><b>Report in clinical trials with patients receiving IMFINZI with gemcitabine and cisplatin</b></p> <ul style="list-style-type: none"> <li>• low number of red blood cells (anaemia)</li> </ul>	<p><b>Speak to your doctor if you have any of these side effects and they worry you.</b></p>

Most common side effects	What do to
<ul style="list-style-type: none"> <li>• low number of white blood cells (neutropenia and leukopenia)</li> <li>• low number of blood platelets (thrombocytopenia)</li> <li>• diarrhea</li> <li>• stomach pain</li> <li>• constipation</li> <li>• nausea</li> <li>• vomiting</li> <li>• feeling tired or weak (fatigue)</li> </ul>	

<b>Most common side effects</b>	<b>What to do to</b>
<ul style="list-style-type: none"> <li>• fever</li> <li>• abnormal liver function tests</li> <li>• decrease in appetite</li> <li>• skin rash and itching</li> </ul>	
<b>Report in clinical trials with patients receiving IMFINZI in combination with tremelimumab:</b> <ul style="list-style-type: none"> <li>• under thyroid</li> </ul>	<b>Speak to your doctor if you have any of these side effects:</b> <b>and they</b>

Most common side effects	What to do
<p>glaucoma that can be caused by tired or weight gain</p> <ul style="list-style-type: none"> <li>• stomach pain</li> <li>• diarrhea</li> <li>• abdominal pain and functional tests</li> <li>• swelling of the legs</li> <li>• fever</li> <li>• abnormal liver function tests</li> <li>• cough</li> </ul>	<p><b>worry you.</b></p>

Most common side effects	What to do
<ul style="list-style-type: none"> <li>• skin rash or itching</li> </ul>	
<p><b>Report in a clinical trial with patient you receive IMFINZI in combination with carboplatin and paclitaxel follow by IMFINZI alone:</b></p> <ul style="list-style-type: none"> <li>• low number of red blood</li> </ul>	<p><b>Speak to your doctor if you have any of these side effects and they worry you.</b></p>

Most common side effects	What do to
<p>cells (anaemia)</p> <ul style="list-style-type: none"> <li>• nausea</li> <li>• feeling tired or weak (fatigue)</li> <li>• low number of white blood cells (neutropenia and leukopenia)</li> <li>• constipation</li> <li>• low number of platelets (thrombocytopenia)</li> <li>• diarrhoea</li> <li>• vomiting</li> </ul>	



Most common side effects	What do to
<ul style="list-style-type: none"> <li>• stomach pain</li> <li>• decreased appetite</li> <li>• swelling of legs</li> <li>• underactive thyroid gland that can cause tired or weight gain (hypothyroidism)</li> <li>• abnormal liver function tests</li> <li>• muscle pain</li> <li>• inflammation of the</li> </ul>	

Most common side effects	What to do
<p>nervousness</p> <p>causing numbness or weakness, tingling or burning pain of the arms and legs</p> <ul style="list-style-type: none"> <li>• cough, productive cough</li> <li>• hair loss</li> <li>• Skin rash or itching</li> </ul>	

Some side effects can only be found when your doctor does blood tests from time to time to check your progress (eg abnormal thyroid gland function test (underactive or overactive), too much calcium in your

blood, not enough sodium in your blood, not enough white blood cells, not enough red blood cells, not enough blood platelets, abnormal liver function tests (eg aspartate aminotransferase (AST) increased; alanine aminotransferase (ALT) increased) or pancreas function tests).

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell. Other less common side effects not listed here may occur in some people.**

## **Reporting side effects**

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of IMFINZI.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

## **7. Product details**

IMFINZI is only available with a doctor's prescription.

### **What IMFINZI contains**

<b>Active ingredient (main ingredient)</b>	durvalumab
--	------------

<b>Other ingredients (inactive ingredients)</b>	histidine histidine hydrochloride monohydrate trehalose dihydrate polysorbate 80 water for injections
---	--

**Do not take IMFINZI if you are allergic to any of these ingredients.**

## **What IMFINZI looks like**

IMFINZI concentrated solution for infusion is a clear to opalescent, colourless to slightly yellow liquid in a glass vial.

Australian Registration Number(s):

IMFINZI, 500 mg (500 mg/10mL) in 10 mL vial for intravenous infusion - AUST R 283216

IMFINZI, 120 mg (120 mg/2.4mL) in 10 mL vial for intravenous infusion - AUST R 283215

## **Who distributes IMFINZI**

AstraZeneca Pty Ltd  
ABN 54 009 682 311  
66 Talavera Road  
MACQUARIE PARK NSW 2113

Telephone: 1800 805 342

This leaflet was prepared 12 August 2025

IMFINZI is a trade mark of the AstraZeneca group of companies.

© AstraZeneca 2025

VV-RIM-01430609 v23