

# Lioresal<sup>®</sup> tablets

---

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

---

### 1. Why am I using Lioresal?

Lioresal contains the active ingredient baclofen. Lioresal belongs to a group of medicines called muscle relaxants.

Lioresal is used to reduce the stiffness or spasms in your muscles to help make you more mobile and able to manage your daily activities.

For more information, see Section [1. Why am I using Lioresal?](#) in the full CMI.

### 2. What should I know before I use Lioresal?

Do not use if you have ever had an allergic reaction to Lioresal or any of the ingredients listed at the end of the CMI.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see Section [2. What should I know before I use Lioresal?](#) in the full CMI.

### **3. What if I am taking other medicines?**

Some medicines may interfere with Lioresal and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### **4. How do I use Lioresal?**

- Your doctor will tell you exactly how many tablets of Lioresal to take.
- Do not exceed the recommended dose prescribed by your doctor.
- Treatment is usually started in hospital with small doses of Lioresal. The dose is then gradually increased to an amount that works best for you

More instructions can be found in Section [4. How do I use Lioresal?](#) in the full CMI.

## 5. What should I know while using Lioresal?

<b>Things you should do</b>	<ul style="list-style-type: none"><li>● Remind any doctor, dentist or pharmacist you visit that you are using Lioresal.</li><li>● Keep all doctor's appointments, so that your progress can be checked.</li></ul>
<b>Things you should not do</b>	<ul style="list-style-type: none"><li>● Do not stop using this medicine suddenly, unless your doctor tells you to</li></ul>
<b>Driving or using machines</b>	<ul style="list-style-type: none"><li>● Be careful before you drive or use any machines or tools until you know how Lioresal affects you.</li><li>● Lioresal may cause sleepiness in some people, especially when you start taking it</li></ul>
<b>Drinking alcohol</b>	<ul style="list-style-type: none"><li>● Tell your doctor if you drink alcohol.</li></ul>
<b>Looking after your medicine</b>	<ul style="list-style-type: none"><li>● Store the tablets in a cool dry place</li></ul>

- Keep out of reach of children

For more information, see Section [5. What should I know while using Lioresal?](#) in the full CMI.

## **6. Are there any side effects?**

Common side effects include daytime sleepiness, lack of energy, tiredness, dizziness or light-headedness, vertigo, confusion, headache, difficulty sleeping or nightmares, nausea, retching or vomiting, constipation, stomach cramps or diarrhoea, loss of appetite, stuffy nose, dry mouth, change in sense of taste, misuse, abuse and dependence, numbness, muscle weakness or spasms, swelling of ankles, blurred vision, ringing in the ears, frequent urination, excessive sweating, weight gain, impotence or inability to ejaculate, increased blood sugar, low body temperature. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

# Lioresal tablets

Active ingredient(s): *Baclofen*

---

## Consumer Medicine Information (CMI)

This leaflet provides important information about using Lioresal tablets. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Lioresal.**

**Where to find information in this leaflet:**

- [1. Why am I using Lioresal?](#)
- [2. What should I know before I use Lioresal?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Lioresal?](#)
- [5. What should I know while using Lioresal?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

## 1. Why am I using Lioresal?

**Lioresal contains the active ingredient baclofen.** Lioresal belongs to a group of medicines called muscle relaxants.

**Lioresal is used to reduce the stiffness or spasms in your muscles to help make you more mobile and able to manage your daily activities.** These spasms

happen in various illnesses such as multiple sclerosis and diseases or injuries of the spinal cord.

## **2. What should I know before I use Lioresal?**

### **Warnings**

#### **Do not use Lioresal if:**

- you are allergic to baclofen, or any of the ingredients listed at the end of this leaflet.

Always check the ingredients to make sure you can use this medicine.

#### **Check with your doctor if you:**

- have any other medical conditions including mental illness, Parkinson's disease, seizures, stiffness and restriction of movement in a group of muscles, stomach ulcers, stroke or blood circulation problems, diabetes, a blood disorder known as porphyria, high blood pressure, difficulty urinating
- have problems with your heart, kidney, liver or lungs
- have a history of alcoholism, drink large amounts of alcohol, or have a history of drug abuse or dependence
- have had thoughts of self-harm or and suicidal thoughts
- take any medicines for any other condition

- have any questions about why this medicine has been prescribed for you.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

## **Pregnancy and breastfeeding**

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

## **3. What if I am taking other medicines?**

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

**Some medicines may interfere with Lioresal and affect how it works. These include:**

- Any medicine that makes you sleepy such as medicines used to help you sleep or calm you, pain relievers, and medicines for cold or allergies
- medicines used to treat mood disorders, including tricyclic antidepressants, lithium and monoamine oxidase inhibitors (MAOIs)
- medicines used to treat diabetes

- medicines that lower blood pressure including those used to treat high blood pressure
- medicines used to treat Parkinson's disease, including selegiline and levodopa and carbidopa.

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Lioresal.**

## **4. How do I use Lioresal?**

### **How much to take**

- Your doctor will tell you exactly how many tablets of Lioresal to take.
- Do not exceed the recommended dose prescribed by your doctor.
- Treatment is usually started in hospital with small doses of Lioresal. The dose is then gradually increased to an amount that works best for you
- If you are under the age of 16 or over 65, or you have kidney disease, your doctor may start you on a lower dose and increase it slowly to prevent unwanted side effects
- Follow the instructions provided and use Lioresal until your doctor tells you to stop.

### **When to take Lioresal**

- Lioresal is usually taken in 3 divided doses throughout the day



- Your doctor may tell you to take it more or less often, depending on your situation.

## **How to take Lioresal**

- Swallow the tablets whole during meals with a glass of water or other liquid.

## **If you forget to take Lioresal**

**If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.**

**Otherwise, take it as soon as you remember and then go back to taking it as you would normally.**

**Do not take a double dose to make up for the dose you missed.**

## **If you take too much Lioresal**

If you think that you have taken too much Lioresal, you may need urgent medical attention.

### **You should immediately:**

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

**You should do this even if there are no signs of discomfort or poisoning.**

## **Symptoms of overdose include:**

- Main symptoms such as drowsiness, breathing difficulties, and being unconscious (coma)
- Other symptoms such as feeling confused, hallucinations, agitation, convulsions, blurred vision, unusual low muscle tone, sudden and involuntary muscle spasm, poor or absent reflexes, high or low blood pressure, slow, fast or irregular heartbeat, low body temperature, nausea, vomiting, diarrhoea, excessive salivation, trouble breathing during sleep (sleep apnoea), severe muscle pain with fever and dark urine (rhabdomyolysis), and ringing in the ears
- If you have kidney disease and have accidentally taken more tablets than your doctor has prescribed, you may feel drowsy, confused, have hallucinations.

## **5. What should I know while using Lioresal?**

### **Things you should do**

**Keep all doctor's appointments, so that your progress can be checked.**

### **Call your doctor straight away if you:**

- Become pregnant or think you might be pregnant.
- Remind any doctor, dentist or pharmacist you visit that you are using Lioresal.

**Call your doctor straight away if you have symptoms such as increased sleepiness, new onset of drowsiness, confusion or muscle jerks.**

These may be signs of a condition known as encephalopathy and your doctor may decide to stop treatment with Lioresal tablets.

**Call your doctor straight away or go to a hospital if you have thoughts of harming or killing yourself.**

Some people treated with Lioresal have had suicidal thoughts. Ask a relative or close friend to tell you if they are worried about any changes in your behaviour.

## **Things you should not do**

- **Do not stop using this medicine suddenly unless your doctor tells you to.**

Stopping Lioresal suddenly may make your condition worse. You may experience the following: nervousness, feeling confused, hallucinations, abnormal thinking or behaviour, convulsions, uncontrollable twitching, jerking or writhing movements, fast heartbeat, high body temperature, pain in muscles, fever, dark urine. The excessive stiffness (spasms) in your muscles may also get worse.

- Do not use this medicine to treat any other problems unless your doctor tells you to
- Do not give this medicine to anyone else. Even if they have the same condition as you.

## **Driving or using machines**

**Be careful before you drive or use any machines or tools until you know how Lioresal affects you.**

Lioresal may cause sleepiness and decreased alertness in some people, especially when you start taking it.

## **Drinking alcohol**

**Tell your doctor if you drink alcohol.**

Alcohol may make you feel more sleepy when taking Lioresal.

## **Looking after your medicine**

- Keep your tablets in the original container until it is time to take them.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

**Keep it where young children cannot reach it.**

## **When to discard your medicine**

**If your doctor tells you to stop taking Lioresal or the expiry date on the medicine has passed, ask your pharmacist what to do with any tablets you have left over.**

## **Getting rid of any unwanted medicine**

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

## **6. Are there any side effects?**

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

Side effects happen mainly at the start of treatment with Lioresal or if the dose is too high or increased too quickly. Your doctor will adjust your dose if necessary.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

## Less serious side effects

Less serious side effects	What to do
<p><b>Brain and nerves:</b></p> <ul style="list-style-type: none"><li>• daytime sleepiness or drowsiness</li><li>• lack of energy, tiredness</li><li>• dizziness or light-headedness</li><li>• spinning sensation (vertigo)</li><li>• mental confusion</li><li>• headache</li><li>• difficulty sleeping or nightmares</li><li>• change in sense of taste</li><li>• blurred or double vision</li><li>• ringing in the ears</li></ul> <p><b>Airways:</b></p> <ul style="list-style-type: none"><li>• stuffy or blocked nose</li><li>• dry mouth</li></ul> <p><b>Stomach problems:</b></p> <ul style="list-style-type: none"><li>• nausea (feeling sick), retching or vomiting</li><li>• constipation, stomach cramps or diarrhoea</li></ul>	<p><b>Speak to your doctor if you have any of these less serious side effects and they worry you.</b></p> <p><b>If these side effects become severe, please tell your doctor, pharmacist or healthcare provider.</b></p>

<b>Less serious side effects</b>	<b>What to do</b>
<ul style="list-style-type: none"> <li>● loss of appetite</li> <li><b>General problems:</b></li> <li>● numbness or tingling in hands and feet</li> <li>● muscle weakness, spasms or pain</li> <li>● problems with coordination and balance</li> <li>● difficulty in speaking</li> <li>● swelling of ankles due to fluid build-up</li> <li>● frequent urination or bed wetting</li> <li>● excessive sweating</li> <li>● weight gain</li> <li>● impotence or not able to ejaculate</li> <li>● increased blood sugar</li> <li>● feeling cold</li> <li>● hair loss.</li> </ul>	

## Serious side effects

Serious side effects	What to do
<p data-bbox="204 432 549 533"><b>Skin or muscle problems:</b></p> <ul data-bbox="204 568 753 1205" style="list-style-type: none"><li data-bbox="204 568 753 965">● signs of allergy such as rash, itching or hives on the skin; swelling of the face, lips, tongue or other parts of the body; shortness of breath or wheezing</li><li data-bbox="204 981 753 1205">● uncontrollable muscle spasms affecting the eyes, head, neck or body</li></ul> <p data-bbox="204 1227 584 1279"><b>Chest problems:</b></p> <ul data-bbox="204 1308 778 1554" style="list-style-type: none"><li data-bbox="204 1308 778 1359">● slow or difficult breathing</li><li data-bbox="204 1375 778 1487">● irregular heart beat (fast or slow)</li><li data-bbox="204 1503 778 1554">● chest pain</li></ul> <p data-bbox="204 1576 743 1628"><b>Neurological problems:</b></p> <ul data-bbox="204 1657 762 1944" style="list-style-type: none"><li data-bbox="204 1657 762 1709">● fainting or seizures (fits)</li><li data-bbox="204 1724 762 1944">● depression or other severe mood, behavioural or mental changes</li></ul>	<p data-bbox="810 432 1378 824"><b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b></p>



<b>Serious side effects</b>	<b>What to do</b>
<ul style="list-style-type: none"> <li>● hallucinations (hearing or seeing things that are not there)</li> <li>● thoughts of suicide</li> <li>● anxiety, confusion, paranoia, and other mental problems following sudden discontinuation of the medicine (drug withdrawal syndrome)</li> <li>● feeling more tired than normal, decreased levels of awareness with uncontrolled muscle jerks.</li> </ul> <p><b>Other problems:</b></p> <ul style="list-style-type: none"> <li>● being unable to urinate or pain when urinating; blood in the urine</li> </ul>	

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

## Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

## 7. Product details

This medicine is only available with a doctor's prescription.

### What Lioresal contains

<b>Active ingredient (main ingredient)</b>	baclofen
<b>Other ingredients (inactive ingredients)</b>	Colloidal anhydrous silica Microcrystalline cellulose Magnesium stearate Povidone Wheat starch

<b>Potential allergens</b>	<p>Tell your doctor if you have an intolerance to gluten.</p> <p>This medicine contains wheat starch.</p> <p>Lioresal tablets do not contain lactose, sucrose, tartrazine or any other azo dyes.</p>
----------------------------	--

**Do not take this medicine if you are allergic to any of these ingredients.**

## **What Lioresal looks like**

Lioresal 10 mg: white, round, flat tablets marked with KJ and a score line on one side and CG on the other side. Bottles of 100 tablets; AUST R 11038.

Lioresal 25 mg: white, round, flat tablets marked with UR and a score line on one side and CG on the other side. Blister packs of 100 tablets; AUST R 68365.

## **Who distributes Lioresal**

Lioresal is supplied in Australia by:

NOVARTIS Pharmaceuticals Australia Pty Limited

ABN 18 004 244 160

54 Waterloo Road

Macquarie Park NSW 2113

Telephone 1800 671 203

[www.novartis.com.au](http://www.novartis.com.au)

® = Registered trademark

This leaflet was prepared in September 2024.

Internal document code

(Irst240924c is based on PI Irst240924i).