

# AUSTRALIAN PRODUCT INFORMATION – MERSYNDOL FORTE (PARACETAMOL, CODEINE PHOSPHATE HEMIHYDRATE, DOXYLAMINE SUCCINATE)

## WARNINGS

### Limitations of use

Because of the risks associated with the use of opioids, Mersyndol Forte should only be used in patients for whom other treatment options, including non-opioid analgesics, are ineffective, not tolerated or otherwise inadequate to provide appropriate management of pain (see section 4.4 Special Warnings and Precautions for Use).

### Hazardous and harmful use

Mersyndol Forte poses risks of hazardous and harmful use which can lead to overdose and death. Assess the patient's risk of hazardous and harmful use before prescribing and monitor the patient regularly during treatment (see section 4.4. Special Warnings and Precautions for Use).

### Life threatening respiratory depression

Serious, life-threatening or fatal respiratory depression may occur with the use of Mersyndol Forte. Be aware of situations which increase the risk of respiratory depression, modify dosing in patients at risk and monitor patients closely, especially on initiation or following a dose increase (see section 4.4 Special Warnings and Precautions for Use).

### Concomitant use of benzodiazepines and other central nervous system (CNS) depressants, including alcohol

Concomitant use of opioids with benzodiazepines, gabapentinoids, antihistamines, tricyclic antidepressants, antipsychotics, cannabis or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Limit dosages and durations to the minimum required; and monitor patients for signs and symptoms of respiratory depression and sedation. Caution patients not to drink alcohol while taking Mersyndol Forte.

## 1 NAME OF THE MEDICINE

Paracetamol, codeine phosphate hemihydrate and doxylamine succinate

## **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains paracetamol 450 mg, codeine phosphate hemihydrate 30 mg, doxylamine succinate 5 mg.

For the full list of excipients, see Section [6.1 LIST OF EXCIPIENTS](#).

## **3 PHARMACEUTICAL FORM**

White tablets

## **4 CLINICAL PARTICULARS**

### **4.1 THERAPEUTIC INDICATIONS**

Mersyndol Forte is indicated for the short-term management of severe pain for which other treatment options have failed, are contraindicated, not tolerated or are otherwise inappropriate to provide sufficient management of pain.

### **4.2 DOSE AND METHOD OF ADMINISTRATION**

Adults and children 12 years of age and older: One or two tablets every 4 to 6 hours as needed for relief. Do not exceed 8 tablets in a 24 - hour period. Mersyndol Forte is contraindicated for use in patients who are:

- younger than 12 years.
- aged between 12 – 18 years in whom respiratory function might be compromised, including post tonsillectomy and/or adenoidectomy for obstructive sleep apnoea. (See Section [4.3 CONTRAINDICATIONS](#) and Section [4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – Paediatric use](#))

### **4.3 CONTRAINDICATIONS**

MERSYNDOL FORTE should not be given to patients with a known hypersensitivity to paracetamol, codeine, doxylamine succinate or any of the excipients listed under Section [6.1 LIST OF EXCIPIENTS](#).

It must not be used in patients with known glucose-6-phosphate-dehydrogenase deficiency, severe respiratory disease, acute respiratory disease and respiratory depression/insufficiency, for example acute asthma, acute exacerbations of chronic obstructive pulmonary disease since codeine may exacerbate the condition.

Concomitant treatment with Monoamine inhibitors (MAOIs) or treatment within 14 days of stopping MAOIs is contraindicated.

Paracetamol should not be used in patients with a history of intolerance to the drug.

Paracetamol should not be used in patients with severe hepatocellular insufficiency.

Due to codeine's structural similarity to morphine and oxycodone, patients experiencing systemic allergy (generalised rash, shortness of breath) to these drugs should not receive codeine.

Codeine is contraindicated in patients with diarrhoea caused by poisoning, until the toxic substance has been eliminated from the gastrointestinal tract, or diarrhoea associated with pseudomembranous colitis caused by antibiotic administration since codeine may slow the elimination of the toxic material or antibiotic.

Codeine is contraindicated in the event of impending childbirth or in case of risk of premature birth (see Section 4.6 FERTILITY, PREGNANCY AND LACTATION – Use in pregnancy).

Mersyndol Forte is contraindicated during breast-feeding (see Section 4.6 FERTILITY, PREGNANCY AND LACTATION – Use in lactation).

Mersyndol Forte is contraindicated for use in patients who are:

- younger than 12 years (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – Paediatric use).
- Aged between 12 – 18 years in whom respiratory function might be compromised, including post tonsillectomy and/or adenoidectomy for obstructive sleep apnoea, due to an increased risk of developing serious life-threatening adverse reactions (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – Paediatric use).

Mersyndol Forte is contraindicated for use in patients who are CYP 2D6 ultra-rapid metabolisers (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – CYP2D6 Metabolism).

#### **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE**

Hepatotoxicity may occur with paracetamol even at therapeutic doses, after short treatment duration and in patients without pre-existing liver dysfunction. In view of the increased risk of hepatotoxicity, the benefit should be weighed against the risk when administering Mersyndol Forte to patients with viral hepatitis or pre-existing hepatic disease. In such patients, hepatic function determinations may be required at periodic intervals during high dose or long-term therapy.

It has been reported that paracetamol may produce symptoms of acute toxicity in adults following the ingestion of more than 15g.

Hepatotoxicity may develop after the ingestion of a single dose of 10 to 15g (200 to 250 mg/kg) and a dose of more than 25g is potentially fatal. Patients may be asymptomatic for several days following ingestion of large doses of paracetamol and laboratory evidence of hepatotoxicity may be delayed for up to one week. Non-fatal hepatic damage is usually reversible. There have been reports of kidney damage, disturbances in clotting mechanisms, metabolic acidosis, hypoglycaemia, agranulocytosis, thrombocytopenia, methaemoglobinaemia and myocardial necrosis.

To avoid the risk of overdose, check that paracetamol is absent from the composition of other medicinal products taken concomitantly.

Caution is advised in patients with underlying sensitivity to aspirin and/or to non-steroidal anti-inflammatory drugs (NSAIDs).

Severe cutaneous adverse reactions (SCARs): Life-threatening cutaneous reactions Stevens Johnson Syndrome (SJS), and Toxic Epidermal Necrolysis (TEN) have been reported with the use of paracetamol. If symptoms or signs of SJS and TEN (e.g. progressive skin rash often with blisters or mucosal lesions) occur, patients should stop immediately paracetamol treatment and seek medical advice.

**Paracetamol should be used with caution in patients with:**

- chronic alcohol use including recent cessation of alcohol intake
- low glutathione reserves
- Gilbert's syndrome

Codeine should be used with caution in patients with CNS depression or decreased respiratory reserve e.g. in emphysema, kyphoscoliosis, hypoxia, hypercapnia or even severe obesity or cor pulmonale, or chronic obstructive pulmonary disease.

Codeine should be administered with caution in patients with impaired cardiac, hepatic or renal function, hypotension, benign prostatic hyperplasia, urethral stenosis, chronic colitis ulcerative, gall bladder conditions, multiple sclerosis, hypothyroidism, adrenocortical insufficiency (e.g. Addison's disease), shock, myxedema, acute alcohol intoxication or delirium tremens since codeine may exacerbate the symptoms or increase the risk of respiratory and/or CNS depression.

Codeine should be administered with great caution in patients with head injury, brain tumour or increased intracranial pressure since codeine may increase the risk of respiratory depression and further elevate intracranial pressure. In addition codeine can produce side effects such as:

- confusion
- miosis
- vomiting

These are important signs in following the clinical course of patients with head injuries.

Extensive use of analgesics to relieve headaches or migraines, especially at high doses, may induce headaches that must not be treated with increased doses of the drug. In such cases the analgesic should not continue to be taken without medical advice.

Monitoring after prolonged use should include blood count, liver function and renal function.

Codeine should only be used with careful risk-benefit assessment and great caution in case of:

- Opioid dependence
- Chronic constipation
- Conditions with elevated intracranial pressure and head trauma. Codeine can increase the pressure of cerebrospinal fluid and may increase the respiratory depressant effect.

Like other narcotics, it causes adverse reactions that can obscure the clinical course of patients with head injury.

- Impaired consciousness
- Compromised respiratory function (due to emphysema, kyphoscoliosis, severe obesity) and chronic obstructive airway disease

Paracetamol should not be used in patients with active alcoholism as chronic excessive alcohol ingestion predisposes patients to paracetamol hepatotoxicity.

Codeine should be administered with caution in patients with acute abdominal conditions since codeine may obscure the diagnosis or the course of the disease. Codeine should be administered with caution in patients with severe inflammatory bowel disease (risk of toxic megacolon may be increased, especially with repeated dosing). Mersyndol Forte should also be used with caution in patients who have had recent gastrointestinal tract surgery.

Patients who have had a cholecystectomy should be treated with caution. The contraction of the sphincter of Oddi can cause symptoms resembling those of myocardial infarction or intensify the symptoms in patients with pancreatitis.

Codeine should be administered with caution in patients with a history of convulsive disorders (convulsions may be induced or exacerbated by codeine).

Codeine should be administered with caution in patients with prostatic hypertrophy, urethral structure or recent urinary tract surgery since codeine may cause urinary retention.

Codeine should be used with caution in elderly or debilitated patients because of the danger of respiratory or cardiac depression.

Patients with known analgesic intolerance or known bronchial asthma must only use Mersyndol Forte after having consulted a physician (hypersensitivity reactions including bronchospasm are possible).

### **Hazardous and harmful use**

Mersyndol Forte contains the opioid codeine and is a potential drug of abuse, misuse and addiction. Addiction can occur in patients appropriately prescribed Mersyndol Forte at recommended doses.

The risk of addiction is increased in patients with a personal or family history of substance abuse (including alcohol and prescription and illicit drugs) or mental illness. The risk also increases the longer the drug is used and with higher doses. Patients should be assessed for their risks for opioid abuse or addiction prior to being prescribed Mersyndol Forte.

There have been reports of drug abuse with codeine, including cases in children and adolescents. Caution is particularly recommended for use in children, adolescents, young adults and in patients with a history of drug and/or alcohol abuse. See Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – Paediatric use.

All patients receiving opioids should be routinely monitored for signs of misuse and abuse. Opioids are sought by people with addiction and may be subject to diversion. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and

advising the patient on the safe storage and proper disposal of any unused drug (see section 6.4 Special precautions for storage and section 6.6 Special precautions for disposal). Caution patients that abuse of oral or transdermal forms of opioids by parenteral administration can result in serious adverse events, which may be fatal.

Patients should be advised not to share Mersyndol Forte with anyone else.

### **Respiratory depression**

Serious, life-threatening or fatal respiratory depression can occur with the use of opioids even when used as recommended. It can occur at any time during the use of Mersyndol Forte but the risk is greatest during initiation of therapy or following an increase in dose. Patients should be monitored closely for respiratory depression at these times.

The risk of life-threatening respiratory depression is also higher in elderly, frail, or debilitated patients, in patients with hepatic and renal impairment (see Use in hepatic impairment and Use in renal impairment) and in patients with existing impairment of respiratory function (e.g. chronic obstructive pulmonary disease; asthma). Opioids should be used with caution and with close monitoring in these patients. The use of opioids is contraindicated in patients with severe respiratory disease, acute respiratory disease and respiratory depression (see section 4.3 Contraindications).

The risk of respiratory depression is greater with the use of high doses of opioids, especially high potency and modified release formulations, and in opioid naïve patients. Initiation of opioid treatment should be at the lower end of the dosage recommendations with careful titration of doses to achieve effective pain relief. Careful calculation of equianalgesic doses is required when changing opioids or switching from immediate release to modified release formulations, together with consideration of pharmacological differences between opioids. Consider starting the new opioid at a reduced dose to account for individual variation in response.

### **Risks from concomitant use of benzodiazepines or other CNS depressants, including alcohol**

Concomitant use of opioids and benzodiazepines or other CNS depressants, including alcohol, may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing of Mersyndol Forte with CNS depressant medicines, such as other opioid analgesics, benzodiazepines, gabapentinoids, cannabis, sedatives, hypnotics, tricyclic antidepressants, antipsychotics, antihistamines, centrally-active anti-emetics and other CNS depressants, should be reserved for patients for whom other treatment options are not possible. If a decision is made to prescribe Mersyndol Forte concomitantly with any of the medicines, the lowest effective dose should be used, and the duration of treatment should be as short as possible. Patients should be followed closely for signs and symptoms of respiratory depression and sedation. Patients and their caregivers should be made aware of these symptoms. Patients and their caregivers should also be informed of the potential harms of consuming alcohol while taking Mersyndol Forte.

### **Use of opioids in chronic (long-term) non-cancer pain (CNCP)**

Opioid analgesics have an established role in the treatment of acute pain, cancer pain and palliative and end-of-life care. Current evidence does not generally support opioid analgesics

in improving pain and function for most patients with chronic non-cancer pain. The development of tolerance and physical dependence and risks of adverse effects, including hazardous and harmful use, increase with the length of time a patient takes an opioid. The use of opioids for long-term treatment of CNCP is not recommended.

The use of an opioid to treat CNCP should only be considered after maximised non-pharmacological and non-opioid treatments have been tried and found ineffective, not tolerated or otherwise inadequate to provide sufficient management of pain. Opioids should only be prescribed as a component of comprehensive multidisciplinary and multimodal pain management.

Opioid therapy for CNCP should be initiated as a trial in accordance with clinical guidelines and after a comprehensive biopsychosocial assessment has established a cause for the pain and the appropriateness of opioid therapy for the patient (see Hazardous and harmful use, above). The expected outcome of therapy (pain reduction rather than complete abolition of pain, improved function and quality of life) should be discussed with the patient before commencing opioid treatment, with agreement to discontinue treatment if these objectives are not met.

Owing to the varied response to opioids between individuals, it is recommended that all patients be started at the lowest appropriate dose and titrated to achieve an adequate level of analgesia and functional improvement with minimum adverse reactions. Immediate-release products should not be used to treat chronic pain, but may be used for a short period in opioid-naïve patients to develop a level of tolerance before switching to a modified-release formulation. Careful and regular assessment and monitoring is required to establish the clinical need for ongoing treatment. Discontinue opioid therapy if there is no improvement of pain and/or function during the trial period or if there is any evidence of misuse or abuse. Treatment should only continue if the trial has demonstrated that the pain is opioid responsive and there has been functional improvement. The patient's condition should be reviewed regularly and the dose tapered off slowly if opioid treatment is no longer appropriate (see Ceasing Opioids).

### **Tolerance, dependence and withdrawal**

Neuroadaptation of the opioid receptors to repeated administration of opioids can produce tolerance and physical dependence. Tolerance is the need for increasing doses to maintain analgesia. Tolerance may occur to both the desired and undesired effects of the opioid.

Physical dependence, which can occur after several days to weeks of continued opioid usage, results in withdrawal symptoms if the opioid is ceased abruptly or the dose is significantly reduced. Withdrawal symptoms can also occur following the administration of an opioid antagonist (e.g. naloxone) or partial agonist (e.g. buprenorphine). Withdrawal can result in some or all of the following symptoms: dysphoria, restlessness/agitation, lacrimation, rhinorrhoea, yawning, sweating, chills, myalgia, mydriasis, irritability, anxiety, increasing pain, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhoea, increased blood pressure, increased respiratory rate and increased heart rate.

When discontinuing Mersyndol Forte in a person who may be physically-dependent, the drug should not be ceased abruptly but withdrawn by tapering the dose gradually (see Ceasing opioids and section 4.2 Dose and Method of Administration).

## **Accidental ingestion/exposure**

Accidental ingestion or exposure of Mersyndol Forte, especially by children, can result in a fatal overdose of codeine. Patients and their caregivers should be given information on safe storage and disposal of unused Mersyndol Forte (see section 6.4 Special precautions for storage and section 6.6 Special precautions for disposal).

## **Hyperalgesia**

Hyperalgesia may occur with the use of opioids, particularly at high doses. Hyperalgesia may manifest as an unexplained increase in pain, increased levels of pain with increasing opioid dosages or diffuse sensitivity not associated with the original pain. Hyperalgesia should not be confused with tolerance (see Tolerance, dependence and withdrawal). If opioid induced hyperalgesia is suspected, the dose should be reduced and tapered off if possible. A change to a different opioid may be required.

## **Opioid-Induced Hyperalgesia or Allodynia**

Opioid-Induced Hyperalgesia (OIH) occurs when an opioid analgesic paradoxically causes an increase in pain (hyperalgesia), or an increase in sensitivity to pain (allodynia). This condition differs from tolerance, which is the need for increasing doses of opioids to maintain a defined effect.

Symptoms of OIH include increased levels of pain upon opioid dosage increase, decreased levels of pain upon opioid dosage decrease, or pain from ordinarily non-painful stimuli (allodynia). The pain experienced may be at the same location of the underlying pain or can be more generalised or widespread in nature. These symptoms may suggest the occurrence of OIH only if there is no evidence of underlying disease progression, opioid tolerance, opioid withdrawal, or addictive behaviour.

If a patient is suspected to be experiencing OIH, carefully consider appropriately decreasing the dose of the current opioid analgesic, or opioid rotation (safety switching the patient to a different opioid moiety).

## **Ceasing opioids**

Abrupt discontinuation or rapid decreasing of the dose in a person physically dependent on an opioid may result in serious withdrawal symptoms and uncontrolled pain (see Tolerance, dependence and withdrawal). Such symptoms may lead the patient to seek other sources of licit or illicit opioids. Opioids should not be ceased abruptly in a patient who is physically dependent but withdrawn by tapering the dose slowly. Factors to take into account when deciding how to discontinue or decrease therapy include the dose and duration of the opioid the patient has been taking, the type of pain being treated and the physical and psychological attributes of the patient. A multimodal approach to pain management should be in place before initiating an opioid analgesic taper. During tapering, patients require regular review and support to manage any increase in pain, psychological distress and withdrawal symptoms.

There are no standard tapering schedules suitable for all patients and an individualised plan is necessary. In general, tapering should involve a dose reduction of no more than 10 percent to 25 percent every 2 to 4 weeks (see section 4.2 Dose and Method of Administration). If the



patient is experiencing increased pain or serious withdrawal symptoms, it may be necessary to go back to the previous dose until stable before proceeding with a more gradual taper.

When ceasing opioids in a patient who has a suspected opioid use disorder, the need for medication assisted treatment and/or referral to a specialist should be considered.

Doxylamine must be administered with caution in patients with:

- urinary retention
- susceptibility to angle closure glaucoma
- benign prostatic hyperplasia

### **CYP2D6 Metabolism**

Mersyndol Forte is contraindicated for use in patients who are CYP2D6 ultra-rapid metabolisers.

Codeine is metabolised by the liver enzyme CYP2D6 into morphine, its active metabolite. If a patient has a deficiency or is completely lacking this enzyme an adequate analgesic effect will not be obtained. However, if the patient is an extensive or ultra-rapid metaboliser there is an increased risk of developing side effects of opioid toxicity even at commonly prescribed doses. These patients convert codeine into morphine rapidly resulting in higher than expected serum morphine levels. General symptoms of opioid toxicity include confusion, somnolence, shallow breathing, small pupils, nausea, vomiting, constipation and lack of appetite. In severe cases this may include symptoms of circulatory and respiratory depression, which may be life-threatening and very rarely fatal. Children are particularly susceptible due to their immature airway anatomy. Deaths have been reported in children with rapid metabolism who were given codeine for analgesia post adenotonsillectomy. Morphine can also be ingested by infants through breast milk, causing risk or respiratory depression to infants of rapid metaboliser mothers who take codeine.

The prevalence of codeine ultra-rapid metabolism by CYP2D6 in children is not known, but is assumed to be similar to that reported in adults. The prevalence of ultra-rapid metabolisers is estimated to be 1% in those of Chinese, Japanese and Hispanic decent, 3% in African Americans and 1%-10% in Caucasians. The highest prevalence (16%-28%) occurs in North African, Ethiopian and Arab populations.

(See Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE - Paediatric Use and Section 4.6 FERTILITY, PREGNANCY AND LACTATION - Use in Lactation.)

This medication may be dangerous when used in large amounts or for long periods.

### **Use in hepatic impairment**

Mersyndol Forte should be administered with caution to patients with hepatic dysfunction, viral hepatitis, and to patients taking other drugs which affect the liver.

### **Use in renal impairment**

Mersyndol Forte should be administered with caution to patients with renal dysfunction.

## Use in the elderly

Elderly people may be more sensitive to the effects of this medicinal product. The elderly are more likely to have hypertrophy, prostatic obstruction and age-related renal impairment and may be more susceptible to the undesirable effects due to opioid-induced urinary retention and the respiratory effects of opioid analgesics. Dose reduction may be required.

## Paediatric use

Mersyndol Forte is contraindicated for use in children:

- younger than 12 years.
- aged between 12 – 18 years in whom respiratory function might be compromised, including post tonsillectomy and/or adenoidectomy for obstructive sleep apnoea. Respiratory depression and death have occurred in some children who received codeine following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolisers of codeine due to a CYP2D6 polymorphism.

(See also Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – CYP2D6 Metabolism.)

## Effects on laboratory tests

*Plasma amylase and lipase activity:* Codeine may cause increased biliary tract pressure, thus increasing plasma amylase and/or lipase concentrations.

*Gastric emptying studies:* Gastric emptying is delayed by codeine so gastric emptying studies will not be valid.

*Uric acid and blood glucose:* Intake of paracetamol may affect the laboratory determination of uric acid by phosphotungstic acid and of blood glucose by glucose oxidase-peroxidase.

## 4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS

*Salicylates and NSAIDs:* Prolonged concurrent use of paracetamol and salicylates or non-steroidal anti-inflammatory drugs may increase the risk of adverse renal effects.

*Coumarins:* Paracetamol may increase the risk of bleeding in patients taking warfarin and other coumarin derivatives (antivitamin K). Monitoring of coagulation and bleeding complications is required.

*Chloramphenicol:* Paracetamol may slow down the excretion of chloramphenicol, entailing the risk of increased toxicity.

*Diflunisal:* Diflunisal may increase the plasma concentrations of paracetamol by 50%.

*Anticholinergics:* Concomitant use of codeine and anticholinergic agents may increase the risk of severe constipation and/or urinary retention. Drugs, which decrease gastric emptying, may decrease the absorption of paracetamol.

*Cholestyramine:* Cholestyramine reduces the absorption of paracetamol if given within one hour of paracetamol administration.

*Chelating resin:* Chelating resin can decrease the intestinal absorption of paracetamol and potentially decrease its efficacy if taken simultaneously. In general, there must be an interval of more than 2 hours between taking the resin and taking paracetamol, if possible.

*Propantheline:* Decreases gastric emptying which may decrease the absorption of paracetamol.

*Rifampicin:* Concomitant use may increase the likelihood of paracetamol toxicity (see Hepatotoxic drugs and liver microsomal enzymes below).

*Flucloxacillin:* Co-administration of flucloxacillin with paracetamol may lead to metabolic acidosis, particularly in patients presenting risk factors of glutathione depletion, such as sepsis, malnutrition or chronic alcoholism.

*Alcohol:* Codeine may potentiate the effects of alcohol and increase the likelihood of paracetamol toxicity (see Hepatotoxic drugs and liver microsomal enzymes below). The concomitant use of alcohol and opioids increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. Concomitant use with alcohol is not recommended (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE).

*Metoclopramide:* Codeine may antagonise the effects of metoclopramide on gastrointestinal motility. Paracetamol absorption is increased by drugs which increase gastric emptying.

*Domperidone:* The absorption rate of paracetamol may be increased by domperidone.

*Opioid analgesics:* Concurrent use of codeine and other opioid agonists is usually inappropriate as additive CNS depression, respiratory depressant and hypotensive effects may occur. Narcotic analgesics may decrease gastric emptying and therefore decrease the absorption of paracetamol.

*Morphinic agonists-antagonists:* Concomitant use of codeine with a partial agonist (e.g. buprenorphine) or antagonist (e.g. naltrexone) can precipitate or delay codeine effects.

*Hepatotoxic drugs and liver microsomal enzyme inducers:* The risk of paracetamol toxicity may be increased in patients receiving other potentially hepatotoxic drugs or drugs that induce liver microsomal enzymes, such as antiepileptics (such as phenobarbital, phenytoin, carbamazepine, topiramate), alcohol, barbiturates and rifampicin. The induced metabolism results in an elevated production of the hepatotoxic oxidative metabolite of paracetamol. Hepatotoxicity will occur if this metabolite exceeds the normal glutathione binding capacity.

*Zidovudine:* When used concurrently with zidovudine, an increased tendency for neutropenia may develop. Combination of Mersyndol Forte and zidovudine should be avoided.

*Antiperistaltic antidiarrhoeals (including kaolin, pectin, loperamide):* Concurrent use of these agents with codeine may increase the risk of severe constipation and CNS depression.

*Tranquillisers, sedatives, hypnotics, General anaesthetics and CNS depressants:* Codeine may potentiate the effects of these drugs. Concomitant use of tranquillisers or sedatives may enhance the potential respiratory depressant effects of codeine (see Section 4.4 SPECIAL

**WARNINGS AND PRECAUTIONS FOR USE – Risks from concomitant use of benzodiazepines or other CNS depressants, including alcohol.)**

*Benzodiazepines:* The concomitant use of benzodiazepines and opioids increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. Limit dosage and duration of concomitant use of benzodiazepines and opioids (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE).

*Monoamine Oxidase Inhibitors:* Non-selective MAOI's intensify the effects of opioid drugs, which can cause anxiety, confusion and significant respiratory depression and can potentiate the central nervous effects and other side effects of unpredictable severity. Severe and sometimes fatal reactions have occurred in patients concurrently administered MAO inhibitors and pethidine. Codeine should not be given to patients taking non-selective MAOI's or within 14 days of stopping such treatment. As it is unknown whether there is an interaction between the selective MAOI's (Reversible Inhibitors of Monoamine Oxidase A) and codeine, caution is advised with this drug combination.

*Tricyclic antidepressants:* A codeine-induced respiratory depression can be potentiated by tricyclic antidepressants (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – Risks from concomitant use of benzodiazepines or other CNS depressants, including alcohol.).

*Antihypertensives:* Hypotensive effects of antihypertensive agents may be potentiated when used concurrently with codeine and lead to orthostatic hypotension.

*Neuromuscular blocking agents:* Codeine may enhance the effects of neuromuscular blocking agents resulting in increased respiratory depression.

Patients receiving other narcotic analgesics, antitussives, antihypertensives, antihistamines, antipsychotics, antianxiety agents, gabapentinoids, cannabis, centrally-active anti-emetics or other CNS depressants (including alcohol) concomitantly with Mersyndol Forte may experience additive CNS depression (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – Risks from concomitant use of benzodiazepines or other CNS depressants, including alcohol.)

*CYP2D6 inhibitors:* Codeine is metabolized by the liver enzyme CYP2D6 to its active metabolite morphine. Medicines that inhibit CYP2D6 activity may reduce the analgesic effect of codeine. Patients taking codeine and moderate to strong CYP2D6 inhibitors (such as quinidine, fluoxetine, paroxetine, bupropion, cinacalcet, methadone) should be adequately monitored for reduced efficacy and withdrawal signs and symptoms. If necessary, an adjustment of the treatment should be considered.

*CYP3A4 inducers:* Medicines that induce CYP3A4 activity may reduce the analgesic effect of codeine. Patients taking codeine and CYP3A4 inducers (such as rifampin) should be adequately monitored for reduced efficacy and withdrawal signs and symptoms. If necessary, an adjustment of the treatment should be considered.

## 4.6 FERTILITY, PREGNANCY AND LACTATION

### Effects on fertility

See Section 5.3 PRECLINICAL SAFETY DATA.

### Use in pregnancy

Category A. There have been no observations of an increase in the frequency of malformations or other direct or indirect harmful effects on the foetus in pregnant women and women of child-bearing age who have taken those drugs found in Mersyndol Forte.

However, prolonged high-dose use of codeine prior to delivery may prolong codeine withdrawal symptoms in the neonate.

Regular use during pregnancy may cause physical dependence in the foetus, leading to withdrawal symptoms in the neonate. Administration of codeine during labour may cause respiratory depression in the newborn infant. Codeine may cause respiratory depression and withdrawal syndrome in neonates born to mothers who use codeine during the third trimester of pregnancy. As a precautionary measure, use of Mersyndol Forte should be avoided during the third trimester of pregnancy and during labour. Codeine is contraindicated in the event of impending childbirth or in case of risk of premature birth (see Section 4.3 CONTRAINDICATIONS). Mersyndol Forte should only be used during pregnancy under medical supervision if the potential benefit justifies the potential risk to the foetus. If administered during pregnancy, morphinomimetic properties of codeine should be taken into account.

### Use in lactation

Mersyndol Forte is contraindicated during breast-feeding (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – CYP2D6 Metabolism) due to risk of respiratory depression in the infant. Codeine, doxylamine and paracetamol are excreted in breast milk. Analgesic doses excreted in breast milk are generally low. However, codeine is partially metabolized by cytochrome P450 2D6 (CYP2D6) into morphine, which is excreted into breast milk. If nursing mothers are CYP2D6 ultra-rapid metabolisers, higher levels of morphine may be present in their breast milk. This may result in symptoms of opioid toxicity in both mother and the breast-fed infant. Life-threatening adverse events or neonatal death may occur even at therapeutic doses (see Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE – CYP2D6 Metabolism).

Therefore, Mersyndol Forte is contraindicated for use during breastfeeding. However, in circumstances where a breastfeeding mother requires codeine therapy, breastfeeding should be suspended and alternative arrangements should be made for feeding the infant for any period during codeine treatment.

Breast feeding mothers should be told how to recognise signs of high morphine levels in themselves and their babies. For example, in a mother, symptoms include extreme sleepiness and trouble caring for the baby. In the baby, symptoms include signs of increased sleepiness (more than usual), difficulty breastfeeding, breathing difficulties, or limpness. Medical advice should be sought immediately.

#### 4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

Mersyndol Forte may cause drowsiness, disturbances of visuomotor coordination and visual acuity and/or dizziness. Due to the preparation's sedative action, psychomotor impairment impacting the mental and/or physical abilities required for the performance of potentially hazardous activities may occur. Patients treated with this medicine should not drive, operate machinery, or drink alcohol whilst taking this medication.

#### 4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)

Reports of adverse reactions are rare. Although the following reactions have been reported when paracetamol and codeine have been administered:

##### *Haematologic*

Less frequent to rare – agranulocytosis, anaemia, thrombocytopenia

##### *Genitourinary*

Less frequent to rare – renal failure, uraemia, urinary retention or hesitance

##### *Hypersensitive*

Less frequent to rare – skin rashes and other allergic reactions, histamine release (hypotension, flushing of the face, tachycardia, breathlessness)

##### *Gastrointestinal*

Common – constipation, nausea, vomiting

##### *Neurological*

Common – drowsiness, dizziness

Less frequent to rare - euphoria, dysphoria. At higher doses codeine may cause respiratory depression.

##### *Hepatic*

Very rare – pancreatitis

##### *Metabolism and nutrition system disorders*

Not known	Pyroglutamic acidosis, in patients with pre-disposing factors for glutathione depletion.
-----------	--

Paracetamol has also been associated with dyspepsia, sweating, erythema, urticaria, anaphylactic shock, angioneurotic oedema, angioedema, leukopenia, neutropenia and pancytopenia. Bronchospasms may be triggered in patients having a tendency of analgesic asthma. Toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), acute generalised exanthematous pustulosis, fixed drug eruption and cytolytic hepatitis, which may lead to acute hepatic failure, have also been reported. Large doses may produce hepatotoxicity.

Prolonged administration of high doses may result in drug dependence.

Haemolytic anaemia, particularly in patients with underlying glucose 6-phosphate-dehydrogenase deficiency has been reported. Kounis syndrome and bronchospasm have also been reported.

Codeine can cause confusional state, dysphoria, seizure, headache, somnolence, sedation, miosis, tinnitus, dry mouth, pruritus, fatigue, and hypotension. Visuomotor coordination and visual acuity may be adversely affected in a dose-dependent manner at higher doses or in particular sensitive patients. Long term use also entails the risk of drug dependence.

The following have been reported in relation to doxylamine: somnolence, paradoxical stimulation, psychomotor impairment, blurred vision, thickened respiratory mucus, gastrointestinal disorders and urinary retention.

### **Reporting suspected adverse effects**

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

## **4.9 OVERDOSE**

Elderly persons, small children, patients with liver disorders, chronic alcohol consumption or chronic malnutrition, as well as patients concomitantly treated with enzyme-inducing drugs are at an increased risk of intoxication, including fatal outcome.

The intake of too high a dose of paracetamol can lead to impairment of liver function due to necrosis of liver cells, and this condition may progress to hepatic coma and may be fatal. Independently of these liver lesions, kidney damage as a result of necrosis of the renal tubules has also been reported.

### **Symptoms and signs of poisoning**

The early stage of acute poisoning is typically characterized by nausea, vomiting, anorexia, pallor, abdominal pain and sweating and general malaise. This is usually followed by a 24 to 48 hour period in which the patient feels better; the symptoms however do not disappear altogether. The following stage is characterized by rapid increase in the size of the liver; serum transaminases and bilirubin rise, prothrombin time increases abnormally; urine excretion decreases and a slight increase in nitrogenous substances may develop. 3 to 5 days after poisoning, the clinical features most typically encountered are commonly jaundice, fever, foetor hepaticus, abnormal bleeding tendency, hypoglycaemia, etc., as well as all stages of hepatic encephalopathy.

Nausea, vomiting, anorexia, pallor and abdominal pain generally appear during the first 24 hours of overdosage with paracetamol. Overdosage with paracetamol may cause hepatic cytolysis which can lead to hepatocellular insufficiency, gastrointestinal bleeding, metabolic acidosis, encephalopathy, disseminated intravascular coagulation, coma and death. Increased levels of hepatic transaminases, lactate dehydrogenase and bilirubin with a reduction in

prothrombin level can appear 12 to 48 hours after acute overdose. It can also lead to pancreatitis, acute renal failure and pancytopenia.

Reactions associated with doxylamine succinate overdose may vary from central nervous depression to stimulation. Stimulation is particularly likely in children. Atropine-like signs and symptoms - dry mouth; fixed, dilated pupils; flushing and gastrointestinal symptoms may also occur. Severe rhabdomyolysis after doxylamine succinate overdose has been reported in humans.

In an evaluation of codeine intoxication in children, symptoms ranked by decreasing order of frequency included: sedation, rash, miosis, vomiting, itching, ataxia and swelling of the skin. Respiratory failure may occur. Blood concentrations of codeine ranged from 1.4 to 5.6 micrograms per mL in eight adults whose deaths were attributed to codeine overdose.

The ingestion of very high doses of codeine can cause initial excitation, anxiety, insomnia followed by drowsiness in certain cases, areflexia progressing to stupor or coma, headache, miosis, alterations in blood pressure, arrhythmias, dry mouth, hypersensitivity reactions, cold clammy skin, bradycardia, tachycardia, convulsions, gastrointestinal disorders, nausea, vomiting and respiratory depression.

Severe intoxication can lead to apnoea, circulatory collapse, cardiac arrest and death.

### **Treatment of poisoning**

Despite lack of significant early symptoms, patients should be referred to hospital urgently for immediate medical attention.

Consists primarily of management of paracetamol toxicity; naloxone hydrochloride is the treatment of choice for codeine intoxication and must be administered intravenously. In cases of overdose, methods of reducing the absorption of ingested drug are important. Prompt administration of 50 g activated charcoal and 500 mL iced mannitol 20% by mouth may reduce absorption.

Determinations of the plasma concentration of paracetamol are recommended. Plasma concentrations of paracetamol should be measured at 4 hours or later after ingestion (earlier concentrations are unreliable).

If the history suggests that 15 g paracetamol or more has been ingested, administer one of the following antidotes:

#### ***Acetylcysteine 20% i.v***

Administer 20% acetylcysteine immediately without waiting for positive urine test or plasma level results: initial dose 150 mg/kg over 15 minutes, followed by continuous infusion of 50mg/kg in 500mL 5% glucose over 4 hours and 100 mg/kg in 1L 5% glucose over 16 hours; or

#### ***Oral Methionine***

2.5 g immediately followed by three further doses of 2.5 g at four hourly intervals. For a 3-year-old child, 1 g methionine 4-hourly for four doses has been used.



If more than ten hours have elapsed since the overdose was taken, the antidote may be ineffective.

In general, treatment for codeine overdose should be symptomatic: re-establish adequate respiratory exchange by ensuring a clear airway and using mechanical ventilation. When treatment for paracetamol toxicity has been initiated naloxone 400 microgram may be administered SC, IM or IV; IV may be repeated at intervals of 2 to 3 minutes if necessary. Assisted respiration may be required.

Further measures will depend on the severity, nature and course of clinical symptoms of intoxication and should follow standard intensive care protocols.

For information on the management of overdose contact the Poisons Information Centre on 131126 (Australia).

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 PHARMACODYNAMIC PROPERTIES**

#### **Mechanism of action**

Codeine is a potent analgesic. Paracetamol is an analgesic and antipyretic. Doxylamine succinate has calmative and antinauseant properties which can be useful in relieving tension and nausea associated with pain.

#### **Clinical trials**

No data available

### **5.2 PHARMACOKINETIC PROPERTIES**

#### **Metabolism**

Patients who metabolise drugs poorly via CYP2D6 are likely to obtain reduced benefit from codeine due to reduced formation of the active metabolite.

### **5.3 PRECLINICAL SAFETY DATA**

#### **Genotoxicity**

No data available

#### **Carcinogenicity**

Toxicity studies in animals have shown that high doses of paracetamol cause testicular atrophy and inhibition of spermatogenesis; the relevance of this finding to use in humans is not known.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 LIST OF EXCIPIENTS

Other ingredients are purified talc, sodium starch glycollate and magnesium stearate.

### 6.2 INCOMPATIBILITIES

No data available.

### 6.3 SHELF LIFE

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

### 6.4 SPECIAL PRECAUTIONS FOR STORAGE

Store below 25°C.

### 6.5 NATURE AND CONTENTS OF CONTAINER


Blister packed white tablets in cartons of 20.

### 6.6 SPECIAL PRECAUTIONS FOR DISPOSAL


In Australia, any unused medicine or waste material should be disposed of by taking to your local pharmacy.

### 6.7 PHYSICOCHEMICAL PROPERTIES


#### Chemical Structure



Paracetamol



Codeine phosphate hemihydrate



Doxylamine succinate

### CAS number

Paracetamol: 103-90-2

Codeine phosphate hemihydrate: 41444-62-6

Doxylamine succinate: 562-10-7

## 7 MEDICINE SCHEDULE (POISONS STANDARD)

Prescription Only Medicine (Schedule 4)

## 8 SPONSOR

sanofi-aventis australia pty ltd  
 12-24 Talavera Road  
 Macquarie Park NSW 2113  
 Freecall: 1800 818 806  
 Email: medinfo.australia@sanofi.com

## 9 DATE OF FIRST APPROVAL

8 July 1991

## 10 DATE OF REVISION

26 August 2024

### SUMMARY TABLE OF CHANGES

Section Changed	Summary of new information
4.4	Safety update to add opioid induced hyperalgesia and/or allodynia