

# PANADOL Optizorb Caplets

# PANADOL Optizorb Tablets

500 mg Paracetamol

---

## INFORMATION LEAFLET

*Please read this information before you start using this medicine.*

---

### What is in this leaflet

This leaflet answers some common questions about PANADOL OPTIZORB. It does not contain all the available information.

It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits.

**If you have any concerns about using this medicine, ask your pharmacist or doctor.**

**Keep this leaflet with the medicine.** You may need to read it again.

---

### What PANADOL OPTIZORB is used for

The active ingredient in these medicines is paracetamol. Paracetamol is used for the fast effective temporary relief of pain and discomfort associated with

- Headache
- Muscular aches
- Period pain
- Arthritis/Osteoarthritis
- Toothache
- Migraine headache
- Colds and flu
- Tension headache
- Sinus pain/headache
- Backache

Paracetamol also reduces fever.

Paracetamol works to stop the pain messages from getting through to the brain. It also acts in the brain to reduce fever.

**Ask your pharmacist or doctor if you have any questions about this medicine.**

Your pharmacist or doctor may have given it for another reason

---

### Before you use PANADOL OPTIZORB

Do not take more than the recommended dose as it may cause serious harm to your liver.

### *When you must not use it*

**Do not use PANADOL OPTIZORB if you have an allergy to:**

- paracetamol or any of the ingredients listed at the end of this leaflet

Do not use this medicine if you are taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu, or to aid sleep.

**Always read and follow the label.**

**Do not use this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.** If it has expired or is damaged, return it to your pharmacist for disposal.

**If you are not sure whether you should start using this medicine, talk to your pharmacist or doctor.**

### *Before you start to use it*

**Tell your pharmacist or doctor if you have or have had any of the following medical conditions:**

- Liver or kidney disease
- Are underweight or malnourished
- Regularly drink alcohol  
You may need to avoid using this product altogether or limit the amount of paracetamol that you take
- You have a severe infection, are severely malnourished or are a chronic heavy alcohol user as this may increase the risk of metabolic acidosis. Signs of metabolic acidosis include:
  - deep, rapid, difficult breathing
  - feeling sick (nausea), being sick (vomiting)
  - loss of appetiteContact a doctor immediately if you get a combination of these symptoms.

Please see your doctor if your symptoms do not improve.

Keep out of sight and reach of children.

**Ask your pharmacist or doctor about using paracetamol if you are pregnant or plan to become pregnant, or breastfeeding.** Paracetamol may be used during pregnancy and if you are breastfeeding but you should always consult your doctor first.

Consider taking the lowest effective dose for the shortest period of time.

**If you have not told your pharmacist or doctor about any of the above, tell them before you use PANADOL OPTIZORB.**

### *Using other medicines*

Tell your pharmacist or doctor if you are using any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and paracetamol may interfere with each other. These include:

- Warfarin, a medicine used to prevent blood clots
- Metoclopramide, a medicine used to control nausea and vomiting
- Medicines used to treat epilepsy or fits
- Chloramphenicol, an antibiotic used to treat ear and eye infections
- Alcohol
- Probenecid, a medicine used to treat gout or sometimes given with an antibiotic
- Cholestyramine, a medicine used to treat high cholesterol levels in the blood

Your pharmacist and doctor will have more information on these and other medicines to be careful with or avoid while using this medicine.

---

## **How to use PANADOL OPTIZORB**

---

**Follow all directions given to you by your pharmacist or doctor carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box, ask your pharmacist or doctor for help.**

**Do not exceed the stated dose.**

**Use the smallest dose that you need to treat your symptoms and use the medicine for the shortest period of time necessary.**

### *How much to use*

Adults and children aged 12 years and over: Take 1 to 2 caplets or tablets every four to six hours as needed. Do not take more than 8 caplets or tablets in 24 hours.

Children 7 to 11 years: Take ½ to 1 caplet or tablet every four to six hours as needed. Do not take more than 4 caplets or tablets in 24 hours.

Do not take more frequently than every 4 hours. Not recommended in children under 7 years.

Swallow with water or other fluid.

### *How long to use it*

Adults and children aged 12 years and over:

Only take paracetamol for a few days at a time unless your doctor tells you to take it for longer.

Children 7 to 11 years:

Only give paracetamol to children for up to 48 hours at a time unless a doctor has told you to give it for longer.

### *If you use too much (overdose)*

**Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26 for Australia, 0800 764 766 for New Zealand) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too**

**much PANADOL OPTIZORB. Do this even if you feel well because of the risk of delayed, serious liver damage if left untreated.**

You may need urgent medical attention.

---

## **While you are using PANADOL OPTIZORB**

---

### *Things you must do*

**Talk to your pharmacist or doctor if your symptoms do not improve.**

Your pharmacist or doctor will assess your condition and decide if you should continue to take the medicine.

### *Things you must not do*

**Children:**

**Do not give paracetamol for more than 48 hours at a time unless a doctor has told you to.**

**Adults:**

**Do not use for more than a few days at a time unless your doctor tells you to.**

**Do not take more than the recommended dose as it may cause serious harm to your liver.**

**Do not use PANADOL OPTIZORB to treat any other complaints unless your pharmacist or doctor tells you to.**

### *Things to be careful of*

**Only drink small quantities of alcohol (beer, wine or spirits) while using paracetamol. Drinking large quantities of alcohol while taking paracetamol may increase the risk of liver side effects.**

---

## Side Effects

---

**Tell your pharmacist or doctor as soon as possible if you do not feel well while you are using PANADOL OPTIZORB.** This medicine helps most people with various types of pain but it may have unwanted side effects. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

**Do not be alarmed by the following lists of side effects. You may not experience any of them.**

**Ask your pharmacist or doctor to answer any questions you may have.**

If any of the following happen, tell your pharmacist or doctor immediately or go to Accident and Emergency at your nearest hospital:

- Shortness of breath
- Wheezing or difficulty breathing
- Swelling of the face, lips, tongue, throat or other parts of the body
- Rash, peeling, itching or hives on the skin or mouth ulcers
- Unexplained bruising or bleeding
- you have experienced breathing problems with aspirin or non-steroidal anti-inflammatories and experience a similar reaction with this product

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are very rare for low doses of this medicine and when used for a short period of time.

**Tell your pharmacist or doctor if you notice anything that is making you feel unwell.**

Other side effects not listed above may also occur in some people.

---

## After using PANADOL OPTIZORB

---

### Storage

**Keep your medicine in the original pack until it is time to take it.**

**Keep your medicine in a cool dry place where the temperature stays below 30°C.**

**Do not store PANADOL OPTIZORB or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in a car.** Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### Disposal

**Ask your pharmacist what to do with any medicine that is left over, or if the expiry date has passed.**

---

## Product Description

---

### What it looks like

**PANADOL OPTIZORB Caplets** are a white, to off-white, film-coated capsule-shaped tablet with convex edges. They are marked with a “P” in a circle on one side and a break-line on the other side. They come in blister packs of 16 and 48 caplets.

**PANADOL OPTIZORB Tablets** are a white, to off-white film-coated round shaped tablet. They are marked with a “P” in a circle on one side and a break-line on the other side. They come in blister packs of 16 and 48 tablets.

Not all presentations may be marketed.

### Ingredients

**PANADOL OPTIZORB Caplets** contain 500 mg of paracetamol as the active ingredient.

They also contain:

- Alginic acid
- Calcium carbonate
- Carnauba wax
- Crospovidone
- Magnesium stearate
- White colouring (Opadry YS-1-7003)
- Povidone
- Silica –colloidal anhydrous
- Starch – pregelatinised maize

**PANADOL OPTIZORB tablets** contain 500 mg of paracetamol as the active ingredient.

They also contain:

- Alginic acid
- Calcium carbonate
- Carnauba wax
- Crospovidone
- Magnesium stearate
- White colouring (Opadry YS-1-7003)
- Povidone
- Silica –colloidal anhydrous
- Starch – pregelatinised maize

### Manufacturer/Supplier

Haleon Australia  
Level 48, 8 Parramatta Square,  
10 Darcy Street, Parramatta NSW  
2150 and Auckland, New Zealand

AUST R 356011 (Caplets)  
AUST R 356022 (Tablets)

Date of preparation: August 2024

Trademarks are owned by or licensed to Haleon.