

PROGRAF®; ADVAGRAF® XL

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using PROGRAF or ADVAGRAF XL?

PROGRAF or ADVAGRAF XL contains the active ingredient tacrolimus. PROGRAF or ADVAGRAF XL is used so that your new liver, kidney, lung or heart will not be attacked or rejected by your body. For more information, see Section [1. Why am I using PROGRAF or ADVAGRAF XL?](#) in the full CMI.

2. What should I know before I use PROGRAF or ADVAGRAF XL?

Do not use if you have ever had an allergic reaction to PROGRAF or ADVAGRAF XL, macrolide antibiotics, or any of the ingredients listed at the end of the CMI. ADVAGRAF XL and PROGRAF are not interchangeable.

For more information, see Section [2. What should I know before I use PROGRAF or ADVAGRAF XL?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with PROGRAF or ADVAGRAF XL and affect how it works.

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop. See [3. What if I am taking other medicines?](#) in the full CMI for more information.

4. How do I use PROGRAF or ADVAGRAF XL?

- PROGRAF capsules should be taken in two doses (e.g. morning and evening) each day.
- ADVAGRAF XL should be taken once a day in the morning.
- PROGRAF or ADVAGRAF XL capsules should be taken at least 1 hour before or 2 to 3 hours after a meal preferably with water and not with grapefruit juice.

More instructions can be found in Section [4. How do I use PROGRAF or ADVAGRAF XL?](#) in the full CMI.

5. What should I know while using PROGRAF or ADVAGRAF XL?

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin	
PROGRAF® and	ADVAGRAF® cancer and other cancers	XL1 3

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 4

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	6

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	7

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 8

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 10

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	XL1	11

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	12

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 13

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	PROGRAF or ADVAGRAF	XL1 14

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	15

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	16

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	PROGRAF or ADVAGRAF	XL1 17

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	PROGRAF or ADVAGRAF	XL1 18

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	19

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 20

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	PROGRAF or ADVAGRAF	XL1 21

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 22

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	PROGRAF or ADVAGRAF	XL1 23

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	PROGRAF or ADVAGRAF	XL1 24

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	25

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 26

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 27

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 28

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	29

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 30

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	PROGRAF or ADVAGRAF	XL1 31

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 32

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	33

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 34

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and ADVAGRAF®	taking PROGRAF or ADVAGRAF	XL1 35

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	36

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	37

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	38

Things you should do	• Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL. PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while	
PROGRAF® and	ADVAGRAF® XL1 PROGRAF or ADVAGRAF	39

For more information, see Section [5. What should I know while using PROGRAF or ADVAGRAF XL?](#) in the full CMI.

6. Are there any side effects?

There are a number of side effects associated with this medicine. It is important to be aware of them so that you can identify any symptoms if they occur (see the full CMI for more details). The serious side effects are: signs of allergy; fever; diabetes / increased blood sugar; swelling, numbness or tingling in your hands and feet; constant "flu-like" symptoms; unusual bleeding or bruising; high blood pressure; palpitations, abnormal heart rhythms, chest pain; new lumps or moles, or changes to existing moles; swelling of the eyelids, hands or feet due to excess fluid; a change in the amount of urine passed or in the number of times you urinate, pain on urinating, or other kidney problems; yellowing of the skin and/or eyes (jaundice); symptoms of anaemia; seizures (fits); buzzing or ringing in the ears, difficulty hearing, skin changes such as new or changing discolouration, lesions or lumps. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

PROGRAF® Capsules, Concentrated Injection ADVAGRAF® XL Prolonged-Release Capsules

Active ingredient(s): *tacrolimus*

Consumer Medicine Information (CMI)

This leaflet provides important information about using PROGRAF capsules, ADVAGRAF XL Prolonged-Release capsules and PROGRAF Concentrated Injection. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking PROGRAF or ADVAGRAF XL against the benefits this medicine is expected to have for you.

You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using PROGRAF or ADVAGRAF XL.

Keep this leaflet with your medicine. You may need to read it again.

Where to find information in this leaflet:

[1. Why am I using PROGRAF or ADVAGRAF XL?](#)

[2. What should I know before I use PROGRAF or ADVAGRAF XL?](#)

[3. What if I am taking other medicines?](#)

[4. How do I use PROGRAF or ADVAGRAF XL?](#)

[5. What should I know while using PROGRAF or ADVAGRAF XL?](#)

[6. Are there any side effects?](#)

[7. Product details](#)

1. Why am I using PROGRAF or ADVAGRAF XL?

PROGRAF or ADVAGRAF XL contains the active ingredient tacrolimus, which is an immunosuppressive agent.

You have been given a new transplanted liver, kidney, lung or heart from another person because your own was no longer healthy. Your body recognises that this new organ is different from your other organs and will try to reject it by attacking it in the same way that it would attack germs that enter your body. This could make you become ill again. PROGRAF or ADVAGRAF XL stops this attack; it is very important to take PROGRAF or ADVAGRAF XL given to you by your doctor regularly so that your new liver, kidney, lung or heart will not be attacked or rejected.

If you have been taking other medicines for this purpose, but are still feeling unwell, your doctor may change your treatment and begin giving you PROGRAF or ADVAGRAF XL.

Your doctor may have prescribed PROGRAF or ADVAGRAF XL for another reason.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

2. What should I know before I use PROGRAF or ADVAGRAF XL?

Warnings

Do not use PROGRAF or ADVAGRAF XL if:

- you are allergic to any medicine containing tacrolimus or other macrolides (these are antibiotics of the erythromycin family - trade names are Eryc, EES, Klacid, Zithromax, Rulide and Biaxsig), or any of the ingredients listed at the end of this leaflet.
- Always check the ingredients to make sure you can use this medicine.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

Do not use PROGRAF or ADVAGRAF XL if the packaging is torn or shows signs of tampering. Do not use PROGRAF or ADVAGRAF XL beyond the expiry date printed on the pack.

Check with your doctor if:

- you are receiving cyclosporin immunosuppressive therapy.

If you have not told your doctor or pharmacist about any of the above, tell them before you start taking or are given PROGRAF or ADVAGRAF XL.

Your doctor will advise you whether or not to take PROGRAF or ADVAGRAF XL or if you need to adjust the dose or adapt your treatment.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

Check with your doctor if you are using oral contraceptives.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with PROGRAF or ADVAGRAF XL and affect how it works.

In particular, you should tell your doctor if you are taking or have recently taken medicines like:

- antifungal medicines and antibiotics, particularly so-called macrolide antibiotics, used to treat infections e.g., ketoconazole, fluconazole, itraconazole, voriconazole, clotrimazole, miconazole, caspofungin, erythromycin, clarithromycin, josamycin, rifampicin, rifabutin, isoniazid and flucloxacillin
- letermovir, used to prevent illness caused by CMV (human cytomegalovirus)
- HIV protease inhibitors (e.g., ritonavir, nelfinavir, saquinavir), the booster medicine cobicistat, and combination tablets, or HIV non-nucleoside reverse transcriptase inhibitors (efavirenz, etravirine, nevirapine) used to treat HIV infection
- HCV protease inhibitors (e.g., telaprevir, boceprevir, the combination ombitasvir/paritaprevir/ritonavir with or without dasabuvir, elbasvir/grazoprevir, and glecaprevir/pibrentasvir), used to treat hepatitis C infection

- mycophenolic acid used to suppress the immune system to prevent transplant rejection
- medicines for stomach ulcer and acid reflux (e.g., omeprazole, lansoprazole or cimetidine)
- antiemetics, used to treat nausea and vomiting (e.g., metoclopramide)
- cisapride or the antacid magnesium-aluminium-hydroxide, used to treat heartburn
- the contraceptive pill or other hormone treatments with ethinylestradiol, hormone treatments with danazol
- medicines used to treat high blood pressure or heart problems (e.g., nifedipine, nicardipine, diltiazem and verapamil)
- anti-arrhythmic drugs (amiodarone) used to control arrhythmia (uneven beating of the heart)
- medicines known as “statins” used to treat elevated cholesterol and triglycerides
- carbamazepine, phenytoin or phenobarbital, used to treat epilepsy
- metamizole, used to treat pain and fever
- the corticosteroids prednisolone and methylprednisolone, belonging to the class of corticosteroids used to treat inflammations or suppress the immune system (e.g., in transplant rejection)
- nefazodone, used to treat depression

- herbal preparations containing St. John's wort (*Hypericum perforatum*) or extracts of *Schisandra sphenanthera*
- cannabidiol.

Tell your doctor if you are taking or need to take ibuprofen, amphotericin B, antibiotics (cotrimoxazole, vancomycin, or so-called aminoglycoside antibiotics such as gentamicin), or antivirals (e.g., acyclovir, ganciclovir, cidofovir, or foscarnet). These may worsen kidney or nervous system problems when taken together with PROGRAF or ADVAGRAF XL.

Your doctor also needs to know if you are taking potassium supplements or potassium-sparing diuretics (e.g., amiloride, triamterene, or spironolactone), or the antibiotics trimethoprim or cotrimoxazole that may increase levels of potassium in your blood, certain pain killers (so-called NSAIDs, e.g., ibuprofen), anticoagulants, or oral medication for diabetic treatment, while you receive PROGRAF or ADVAGRAF XL.

Your doctor or pharmacist can tell you what to do if you are taking any of these medicines.

4. How do I use PROGRAF or ADVAGRAF XL?

How much to take

- You can only get PROGRAF or ADVAGRAF XL from your doctor. Your dose will be calculated according

to your weight, age, and medical condition. As your health and the function of your new liver, kidney, lung or heart can be affected by how much medicine you take, it is normal that your doctor tests samples of your blood and urine at regular intervals. This is in order to test whether your medicine requires adjustment.

When to take PROGRAF or ADVAGRAF XL

- PROGRAF should be taken in two doses (e.g., morning and evening) each day.
- ADVAGRAF XL should be taken once a day in the morning.

How to take PROGRAF or ADVAGRAF XL?

- Take the capsule from the blister pack and swallow it whole with plenty of water. Do not use grapefruit juice as it contains substances that interfere with the action of PROGRAF or ADVAGRAF XL.
- PROGRAF or ADVAGRAF XL capsules should generally be taken on an empty stomach or at least 1 hour before or 2 to 3 hours after a meal.
- If you receive PROGRAF Concentrated Injection by intravenous infusion, it will be added to a larger volume of diluent and infused over a prolonged period of time (usually 24 hours).
- You must never change the dose yourself even if you are feeling better. It is very important that you keep taking this medicine so that your body will not reject your new liver, kidney, lung or heart.

- If you accidentally take a larger dose than recommended, tell your doctor immediately.
- If you do not understand the instructions provided with this medicine, ask your doctor or pharmacist for help.

If you forget to take PROGRAF or ADVAGRAF XL

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed.

If you have missed more than one dose, or are not sure what to do, check with your doctor or pharmacist.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you take too much PROGRAF or ADVAGRAF XL

If you think that you have used too much PROGRAF or ADVAGRAF XL, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

5. What should I know while using PROGRAF or ADVAGRAF XL?

Things you should do

- Always follow your doctor's instructions carefully.
- Tell your doctor if you become pregnant while taking PROGRAF or ADVAGRAF XL.
- If you are about to start taking a new medicine, tell your doctor and pharmacist that you are taking PROGRAF or ADVAGRAF XL.
- PROGRAF or ADVAGRAF XL suppresses your immune system by lowering your body's immune defence system. This increases your risk of skin cancer and other cancers while taking PROGRAF or ADVAGRAF XL. You should always protect yourself from the sun, wear sunscreen, a hat and protective clothing.

Things you should not do

- Do not take PROGRAF or ADVAGRAF XL to treat any other complaint unless your doctor says so.
- Do not give this medicine to anyone else, even if their symptoms are similar to yours.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how PROGRAF or ADVAGRAF XL affects you.

PROGRAF or ADVAGRAF XL may cause visual or nervous disturbances. If affected, do not drive or operate machinery.

Effects of food and alcohol

Food reduces the absorption of PROGRAF or ADVAGRAF XL. Therefore, the capsules should be taken at least 1 hour before or 2 to 3 hours after a meal.

Drinking alcohol

Tell your doctor if you drink alcohol.

Looking after your medicine

Use all the PROGRAF capsules within 3 months of opening the aluminium wrapper.

Use all the ADVAGRAF XL capsules within 12 months of opening the aluminium wrapper.

Keep PROGRAF or ADVAGRAF XL capsules in the blisters until it is time to take them.

Keep PROGRAF capsules in a cool dry place where the temperature is below 30°C.

Keep ADVAGRAF XL capsules in a cool dry place where the temperature is below 25°C.

Keep PROGRAF Concentrated Injection away from light and store below 25°C.

Keep your medicines where children cannot reach them. A locked cupboard at least one-and-a-half metres (1.5 m) above the ground is a good place to store medicines.

Do not store PROGRAF or ADVAGRAF XL, or any other medicine, in the bathroom or near a sink. Do not leave medicines in the car or on window sills. Heat and dampness can destroy some medicines.

Keep it where young children cannot reach it.

When to discard your medicine (as relevant)

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Less serious side effects

Less serious side effects	What to do
• tiredness, lack of energy • stomach upset, including nausea (feeling sick), vomiting, loss of appetite, diarrhoea, stomach cramps • tremor (shaking) • headache • feeling depressed (sad) • sleeping difficulties • blurred vision or sensitive to light • muscle cramps, tenderness or weakness	Speak to your doctor if you have any of these less serious side effects and they worry you.

Serious side effects

Serious side effects	What to do
• signs of allergy such as rash, itching or hives on the skin; swelling of the face, lips, tongue or other part of the body; shortness of breath, wheezing or troubled breathing • fever • diabetes / increased blood sugar • swelling, numbness or tingling (pins and needles) in your hands and feet • constant "flu-like" symptoms such as chills, sore throat, aching joints, swollen glands, or any other signs of infection • unusual bleeding or bruising • high blood pressure	Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.

Serious side effects	What to do
• palpitations, abnormal heart rhythms, chest pain • new lumps or moles, or changes to existing moles, anywhere on the body • swelling of the eyelids, hands or feet due to excess fluid • a change in the amount of urine passed or in the number of times you urinate, pain on urinating, or other kidney problems • yellowing of the skin and/or eyes (jaundice) often accompanied by generally feeling unwell (for example, tiredness, lack of energy, loss of appetite, nausea and vomiting, pain in the abdomen) • symptoms of anaemia, such as shortness of breath, tiredness or dizziness • seizures (fits)	

Serious side effects	What to do
• buzzing or ringing in the ears, difficulty hearing • fever and bruising under the skin that may appear as red pinpoint dots, with or without unexplained extreme tiredness, confusion, yellowing of the skin or eyes (jaundice), with symptoms of acute renal failure (low or no urine output) • changes in skin such as new or changing discolouration, lesions or lumps as these could be a symptom of skin or other cancers	

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at

www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What PROGRAF or ADVAGRAF XL contains

Active ingredient (main ingredient)	tacrolimus
Other ingredients (inactive ingredients)	PROGRAF capsules contain hypromellose, croscarmellose sodium, lactose monohydrate, and magnesium stearate. The capsule shell contains gelatin, titanium oxide, purified water (0.5 mg and 1 mg only), iron oxide yellow (0.5 mg capsules only) and iron oxide red (5 mg capsules only). The 0.5 mg and 1 mg capsules have a trace of

printing ink: OPACODE
monogramming ink
S-1-15083 RED (ARTG
PI No: 12388). The 5 mg
capsules have a trace of
printing ink: OPACODE
monogramming ink
S-1-18086 WHITE (ARTG
PI No: 107579).

ADVAGRAF XL
prolonged-release
capsules contain
hypromellose,
ethylcellulose, lactose
monohydrate and
magnesium stearate. The
capsule shells contain
gelatin, titanium dioxide,
iron oxide yellow, iron
oxide red and sodium
lauryl sulfate. The
capsules also have a trace
of printing ink, OPACODE
monogramming ink
S-1-15083 RED (ARTG PI
NO: 12388).

PROGRAF Concentrated
Injection contains
polyoxyethylene (PEG-60)
hydrogenated castor oil
and ethanol.

	PROGRAF and ADVAGRAF XL capsules contain excipients with known effect: sugars as lactose and soya bean products.
--	---

Do not take this medicine if you are allergic to any of these ingredients.

What PROGRAF or ADVAGRAF XL looks like

PROGRAF

- PROGRAF 0.5 mg capsules (AUST R 77280) are light yellow with "0.5 mg" and "[f]607" printed in red, packed in blister sheets of ten capsules and sealed in an aluminium wrapper (Pack size 100 capsules).
- PROGRAF 1 mg capsules (AUST R 58930) are white with "1 mg" and "[f]617" printed in red, packed in blister sheets of ten capsules and sealed in an aluminium wrapper (Pack size 100 capsules).
- PROGRAF 5 mg capsules (AUST R 58931) are greyish-red with "5 mg" and "[f]657" printed in white (Pack size 50 capsules).

ADVAGRAF XL

- ADVAGRAF XL 0.5 mg prolonged-release capsules (AUST R 269212) are oblong capsules with a light yellow cap imprinted with "0.5 mg" and an orange body imprinted with "647" (Pack size 30 capsules).

- ADVAGRAF XL 1 mg prolonged-release capsules (AUST R 269214) are oblong capsules with a white cap imprinted with "1 mg" and an orange body imprinted with "677" (Pack size 60 capsules).
- ADVAGRAF XL 3 mg prolonged-release capsules (AUST R 306501) are oblong capsules with an orange cap imprinted with "3 mg" and an orange body imprinted with "637" (Pack size: 50 capsules).
- ADVAGRAF XL 5 mg prolonged-release capsules (AUST R 269215) are oblong capsules with a greyish red cap imprinted with "5 mg" and an orange body imprinted with "687" (Pack size 30 capsules).

PROGRAF Concentrated Injection

- PROGRAF Concentrated Injection ampoules (AUST R 58932) contain 5 mg tacrolimus in 1 mL (Pack size 10 ampoules).

Who distributes PROGRAF or ADVAGRAF XL

PROGRAF and ADVAGRAF XL are supplied in Australia by:

Astellas Pharma Australia Pty Ltd

Suite 2.01, 2 Banfield Road

Macquarie Park NSW 2113

Tel: 1800 751 755 (Medical Information)

Email: aaumedinfo@astellas.com (Medical Information)

Website: www.astellas.com/au

® = Registered Trademark

This leaflet was prepared in September 2026.