

PACKAGE INSERT – Stamaril – Powder and diluent for suspension for injection

NAME OF THE MEDICINE

Yellow Fever Vaccine (Live)

DOSAGE FORM

Powder and diluent for suspension for injection.

METHOD OF PREPARATION

Preparation:

The freeze-dried powder is reconstituted with the accompanying 0.4% sodium chloride diluent contained in the syringe. The vial is shaken and, after complete dissolution, the suspension obtained is withdrawn into a separate syringe for injection.

Before administration, the reconstituted vaccine should be shaken vigorously.

Use immediately after reconstitution.

Product is for single use in one patient only. Discard any residue.

Use a separate, sterile syringe and needle for each patient to prevent transmission of blood borne infectious agents. Do not recap needles. Dispose of needles and syringes according to biohazard waste guidelines.

Compatibility:

Stamaril must not be mixed with any other injectable vaccine(s) or medical product(s).

Do not administer by intravascular injection.

IN-USE STORAGE CONDITIONS

Use immediately after reconstitution. Product is for single use in one patient only. Discard any residue.

SPONSOR

sanofi-aventis australia Pty Ltd
International Tower 3, Level 23
300 Barangaroo Avenue
Sydney NSW 2000
Phone: 1800 818 806
Email: medinfo.australia@sanofi.com

VERSION/DATE OF REVISION

Version 2.0 – 24 June 2026