

DICLOXACILLIN VIATRIS

dicloxacillin sodium

CONSUMER MEDICINE INFORMATION

What is in this leaflet

This leaflet answers some common questions about DICLOXACILLIN VIATRIS.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you taking DICLOXACILLIN VIATRIS against the benefits expected for you.

If you have any concerns about taking this medicine, talk to your doctor or pharmacist.

Keep this leaflet with your medicine.

You may need to read it again.

What DICLOXACILLIN VIATRIS is used for

DICLOXACILLIN VIATRIS is used to treat infections in different parts of the body caused by bacteria.

DICLOXACILLIN VIATRIS is an antibiotic that belongs to a group of medicines called penicillins. These medicines work by killing the bacteria that are causing your infection.

DICLOXACILLIN VIATRIS will not work against infections caused by viruses, such as colds or flu.

Ask your doctor if you have any questions about why

DICLOXACILLIN VIATRIS has been prescribed for you.

Your doctor may have prescribed DICLOXACILLIN VIATRIS for another purpose.

DICLOXACILLIN VIATRIS is not recommended for use in newborn infants as its safety and effectiveness have not been established in this age group.

DICLOXACILLIN VIATRIS is available only with a doctor's prescription.

There is no evidence that DICLOXACILLIN VIATRIS is addictive.

Before you take DICLOXACILLIN VIATRIS

When you must not take it

Do not take DICLOXACILLIN VIATRIS if you have had an allergic reaction to:

- medicines containing dicloxacillin or any other penicillin e.g. amoxicillin
- cephalosporin antibiotics
- any of the ingredients listed at the end of this leaflet

Some of the symptoms of an allergic reaction may include skin rash, itching or hives, swelling of the face, lips, mouth, throat or other parts of the body, shortness of breath, wheezing or troubled breathing.

Check with your doctor or pharmacist if you are not sure about any of the above.

Do not take DICLOXACILLIN VIATRIS if the expiry date (EXP.) printed on the bottle has passed

If you take this medicine after the expiry date, it may not work as well.

Do not take DICLOXACILLIN VIATRIS if the packaging shows signs of tampering or the capsules do not look quite right.

Before you start to take it

Be sure to tell your doctor if you have had any type of allergic reaction to:

- other penicillins, e.g. amoxicillin
- cephalosporin antibiotics

You may have an increased chance of being allergic to DICLOXACILLIN VIATRIS if you are allergic to cephalosporins.

Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.

Tell your doctor if you have, or have had, any other health problems or medical conditions, especially the following:

- stomach problems such as feeling sick (nausea), vomiting, indigestion
- problems with food absorption
- liver problems
- kidney problems
- cystic fibrosis

Your doctor may want to take special care or adjust your dose if you have any of these conditions.

Tell your doctor if you are over 55 years old.

Some side effects occur more frequently in this age group. Your doctor will discuss the risks and benefits of taking DICLOXACILLIN VIATRIS with you.

Tell your doctor if you are pregnant or plan to become pregnant.

Your doctor will discuss the risks and benefits of taking DICLOXACILLIN VIATRIS during pregnancy.

Tell your doctor if you are breastfeeding or wish to breastfeed.

Like many other medicines, DICLOXACILLIN VIATRIS can pass into breast milk. Your doctor will discuss the risks and benefits of taking DICLOXACILLIN VIATRIS when breastfeeding.

If you have not told your doctor or pharmacist about any of the above, tell them before you start taking DICLOXACILLIN VIATRIS.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines may be affected by DICLOXACILLIN VIATRIS, or may affect how well it works. These include:

- probenecid, a medicine used to treat gout
- warfarin, a medicine used to prevent blood clots
- phenytoin, a medicine used to treat epilepsy

Your doctor can tell you what to do if you are taking any of these medicines.

If you are not sure whether you are taking any of these medicines,

check with your doctor or pharmacist.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking DICLOXACILLIN VIATRIS.

How to take DICLOXACILLIN VIATRIS

Follow all directions given to you by your doctor and pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the bottle, ask your doctor or pharmacist.

How much to take

The dose varies from person to person.

Your doctor will tell you how much DICLOXACILLIN VIATRIS you need to take each day.

For most infections, the usual dose for adults and children over 12 years is 250 mg to 500 mg every 6 hours.

How to take DICLOXACILLIN VIATRIS

Swallow the capsules whole with a glass of water.

Take DICLOXACILLIN VIATRIS on an empty stomach, 1 to 2 hours before food.

Taking the capsules on an empty stomach means that more of the medicine is absorbed into your body.

Take your medicine at about the same time each day.

Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

How long to take DICLOXACILLIN VIATRIS for

Keep taking DICLOXACILLIN VIATRIS until you finish your

bottle, or for as long as your doctor recommends.

Do not stop taking DICLOXACILLIN VIATRIS, even if you feel better after a few days, unless advised by your doctor.

Your infection may not clear completely if you stop taking your medicine too soon.

If you forget to take DICLOXACILLIN VIATRIS

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your capsules as you would normally.

Do not take a double dose to make up for the dose you missed.

This may increase the chance of you getting an unwanted side effect.

If you miss more than one dose, or are not sure what to do, ask your doctor or pharmacist.

If you take too much DICLOXACILLIN VIATRIS (overdose)

Immediately telephone your doctor, or the Poisons Information Centre (telephone 13 11 26), if you think you or anyone else may have taken too much DICLOXACILLIN VIATRIS.

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention

While you are taking DICLOXACILLIN VIATRIS

Things you must do

Tell all the doctors, dentists and pharmacists who are treating you

that you are taking
DICLOXACILLIN VIATRIS.

Before starting any new medicine, tell your doctor or pharmacist that you are taking DICLOXACILLIN VIATRIS.

If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.

If you get severe diarrhoea, tell your doctor or pharmacist immediately. Do this even if it occurs several weeks after stopping DICLOXACILLIN VIATRIS.

Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care.

Do not take any medicine to stop your diarrhoea without first checking with your doctor.

If you develop a yellow discolouration of the skin or eyes, your stools become pale and you feel unwell, tell your doctor immediately.

These symptoms may mean that you have a serious problem with your liver. You may need urgent medical care.

If you get a sore, white mouth or tongue while taking or soon after stopping DICLOXACILLIN VIATRIS, tell your doctor. Also tell your doctor if you get vaginal itching or discharge.

This may mean you have a fungal infection called thrush. Sometimes, the use of DICLOXACILLIN VIATRIS allows fungi to grow and these symptoms to occur. DICLOXACILLIN VIATRIS does not work against fungi.

If you are taking more than one or two courses of DICLOXACILLIN VIATRIS, you may need to have tests to check your liver, kidneys or blood.

Always follow your doctor's instructions and have the tests performed as directed.

Things you must not do

Do not stop taking DICLOXACILLIN VIATRIS, even if you feel better after a few days, unless advised to by your doctor.

If you stop taking DICLOXACILLIN VIATRIS too soon, your infection may not clear completely and your symptoms may return.

Do not use DICLOXACILLIN VIATRIS to treat any other conditions unless advised to by your doctor.

Do not give this medicine to anyone else, even if their symptoms seem similar to yours.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking DICLOXACILLIN VIATRIS.

DICLOXACILLIN VIATRIS treats infections in most people, but it may have unwanted side effects in some people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you may have.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

While you are taking DICLOXACILLIN VIATRIS

Tell your doctor if you notice any of the following and they worry you:

- nausea (feeling sick), vomiting, stomach upset, wind, loose stools
- oral thrush (white, furry sore tongue and mouth)

- vaginal thrush (sore and itchy vagina, vaginal discharge)

Tell your doctor immediately or go to casualty at the nearest hospital if you notice any of the following:

- watery and severe diarrhoea, which may also be bloody
- blood in the urine
- excess protein in the urine
- severe stomach cramps
- yellowing of the skin or eyes (jaundice), pale stools
- kidney problems

Stop taking DICLOXACILLIN VIATRIS and tell your doctor immediately or go to casualty at the nearest hospital if any of the following happen:

- any type of skin rash, itching or hives
- swelling of the face, lips, mouth, throat or other parts of the body
- shortness of breath, wheezing or difficulty breathing

After you have finished taking DICLOXACILLIN VIATRIS

Tell your doctor immediately if you notice any of the following, even if they occur several weeks after stopping treatment with DICLOXACILLIN VIATRIS:

- watery and severe diarrhoea, which may also be bloody
- severe stomach cramps

This is not a complete list of all possible side effects. Others may occur in some people and there may be some side effects not yet known.

Tell your doctor if you notice anything that is making you feel unwell while you are taking, or soon after you have finished taking DICLOXACILLIN VIATRIS, even if it is not on this list.

After taking DICLOXACILLIN VIATRIS

Storage

Keep DICLOXACILLIN VIATRIS where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Keep your capsules in the bottle until it is time to take them.

If you take the capsules out of the bottle they will not keep well.

Keep DICLOXACILLIN VIATRIS in a cool dry place where the temperature stays below 25°C.

Do not keep DICLOXACILLIN VIATRIS or any other medicine in the bathroom or near a sink.

Do not leave DICLOXACILLIN VIATRIS in the car or on window sills.

Heat and dampness can destroy some medicines.

Disposal

If your doctor tells you to stop taking DICLOXACILLIN VIATRIS, or your medicine has passed its expiry date, ask your pharmacist what to do with any capsules that are left over.

Product description

What it looks like

DICLOXACILLIN VIATRIS comes in two strengths of capsules:

- DICLOXACILLIN VIATRIS 250 - Size 2 capsule with white opaque body and cap, printed 'DX' on the cap and '250' on the body in black
- DICLOXACILLIN VIATRIS 500 - Size 0 capsule with white opaque body and cap, printed 'DX' on the cap and '500' on the body in black

Each bottle contains 24 capsules.

Ingredients

The active ingredient in DICLOXACILLIN VIATRIS is dicloxacillin (as dicloxacillin sodium)

- each DICLOXACILLIN VIATRIS 250 capsule contains 250 mg of dicloxacillin
- each DICLOXACILLIN VIATRIS 500 capsule contains 500 mg of dicloxacillin

The capsules also contain:

- colloidal anhydrous silica
- magnesium stearate
- gelatin
- titanium dioxide
- purified water
- TekPrint SW-9008 Black Ink (ARTG PI No: 2328)

DICLOXACILLIN VIATRIS 250 mg tablets contains sulfites and 21.2 mg of sodium per dose.

DICLOXACILLIN VIATRIS 500 mg tablets contains sulfites and 42.5 mg of sodium per dose.

Supplier

DICLOXACILLIN VIATRIS is supplied by:

Alphapharm Pty Ltd trading as Viatris

Level 1, 30 The Bond
30 - 34 Hickson Road
Millers Point NSW 2000
www.viatris.com.au
Phone: 1800 274 276

This leaflet was prepared in November 2022.

Australian registration numbers:

DICLOXACILLIN VIATRIS 250:
AUST R 289105

DICLOXACILLIN VIATRIS 500:
AUST R 289107

DICLOXACILLIN VIATRIS_cmi
\Nov22/00