

# AUSTRALIAN PRODUCT INFORMATION – DBL™ Pentamidine Isethionate for Injection (Pentamidine isetionate)

## 1. NAME OF THE MEDICINE

Pentamidine isetionate

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial of DBL™ Pentamidine Isethionate for Injection contains 300 mg pentamidine isetionate as a freeze dried powder or plug.

Each 4 mg of pentamidine isetionate is equivalent to 2.3 mg pentamidine base.

For the full list of excipients, see Section 6.1.

## 3. PHARMACEUTICAL FORM

Powder for injection.

DBL™ Pentamidine Isethionate for Injection is a white or almost white, odourless or almost odourless hygroscopic powder.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Pentamidine isetionate is indicated for intravenous administration in the treatment of the following conditions:

- As an alternative first line treatment for *Pneumocystis carinii* infection in AIDS patients;
- As second line treatment for *Pneumocystis carinii* infection in non-AIDS patients;
- As second line treatment of Leishmaniasis (visceral and cutaneous), except *Leishmania aethiopica* where it may be used as first line therapy;
- As second line treatment for Trypanosomiasis (except for the *Trypanosomiasis rhodesiense* strain due to lack of efficacy).

### 4.2 Dose and method of administration

#### Dosage

The following dosage regimens are recommended.

### ***P. carinii pneumonia***

4 mg/kg bodyweight Pentamidine Isetionate once daily for 14 days, preferably by slow intravenous infusion.

### ***Leishmaniasis***

On the basis of current knowledge the following dosage are suggested however the optimal treatment regimen has yet to be established.

- Visceral (Kala-azar): 3 to 4 mg/kg bodyweight Pentamidine Isetionate on alternate days (3 times a week) to a maximum of 10 injections.
- Cutaneous: 3 to 4 mg/kg bodyweight Pentamidine Isetionate once or twice weekly, until the condition resolves.

### ***Trypanosomiasis***

Haemolymphete stage only.

4 mg/kg bodyweight Pentamidine Isetionate daily or on alternate days to a total of 7 to 10 injections.

### **Dosage adjustment**

#### ***Renal impairment***

Creatinine clearance <35 mL/min: There is little information on the kinetics or the adverse effects profile of pentamidine in patients with impaired renal function.

#### ***Hepatic impairment***

No information available.

### **Method of administration**

DBL™ Pentamidine Isethionate for Injection should be given as a slow intravenous infusion with a patient in a supine position in order to reduce the incidence of sudden severe hypotension. Direct bolus intravenous injection or rapid administration must not be used.

#### ***Reconstitution***

The contents of a 300 mg vial should be dissolved in 3 mL to 5 mL of Water for Injections. The required dose of Pentamidine Isetionate should then be diluted further in 50 to 250 mL of Glucose Intravenous Infusion 5% or Sodium Chloride Intravenous Infusion 0.9%. The reconstituted solutions should be visually examined before use. Any solutions which are hazy, discoloured or contain visible particulate matter should not be used. Diluted solutions containing Pentamidine Isetionate should be infused over a period of at least 60 minutes under close medical supervision, whilst the patient is kept lying down.

#### ***Compatibilities***

Reconstituted solutions at concentrations of 100 mg/mL and 60 mg/mL are chemically stable

for 48 hours when stored at 2 to 8°C and room temperature under fluorescent light. DBL™ Pentamidine Isethionate for Injection when reconstituted with Water for Injections and diluted to 1.0 mg/mL and 2.5 mg/mL in Sodium Chloride Intravenous Infusion 0.9% and Glucose Intravenous Infusion 5% retained its potency for at least 48 hours when stored under fluorescent light at 21 ± 2°C. However, to avoid microbial contamination, the prepared solution should be used within 24 hours.

### **4.3 Contraindications**

Patients with a known hypersensitivity to pentamidine.

Pentamidine should not be administered to patients who are pregnant or breastfeeding unless considered essential by the physician.

### **4.4 Special warnings and precautions for use**

#### **Serious Warnings and Precautions**

Severe, sometimes fatal, hypotension, hypoglycaemia, acute pancreatitis, renal impairment and cardiac arrhythmias have been reported. Other life threatening reactions requiring immediate corrective measures and withdrawal of treatment have included leucopenia (less than 1,000 per cubic millimetre), thrombocytopenia (less than 20,000 per cubic millimetre), acute renal failure, hypocalcaemia, and ventricular tachycardia. A possible case of Stevens-Johnson syndrome has been reported.

Fatalities have been documented following pentamidine administration. The ratio of therapeutic to toxic dose of pentamidine is very low and adverse effects, some of which may be life threatening, occur frequently during its use. Pentamidine Isetionate for Injection should be used only in a hospital setting with facilities to monitor blood glucose, blood count, renal function and hepatic function. Electrocardiograms should be carried out at regular intervals (see Section 4.4 Special warnings and precautions for use: Monitoring and Laboratory Tests).

Extravasation reactions may result in ulceration, tissue necrosis and long-term sequelae (see Section 4.8 Adverse effects (undesirable effects))

DBL Pentamidine is not approved for inhaled use. Bronchospasm has been reported to occur following the use of inhaled product (see section 4.8 Adverse effects (undesirable effects)). This has been particularly noted in patients who have a history of smoking or asthma. This can be controlled by prior use of bronchodilators.

Pentamidine isetionate should be used with caution in patients with the following:

1. Malnutrition
2. Hyperglycaemia or hypoglycaemia
3. Hepatic dysfunction
4. Renal dysfunction
5. Hypertension or hypotension

## 6. Anaemia, leucopenia or thrombocytopenia

### **General**

Pentamidine Isetionate for Injection should only be administered under close medical supervision, and the patient should be very carefully monitored for the development of serious adverse reactions (see Section 4.8 Adverse effects (undesirable effects)).

The administration of Pentamidine Isetionate for Injection should be limited to patients in whom *Pneumocystis jirovecii* infection has been confirmed.

Some patients may become nauseated or develop fever after taking each dose of Pentamidine Isetionate for Injection. In such cases, the prophylactic use of an antiemetic and/or paracetamol may be considered.

### **Cardiovascular**

Patients may develop sudden, severe hypotension after receiving a single intramuscular or intravenous dose of pentamidine isetionate. Therefore, patients receiving Pentamidine Isetionate for Injection should be in a supine position and the blood pressure monitored closely during administration of the drug and several times thereafter until the blood pressure is stable. Equipment for emergency resuscitation should be readily available. Pentamidine Isetionate for Injection should be infused over a period of at least 60 minutes to minimise the risk of hypotension.

Severe hypotension with accompanying bradycardia has been observed in a patient after the sixth dose of pentamidine isetionate. This hypotension did not respond to intravenous colloids, graded compression stockings or corticosteroids but resolved within four days of stopping treatment.

Ventricular tachycardia and ECG abnormalities may develop in patients receiving pentamidine isetionate. ECG's may be required at regular intervals if signs of cardiotoxicity develop.

### **Endocrine and Metabolism**

Pentamidine isetionate can produce hypoglycaemia, which may be severe, life-threatening and/or prolonged. It generally occurs after 5 to 7 days of therapy but can even occur up to several days after the drug is discontinued. The duration appears quite variable, persisting for one day to several weeks. Pentamidine isetionate-induced hypoglycemia has been associated with pancreatic islet cell necrosis and inappropriately high plasma insulin concentrations. Hyperglycemia and diabetes mellitus, with or without preceding hypoglycemia, have also occurred, sometimes several months after termination of therapy with pentamidine isetionate.

Hypoglycaemia induced by pentamidine isetionate may be controlled by intravenous administration of dextrose or (oral) diazoxide, but it is not known if such therapy can prevent the subsequent development of diabetes mellitus.

### **Gastrointestinal**

Cases of nausea and vomiting have been observed with pentamidine isetionate treatment.

## **Haematologic**

Leucopenia and thrombocytopenia, which can be severe (e.g. leucocyte count less than 1,000 per cubic millimetre, platelet count less than 20,000 per cubic millimetre), occur occasionally in patients receiving pentamidine isetionate. Cases of anaemia have been observed. In a few cases, pentamidine isetionate therapy has been associated with neutropenia.

## **Hepatic/Biliary/Pancreatic**

Abnormal liver function tests may occur. Liver function should be routinely monitored in patients receiving Pentamidine Isetionate for Injection (see Section 4.4 Special warnings and precautions for use: Monitoring and laboratory tests). Cases of acute pancreatitis have been observed with pentamidine isetionate treatment.

## **Immune**

Hypersensitivity reactions at the injection site, such as skin rash and erythema, may occur (see Section 4.4 Special warnings and precautions for use: Skin and Section 4.8 Adverse effects (undesirable effects)).

## **Skin**

Intramuscular injections are often associated with pain, tenderness, erythema, and induration at the site of injection. Sterile abscesses have been observed. Therefore, intramuscular administration should be reserved for patients with adequate muscle mass and limited to the rare situations where intravenous infusion is not feasible.

## **Vascular**

Phlebitis can occur after intravenous injection.

## **Monitoring and Laboratory Tests**

In order to monitor for possible toxicity, the following tests should be performed before, during and after treatment.

1. Blood urea nitrogen and serum creatinine daily during therapy.
2. Complete blood and platelet counts daily during therapy.
3. Fasting blood glucose measurements should be taken before, daily during therapy, and at regular intervals after completion of therapy. Hyperglycaemia and diabetes mellitus, with or without preceding hypoglycaemia, have occurred up to several months after cessation of therapy.
4. Liver function tests (LFTs) including serum bilirubin, alkaline phosphatase, aspartate aminotransferase (AST/SGOT), and alanine aminotransferase (ALT/SGPT). If baseline measurements are normal and remain so during therapy, test weekly thereafter. When there is baseline elevation in LFTs or LFTs increase during therapy, continue monitoring weekly unless the patient is on other hepatotoxic agents, when monitoring every 3 to 5 days is appropriate.

5. Serum calcium, test weekly.
6. Urine analysis and serum electrolytes daily during therapy.
7. Electrocardiograms at regular intervals.

### **Use in renal impairment**

Severe renal impairment resulting in death may also occur in the presence of various clinical complications (e.g. bacterial sepsis), concurrent administration of other nephrotoxic antibiotic agents or previous evidence of renal disease.

### **Use in the elderly**

There is no information on the safe use of pentamidine in the elderly.

### **Paediatric use**

The safety and efficacy of Pentamidine Isetionate for Injection has not been established in the paediatric population, and pharmacokinetic data are extremely limited.

### **Effects on laboratory tests**

No data available.

## **4.5 Interactions with other medicines and other forms of interactions**

Since nephrotoxic effects may be additive, the concurrent or sequential use of pentamidine isetionate with drugs having a nephrotoxic potential (e.g. aminoglycosides, amphotericin B, cisplatin, methoxyflurane or vancomycin) should be undertaken with caution. Pentamidine Isetionate for Injection should be administered with caution to patients who are receiving drugs with hepatotoxic potential or medication that can impair the haematopoietic system.

## **4.6 Fertility, pregnancy and lactation**

### **Effects on fertility**

No data available.

### **Use in pregnancy – Category B3**

Pentamidine has been found to cross the placenta in rats given high doses late in pregnancy. Studies in rabbits have also shown pentamidine to be mildly embryotoxic, with an increase in post-implantation losses and delayed fetal ossification at doses of 1, 3 and 8 mg/kg of body weight.

It is not known whether pentamidine isetionate crosses the placenta or causes fetal harm when administered to pregnant women; therefore, pentamidine is contraindicated during pregnancy and should only be used when considered essential.

## **Use in lactation**

Since it is not known whether pentamidine isetionate is distributed into milk, the medicine is contraindicated during lactation and should only be administered to nursing mothers when considered essential.

## **4.7 Effects on ability to drive and use machines**

The effects of this medicine on a person's ability to drive and use machines were not assessed as part of its registration.

## **4.8 Adverse effects (undesirable effects)**

Fatalities due to severe hypotension, hypoglycemia, acute pancreatitis, renal impairment and cardiac arrhythmias have been reported in patients treated with pentamidine isetionate.

### **Life threatening reactions**

#### ***Blood and lymphatic system disorders***

Leucopenia (less than 1,000 cells per cubic millimetre), thrombocytopenia (less than 20,000 cells per cubic millimetre)

#### ***Immune system disorders***

Herxheimer reaction

#### ***Metabolism and nutrition disorders***

Severe hypoglycaemia, hypocalcaemia

#### ***Psychiatric disorders***

Toxic delirium

#### ***Nervous system disorders***

Syncope

#### ***Cardiac disorders***

Cardiac arrhythmias and cardiac arrest, ventricular tachycardia

#### ***Vascular disorders***

Severe hypotension

#### ***Gastrointestinal disorders***

Acute pancreatitis

#### ***Skin and subcutaneous tissue disorders***

Stevens-Johnson syndrome (single possible case)

### ***Renal and urinary disorders***

Acute renal failure

The above mentioned adverse effects can be severe, sometimes fatal, and require immediate corrective measures and withdrawal of treatment.

### **Other reactions**

Other adverse reactions reported are listed in this section per MedDRA system organ class.

### ***Blood and lymphatic system disorders***

Anaemia, thrombocytopenia, leucopenia

### ***Metabolism and nutrition disorders***

Hypocalcaemia, hypoglycaemia, hyperglycaemia, hyperkalaemia, hyponatraemia, diabetes mellitus

### ***Psychiatric disorders***

Hallucinations

### ***Nervous system disorders***

Taste disturbances, dizziness, syncope

### ***Cardiac disorders***

Tachycardia, bradycardia

### ***Vascular disorders***

Hypotension, facial flushing, venous thrombosis

### ***Respiratory, thoracic and mediastinal disorders***

Breathlessness

Local reactions ranging in severity from cough, breathlessness, wheezing, bronchospasms (with inhaled use), particularly in patients with a history of smoking or asthma, which can usually be controlled by prior use of bronchodilator.

### ***Gastrointestinal disorders***

Nausea, vomiting

### ***Hepatobiliary disorders***

Abnormal liver function (hepatic dysfunction)



### ***Skin and subcutaneous tissue disorders***

Abscess and/or necrosis, rash, itching, alopecia, erythema multiforme

### ***Renal and urinary disorders***

Azotaemia, albuminuria, glycosuria, increased creatinine levels

### ***General disorders and administration site conditions***

Local reactions at the injection site including abscess, pain, thrombophlebitis

### ***Investigations***

Depressed serum folate

### **Reporting suspected adverse effects**

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

## **4.9 Overdose**

There is no information available concerning the treatment of overdose. There is no specific antidote. In general, overdose would be expected to produce effects that are an extension of common adverse effects or of the serious metabolic sequelae observed. Treatment should be symptomatic and supportive. Neither peritoneal dialysis nor haemodialysis appear to remove the drug rapidly enough to cause a precipitous decline in the plasma concentration of pentamidine isetionate.

For information on the management of overdose, contact the Poisons Information Centre on 13 11 26 (Australia).

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

#### **Mechanism of action**

Pentamidine isetionate exhibits antiprotozoal activity against *Pneumocystis carinii*, *Leishmania* and some species of *Trypanosoma*.

The exact mechanism of antiprotozoal action of pentamidine has not been fully elucidated. Several mechanisms of action may be involved, and the role of the mechanism(s) may vary among the different types of protozoa (e.g., trypanosome, sporozoons). Most information on the antiprotozoal activity of pentamidine has been derived from studies involving trypanosomes. *In vitro* studies indicate that the drug interferes with nuclear metabolism.

#### **Microbiology**

Pentamidine isetionate exhibits antiprotozoal activity against *Pneumocystis carinii*, *Leishmania* and some species of *Trypanosoma*.

### **Clinical trials**

No data available.

## **5.2 Pharmacokinetic properties**

Limited information is available concerning the pharmacokinetics of pentamidine isetionate.

### **Absorption**

Following a single 4 mg/kg IV dose of pentamidine isetionate (given as a 2 hour infusion), peak plasma pentamidine concentrations averaged 612 nanogram/mL after completion of the IV infusion.

### **Distribution**

Distribution of pentamidine into human body tissues and fluids has not been well characterised, but the drug appears to be rapidly and extensively distributed and/or bound to tissues. Pentamidine has a distribution half life of 5 to 15 minutes after intravenous administration. Following parenteral administration, highest concentrations have been found in the liver, followed by the kidneys, adrenals, spleen, lungs and pancreas. Pentamidine penetrates the CNS only very poorly after prolonged therapy.

*In vitro*, pentamidine is reportedly 69% bound to serum proteins.

It is not known whether pentamidine isetionate crosses the placenta or is distributed into breast milk.

### **Excretion**

Little is known about the elimination in humans. Plasma concentrations of pentamidine have been found to decline in a biphasic manner following a single IV infusion in patients with normal renal function. The mean elimination half life was found to be 18 minutes in the initial phase and 6.4 hours in the terminal phase. Pentamidine appears to be eliminated very slowly from tissues in which the drug principally accumulates (e.g., liver, lungs). The half life of pentamidine may be prolonged in patients with impaired renal function, however no correlation between renal function and plasma clearance of pentamidine has been found. It is not known if the drug is excreted in faeces.

Only a small amount (approximately 6%) of the administered dose is excreted unchanged in the urine over a 15 day period.

Following a single 4 mg/kg IV dose of pentamidine isetionate in patients with AIDS or pneumocystis pneumonia who had normal renal function, about 2.5 to 5% of the dose was excreted in urine as unchanged drug over 24 hours, mainly within the first 8 hours after administration. Similar amounts (about 1 to 4% of the dose) were also excreted in urine as unchanged drug in 24 hours in patients with mild to moderate renal impairment.

Limited data suggest that pentamidine is not appreciably removed by haemodialysis or peritoneal dialysis.

### **5.3 Preclinical safety data**

#### **Genotoxicity**

No data available.

#### **Carcinogenicity**

No data available.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

None

### **6.2 Incompatibilities**

After reconstitution with Water for Injections, DBL™ Pentamidine Isethionate for Injection should not be mixed with any injection solution other than Glucose Intravenous Infusion 5% or Sodium Chloride Intravenous Infusion 0.9%.

### **6.3 Shelf life**

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

### **6.4 Special precautions for storage**

Store below 25°C.

### **6.5 Nature and contents of container**

DBL™ Pentamidine Isethionate for Injection is available in 1 vial.


<b>Strength</b>	<b>Pack Size</b>
300 mg/vial	1's

### **6.6 Special precautions for disposal**

In Australia, any unused medicine or waste material should be disposed of by taking to your local pharmacy.

## 6.7 Physicochemical properties

### Chemical structure



### CAS number

140-64-7

## 7. MEDICINE SCHEDULE (POISONS STANDARD)

S4 – Prescription Only Medicine

## 8. SPONSOR

Pfizer Australia Pty Ltd  
Level 17, 151 Clarence Street  
Sydney NSW 2000  
Toll Free Number: 1800 675 229  
www.pfizer.com.au

## 9. DATE OF FIRST APPROVAL

08 October 1991

## 10. DATE OF REVISION

07 January 2022

### Summary Table of Changes

Section changed	Summary of new information
4.4 and 4.8	Update to include information on bronchospam